



THE
POWER
TO KNOW.

SAS/ETS[®] 13.2 User's Guide The QLIM Procedure

This document is an individual chapter from *SAS/ETS® 13.2 User's Guide*.

The correct bibliographic citation for the complete manual is as follows: SAS Institute Inc. 2014. *SAS/ETS® 13.2 User's Guide*. Cary, NC: SAS Institute Inc.

Copyright © 2014, SAS Institute Inc., Cary, NC, USA

All rights reserved. Produced in the United States of America.

For a hard-copy book: No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without the prior written permission of the publisher, SAS Institute Inc.

For a Web download or e-book: Your use of this publication shall be governed by the terms established by the vendor at the time you acquire this publication.

The scanning, uploading, and distribution of this book via the Internet or any other means without the permission of the publisher is illegal and punishable by law. Please purchase only authorized electronic editions and do not participate in or encourage electronic piracy of copyrighted materials. Your support of others' rights is appreciated.

U.S. Government License Rights; Restricted Rights: The Software and its documentation is commercial computer software developed at private expense and is provided with RESTRICTED RIGHTS to the United States Government. Use, duplication or disclosure of the Software by the United States Government is subject to the license terms of this Agreement pursuant to, as applicable, FAR 12.212, DFAR 227.7202-1(a), DFAR 227.7202-3(a) and DFAR 227.7202-4 and, to the extent required under U.S. federal law, the minimum restricted rights as set out in FAR 52.227-19 (DEC 2007). If FAR 52.227-19 is applicable, this provision serves as notice under clause (c) thereof and no other notice is required to be affixed to the Software or documentation. The Government's rights in Software and documentation shall be only those set forth in this Agreement.

SAS Institute Inc., SAS Campus Drive, Cary, North Carolina 27513.

August 2014

SAS provides a complete selection of books and electronic products to help customers use SAS® software to its fullest potential. For more information about our offerings, visit support.sas.com/bookstore or call 1-800-727-3228.

SAS® and all other SAS Institute Inc. product or service names are registered trademarks or trademarks of SAS Institute Inc. in the USA and other countries. ® indicates USA registration.

Other brand and product names are trademarks of their respective companies.



Gain Greater Insight into Your SAS[®] Software with SAS Books.

Discover all that you need on your journey to knowledge and empowerment.

 support.sas.com/bookstore
for additional books and resources.


THE POWER TO KNOW.®

SAS and all other SAS Institute Inc. product or service names are registered trademarks or trademarks of SAS Institute Inc. in the USA and other countries. ® indicates USA registration. Other brand and product names are trademarks of their respective companies. © 2013 SAS Institute Inc. All rights reserved. S107969US.0613

Chapter 22

The QLIM Procedure

Contents

Overview: QLIM Procedure	1520
Getting Started: QLIM Procedure	1521
Introductory Example: Binary Probit and Logit Models	1523
Syntax: QLIM Procedure	1527
Functional Summary	1528
PROC QLIM Statement	1531
BAYES Statement	1536
BOUNDS Statement	1541
BY Statement	1542
CLASS Statement	1542
ENDOGENOUS Statement	1543
FREQ Statement	1546
HETERO Statement	1546
INIT Statement	1547
MODEL Statement	1548
NLOPTIONS Statement	1549
OUTPUT Statement	1550
PRIOR Statement	1551
RESTRICT Statement	1551
TEST Statement	1552
WEIGHT Statement	1554
Details: QLIM Procedure	1554
Ordinal Discrete Choice Modeling	1554
Limited Dependent Variable Models	1557
Stochastic Frontier Production and Cost Models	1561
Heteroscedasticity and Box-Cox Transformation	1563
Bivariate Limited Dependent Variable Modeling	1564
Selection Models	1565
Multivariate Limited Dependent Models	1569
Variable Selection	1569
Tests on Parameters	1569
Endogeneity and Instrumental Variables	1570
Bayesian Analysis	1577
Prior Distributions	1584
Automated MCMC	1587
Output to SAS Data Set	1590

OUTEST= Data Set	1595
Naming	1595
ODS Table Names	1597
ODS Graphics	1598
Examples: QLIM Procedure	1600
Example 22.1: Ordered Data Modeling	1600
Example 22.2: Tobit Analysis	1603
Example 22.3: Bivariate Probit Analysis	1605
Example 22.4: Sample Selection Model	1606
Example 22.5: Sample Selection Model with Truncation and Censoring	1607
Example 22.6: Types of Tobit Models	1610
Example 22.7: Stochastic Frontier Models	1616
Example 22.8: Bayesian Modeling	1621
References	1628

Overview: QLIM Procedure

The QLIM (qualitative and limited dependent variable model) procedure analyzes univariate and multivariate limited dependent variable models in which dependent variables take discrete values or in which dependent variables are observed only in a limited range of values. These models include logit, probit, tobit, selection, and multivariate models. The multivariate model can contain discrete choice and limited endogenous variables in addition to continuous endogenous variables.

The QLIM procedure supports the following models:

- linear regression model with heteroscedasticity
- Box-Cox regression with heteroscedasticity
- probit with heteroscedasticity
- logit with heteroscedasticity
- tobit (censored and truncated) with heteroscedasticity
- bivariate probit
- bivariate tobit
- sample selection and switching regression models
- multivariate limited dependent variables
- stochastic frontier production and cost models

In the linear regression models with heteroscedasticity, the assumption that error variance is constant across observations is relaxed. The QLIM procedure allows for a number of different linear and nonlinear variance specifications. Another way to make the linear model more appropriate to fit the data and reduce skewness is to apply Box-Cox transformation. If the nature of the data is such that the dependent variable is discrete and it takes only two possible values, ordinary least squares (OLS) estimates are inconsistent. The QLIM procedure offers probit and logit models to overcome these estimation problems. Assumptions about the error variance can also be relaxed in order to estimate probit or logit with heteroscedasticity.

The QLIM procedure also offers a class of models in which the dependent variable is censored or truncated from below or above or both. When a continuous dependent variable is observed only within a certain range and values outside this range are not available, the QLIM procedure offers a class of models that adjust for truncation. In some cases, the dependent variable is continuous only in a certain range and all values outside this range are reported as being on its boundary. For example, if it is not possible to observe negative values, the value of the dependent variable is reported as equal to 0. Because the data are censored, OLS results are inconsistent, and it cannot be guaranteed that the predicted values from the model fall in the appropriate region.

Most of the models in the QLIM procedure can be extended to accommodate bivariate and multivariate scenarios. The assumption that one variable is observed only if another variable takes on certain values lead to the introduction of sample selection models. If the dependent variables are mutually exclusive and observed only for certain ranges of the selection variable, the sample selection can be extended to include cases of switching regression. Stochastic frontier production and cost models allow for random shocks of the production or cost. They include a systematic positive component in the error term that adjusts for technological or cost inefficiency.

The QLIM procedure can use the maximum likelihood method for both univariate and multivariate models. PROC QLIM can use the Bayesian method for univariate models and for multivariate models in which the likelihood function is available in closed form. Initial starting values for the nonlinear optimizations are typically calculated by OLS.

Getting Started: QLIM Procedure

The QLIM procedure is similar in use to the other regression or simultaneous equations model procedures in the SAS System. For example, the following statements are used to estimate a binary choice model by using the probit probability function:

```
proc qlim data=a;
  model y = x1;
  endogenous y ~ discrete;
run;
```

The response variable, *y*, is numeric and has discrete values. PROC QLIM enables the user to specify the type of endogenous variables in the ENDOGENOUS statement. The binary probit model can be also specified as follows:

```
model y = x1 / discrete;
```

When multiple endogenous variables are specified in the QLIM procedure, these equations are estimated as a system. Multiple endogenous variables can be specified with one MODEL statement in the QLIM procedure when these models have the same exogenous variables:

```
model y1 y2 = x1 x2 / discrete;
```

The preceding specification is equivalent to the following statements:

```
proc qlim data=a;
  model y1 = x1 x2;
  model y2 = x1 x2;
  endogenous y1 y2 ~ discrete;
run;
```

Some equations in multivariate models can be continuous while other equations can be discrete. A bivariate model with a discrete and a continuous equation is specified as follows:

```
proc qlim data=a;
  model y1 = x1 x2;
  model y2 = x3 x4;
  endogenous y1 ~ discrete;
run;
```

The standard tobit model is estimated by specifying the endogenous variable to be truncated or censored. The limits of the dependent variable can be specified with the CENSORED or TRUNCATED option in the ENDOGENOUS or MODEL statement when the data are limited by specific values or variables. For example, the two-limit censored model requires two variables that contain the lower (bottom) and upper (top) bound:

```
proc qlim data=a;
  model y = x1 x2 x3;
  endogenous y ~ censored(lb=bottom ub=top);
run;
```

The bounds can be numbers if they are fixed for all observations in the data set. For example, the standard tobit model can be specified as follows:

```
proc qlim data=a;
  model y = x1 x2 x3;
  endogenous y ~ censored(lb=0);
run;
```


Introductory Example: Binary Probit and Logit Models

The following example illustrates the use of PROC QLIM. The data were originally published by Mroz (1987) and downloaded from Wooldridge (2002). This data set is based on a sample of 753 married white women. The dependent variable is a discrete variable of labor force participation (`inlf`). Explanatory variables are the number of children ages 5 or younger (`kidslt6`), the number of children ages 6 to 18 (`kidsge6`), the woman's age (`age`), the woman's years of schooling (`educ`), wife's labor experience (`exper`), square of experience (`expersq`), and the family income excluding the wife's wage (`nwifeinc`). The program (with data values omitted) is as follows:

```
/*-- Binary Probit --*/
proc qlim data=mroz plots=predicted;
  model inlf = nwifeinc educ exper expersq
              age kidslt6 kidsge6 / discrete;
run;
```

Results of this analysis are shown in the following four figures. In the first table, shown in [Figure 22.1](#), PROC QLIM provides frequency information about each choice. In this example, 428 women participate in the labor force (`inlf`=1).

Figure 22.1 Choice Frequency Summary

Binary Data

The QLIM Procedure

Discrete Response Profile of <code>inlf</code>		
Index	Value	Total Frequency
1	0	325
2	1	428

The second table is the estimation summary table shown in [Figure 22.2](#). Included are the number of dependent variables, names of dependent variables, the number of observations, the log-likelihood function value, the maximum absolute gradient, the number of iterations, AIC, and Schwarz criterion.

Figure 22.2 Fit Summary Table of Binary Probit

Model Fit Summary	
Number of Endogenous Variables	1
Endogenous Variable	inlf
Number of Observations	753
Log Likelihood	-401.30219
Maximum Absolute Gradient	0.0000669
Number of Iterations	15
Optimization Method	Quasi-Newton
AIC	818.60439
Schwarz Criterion	855.59691

Goodness-of-fit measures are displayed in Figure 22.3. All measures except McKelvey-Zavoina's definition are based on the log-likelihood function value. The likelihood ratio test statistic has chi-square distribution conditional on the null hypothesis that all slope coefficients are zero. In this example, the likelihood ratio statistic is used to test the hypothesis that $\text{kidslt6} = \text{kidsge6} = \text{age} = \text{educ} = \text{exper} = \text{expersq} = \text{nwifeinc} = 0$.

Figure 22.3 Goodness of Fit

Goodness-of-Fit Measures		
Measure	Value	Formula
Likelihood Ratio (R)	227.14	$2 * (\text{LogL} - \text{LogL0})$
Upper Bound of R (U)	1029.7	$2 * \text{LogL0}$
Aldrich-Nelson	0.2317	$R / (R+N)$
Cragg-Uhler 1	0.2604	$1 - \exp(-R/N)$
Cragg-Uhler 2	0.3494	$(1 - \exp(-R/N)) / (1 - \exp(-U/N))$
Estrella	0.2888	$1 - (1 - R/U)^{(U/N)}$
Adjusted Estrella	0.2693	$1 - ((\text{LogL} - K) / \text{LogL0})^{(-2/N * \text{LogL0})}$
McFadden's LRI	0.2206	R / U
Veall-Zimmermann	0.4012	$(R * (U+N)) / (U * (R+N))$
McKelvey-Zavoina	0.4025	

N = # of observations, K = # of regressors


The parameter estimates and standard errors are shown in Figure 22.4.

Figure 22.4 Parameter Estimates of Binary Probit

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
Intercept	1	0.270077	0.508590	0.53	0.5954
nwifeinc	1	-0.012024	0.004840	-2.48	0.0130
educ	1	0.130905	0.025255	5.18	<.0001
exper	1	0.123348	0.018720	6.59	<.0001
expersq	1	-0.001887	0.000600	-3.14	0.0017
age	1	-0.052853	0.008477	-6.24	<.0001
kidslt6	1	-0.868329	0.118519	-7.33	<.0001
kidsge6	1	0.036005	0.043477	0.83	0.4076

Finally, the QLIM procedure profiles the predicted outcome with respect to the regressors. For example, [Output 22.5](#) shows the predicted values profiled with respect to `nwifeinc`, `educ`, `exper`, `expersq`, `age`, and `kidslt6`.

Figure 22.5 Predictions by Regressors: `nwifeinc`, `educ`, `exper`, `expersq`, `age`, and `kidslt6`



When the error term has a logistic distribution, the binary logit model is estimated. To specify a logistic distribution, add `D=LOGIT` option as follows:

```
/*-- Binary Logit --*/
proc qlim data=mroz;
  model inlf = nwifeinc educ exper expersq
              age kidslt6 kidsge6 / discrete(d=logit);
run;
```

The estimated parameters are shown in [Figure 22.6](#).

Figure 22.6 Parameter Estimates of Binary Logit
Binary Data**The QLIM Procedure**

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
Intercept	1	0.425452	0.860365	0.49	0.6210
nwifeinc	1	-0.021345	0.008421	-2.53	0.0113
educ	1	0.221170	0.043441	5.09	<.0001
exper	1	0.205870	0.032070	6.42	<.0001
expersq	1	-0.003154	0.001017	-3.10	0.0019
age	1	-0.088024	0.014572	-6.04	<.0001
kidslt6	1	-1.443354	0.203575	-7.09	<.0001
kidsge6	1	0.060112	0.074791	0.80	0.4215

The heteroscedastic logit model can be estimated using the HETERO statement. If the variance of the logit model is a function of the family income level excluding wife's income (nwifeinc), the variance can be specified as

$$\text{Var}(\epsilon_i) = \sigma^2 \exp(\gamma * \text{nwifeinc}_i)$$

where σ^2 is normalized to 1 because the dependent variable is discrete. The following SAS statements estimate the heteroscedastic logit model:

```

/*-- Binary Logit with Heteroscedasticity --*/
proc qlim data=mroz;
  model inlf = nwifeinc educ exper expersq
              age kidslt6 kidsge6 / discrete(d=logit);
  hetero inlf ~ nwifeinc / noconst;
run;

```

The parameter estimate, γ , of the heteroscedasticity variable is listed as _H.nwifeinc; see [Figure 22.7](#).

Figure 22.7 Parameter Estimates of Binary Logit with Heteroscedasticity

Binary Data					
The QLIM Procedure					
Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
Intercept	1	0.510445	0.983538	0.52	0.6038
nwifeinc	1	-0.026778	0.012108	-2.21	0.0270
educ	1	0.255547	0.061728	4.14	<.0001
exper	1	0.234105	0.046639	5.02	<.0001
expersq	1	-0.003613	0.001236	-2.92	0.0035
age	1	-0.100878	0.021491	-4.69	<.0001
kidslt6	1	-1.645206	0.311296	-5.29	<.0001
kidsge6	1	0.066941	0.085633	0.78	0.4344
_H.nwifeinc	1	0.013280	0.013606	0.98	0.3291

Syntax: QLIM Procedure

The QLIM procedure is controlled by the following statements:

```

PROC QLIM options ;
  BAYES <options> ;
  BOUNDS bound1 < , bound2 ... > ;
  BY variables ;
  CLASS variables ;
  FREQ variable ;
  ENDOGENOUS variables ~ options ;
  HETERO dependent variables ~ exogenous variables / options ;
  INIT initvalue1 < , initvalue2 ... > ;
  MODEL dependent variables = regressors / options ;
  NLOPTIONS options ;
  OUTPUT options ;
  PRIOR variables ~ distributions ;
  RESTRICT restriction1 < , restriction2 ... > ;
  TEST options ;
  WEIGHT variable ;

```

At least one MODEL statement is required. If more than one MODEL statement is used, the QLIM procedure estimates a system of models. If a FREQ or WEIGHT statement is specified more than once, the variable specified in the first instance is used. Main effects and higher-order terms can be specified in the MODEL statement, as in the GLM procedure and PROBIT procedure in SAS/STAT. If a CLASS statement is used, it must precede the MODEL statement.

Functional Summary

Table 22.1 summarizes the statements and options used with the QLIM procedure.

Table 22.1 PROC QLIM Functional Summary

Description	Statement	Option
Data Set Options		
Specifies the input data set	QLIM	DATA=
Writes parameter estimates to an output data set	QLIM	OUTEST=
Writes predictions to an output data set	OUTPUT	OUT=
Declaring the Role of Variables		
Specifies BY-group processing	BY	
Specifies classification variables	CLASS	
Specifies a frequency variable	FREQ	
Specifies a weight variable	WEIGHT	NONNORMALIZE
Printing Control Options		
Requests all printing options	QLIM	PRINTALL
Prints correlation matrix of the estimates	QLIM	CORRB
Prints covariance matrix of the estimates	QLIM	COVB
Prints a summary iteration listing	QLIM	ITPRINT
Suppresses the normal printed output	QLIM	NOPRINT
Plotting Options		
Displays plots	QLIM	PLOTS=
Options to Control the Optimization Process		
Specifies the optimization method	QLIM	METHOD=
Specifies the optimization options	NLOPTIONS	see Chapter 6, “Nonlinear Optimization Methods,”
Sets initial values for parameters	INIT	
Specifies upper and lower bounds for the parameter estimates	BOUNDS	
Specifies linear restrictions on the parameter estimates	RESTRICT	
Model Estimation Options		
Specifies options specific to Box-Cox transformation	MODEL	BOXCOX()
Suppresses the intercept parameter	MODEL	NOINT
Specifies variable selection	MODEL	SELECTVAR=()
Specifies the type of random number generators	MODEL	RANDNUM=
Specifies that initial values are generated using random numbers	MODEL	RANDOMINIT

Table 22.1 *continued*

Description	Statement	Option
Specifies a seed for pseudo-random number generation	QLIM	SEED=
Specifies the number of draws for Monte Carlo integration	QLIM	NDRAW=
Specifies the method to calculate parameter covariance	QLIM	COVEST=
Requests estimation by Heckman's two-step method	QLIM	HECKIT
Bayesian MCMC Options		
Controls the aggregation of multiple posterior chains	BAYES	AGGREGATION=
Automates the initialization of the MCMC algorithm	BAYES	AUTOMCMC()
Specifies the initial values of the MCMC	INIT	
Specifies the maximum number of tuning phases	BAYES	MAXTUNE=
Specifies the minimum number of tuning phases	BAYES	MINTUNE=
Specifies the number of burn-in iterations	BAYES	NBI=
Specifies the number of iterations during the sampling phase	BAYES	NMC=
Specifies the number of samples for the prior predictive analysis	BAYES	NMCPRIOR=
Specifies the number of threads to use during the sampling phase	BAYES	NTRDS=
Specifies the number of iterations during the tuning phase	BAYES	NTU=
Controls options for constructing the initial proposal covariance matrix	BAYES	PROPCOV=
Specifies the sampling scheme	BAYES	SAMPLING=
Specifies the random number generator seed	BAYES	SEED=
Prints the time required for the MCMC sampling	BAYES	SIMTIME
Controls the thinning of the Markov chain	BAYES	THIN=
Bayesian Summary Statistics and Convergence Diagnostics		
Displays convergence diagnostics	BAYES	DIAGNOSTICS=
Displays summary statistics of the posterior samples	BAYES	STATISTICS=
Bayesian Prior and Posterior Samples		
Specifies a SAS data set for the posterior samples	BAYES	OUTPOST=
Specifies a SAS data set for the prior samples	BAYES	OUTPRIOR=
Bayesian Analysis		
Specifies normal prior distribution	PRIOR	NORMAL(MEAN=, VAR=)
Specifies gamma prior distribution	PRIOR	GAMMA(SHAPE=, SCALE=)
Specifies inverse gamma prior distribution	PRIOR	IGAMMA(SHAPE=, SCALE=)
Specifies uniform prior distribution	PRIOR	UNIFORM(MIN=, MAX=)
Specifies beta prior distribution	PRIOR	BETA(SHAPE1=, SHAPE2=, MIN=, MAX=)

Table 22.1 *continued*

Description	Statement	Option
Specifies t prior distribution	PRIOR	T(LOCATION=, DF=)
Endogenous Variable Options		
Specifies discrete variable	ENDOGENOUS	DISCRETE()
Specifies censored variable	ENDOGENOUS	CENSORED()
Specifies truncated variable	ENDOGENOUS	TRUNCATED()
Specifies variable selection condition	ENDOGENOUS	SELECT()
Specifies stochastic frontier variable	ENDOGENOUS	FRONTIER()
Endogeneity and Overidentification Test Options		
Requests the variable addition test for endogeneity	ENDOGENOUS	ENDOTEST()
Requests the overidentification test	ENDOGENOUS	OVERID()
Heteroscedasticity Model Options		
Specifies the function for heteroscedasticity models	HETERO	LINK=
Squares the function for heteroscedasticity models	HETERO	SQUARE
Specifies no constant for heteroscedasticity models	HETERO	NOCONST
Output Control Options		
Outputs predicted values	OUTPUT	PREDICTED
Outputs structured part	OUTPUT	XBETA
Outputs residuals	OUTPUT	RESIDUAL
Outputs error standard deviation	OUTPUT	ERRSTD
Outputs marginal effects	OUTPUT	MARGINAL
Outputs probability for the current response	OUTPUT	PROB
Outputs probability for all responses	OUTPUT	PROBALL
Outputs expected value	OUTPUT	EXPECTED
Outputs conditional expected value	OUTPUT	CONDITIONAL
Outputs inverse Mills ratio	OUTPUT	MILLS
Outputs technical efficiency measures	OUTPUT	TE1
	OUTPUT	TE2
Includes covariances in the OUTEST= data set	QLIM	COVOUT
Includes correlations in the OUTEST= data set	QLIM	CORROUT
Test Request Options		
Requests Wald, Lagrange multiplier, and likelihood ratio tests	TEST	ALL
Requests the WALD test	TEST	WALD
Requests the Lagrange multiplier test	TEST	LM
Requests the likelihood ratio test	TEST	LR

PROC QLIM Statement

PROC QLIM *options* ;

The following options can be used in the PROC QLIM statement.

Data Set Options

DATA=SAS-data-set

specifies the input SAS data set. If the DATA= option is not specified, PROC QLIM uses the most recently created SAS data set.

Output Data Set Options

OUTEST=SAS-data-set

writes the parameter estimates to an output data set.

COVOUT

writes the covariance matrix for the parameter estimates to the OUTEST= data set. This option is valid only if the OUTEST= option is specified.

CORROUT

writes the correlation matrix for the parameter estimates to the OUTEST= data set. This option is valid only if the OUTEST= option is specified.

Printing Options

NOPRINT

suppresses the normal printed output but does not suppress error listings. If NOPRINT option is set, then any other print option is turned off.

PRINTALL

turns on all the printing-control options. The options set by PRINTALL are COVB and CORRB.

CORRB

prints the correlation matrix of the parameter estimates.

COVB

prints the covariance matrix of the parameter estimates.

ITPRINT

prints the initial parameter estimates, convergence criteria, and all constraints of the optimization. At each iteration, objective function value, step size, maximum gradient, and slope of search direction are printed as well.

Model Estimation Options

COVEST=covariance-option

specifies the method to calculate the covariance matrix of parameter estimates. The supported covariance types are as follows:

OP	specifies the covariance from the outer product matrix.
HESSIAN	specifies the covariance from the inverse Hessian matrix.
QML	specifies the covariance from the outer product and Hessian matrices (the quasi-maximum likelihood estimates).

The default is COVEST=HESSIAN.

HECKIT <(heckit-options)>

requests that the selection model be estimated by Heckman's two-step estimation method. You must specify exactly two MODEL statements when you use the HECKIT option. One of the models must be a binary probit model; therefore, you must specify the DISCRETE option in the MODEL or ENDOGENOUS statement. You base the selection on the binary probit model for the second model; therefore, you must specify the SELECT option for this model.

You can specify one or both of the following *heckit-options*:

SECONDSTAGE=OLS | ML

specifies the estimation method of the second stage of Heckman's two-step method.

SECONDSTAGE=OLS requests the ordinary least squares method for the second stage. If you specify SECONDSTAGE=OLS, then the model of interest—that is, the model that uses the SELECT option—must be linear and contain a continuous dependent variable. Therefore, you cannot specify the DISCRETE, CENSORED, or TRUNCATED option along with the SELECT option for the model of interest. When you specify the SECONDSTAGE=OLS option, you cannot test or restrict the parameters of the model of interest. However, you can test or restrict the parameters of the selection model—that is, the model that defines the selection rule.

If you specify SECONDSTAGE=ML, then PROC QLIM uses the maximum likelihood method in the second stage, as it does in the first stage. When you specify SECONDSTAGE=ML, the model of interest can be nonlinear. Moreover, you can also use the TEST or RESTRICT statement to test or restrict the parameters of the model of interest.

By default, SECONDSTAGE=OLS.

UNCORRECTED

requests the conventional OLS standard errors when the second-stage estimation method is the ordinary least squares method. If you do not specify the UNCORRECTED option, PROC QLIM reports the corrected OLS standard errors. For more information about the corrected standard errors, see the section “[Heckman's Two-Step Selection Method](#)” on page 1567.

If you specify both the UNCORRECTED and SECONDSTAGE=ML options, PROC QLIM ignores the UNCORRECTED option, because the UNCORRECTED option is related to the OLS standard errors.

NDRAW=*value*

specifies the number of draws for Monte Carlo integration.

SEED=*value*

specifies a seed for pseudo-random number generation in Monte Carlo integration.

Optimization Process Control Options

PROC QLIM uses the nonlinear optimization (NLO) subsystem to perform nonlinear optimization tasks. You can use any of the NLO options in the NLOPTIONS statement. For more information, see Chapter 6, “Nonlinear Optimization Methods.”

METHOD=*value*

specifies the optimization method. If this option is specified, it overwrites the TECH= option in NLOPTIONS statement. Valid values are as follows:

CONGRA	performs a conjugate-gradient optimization
DBLDOG	performs a version of double-dogleg optimization
NMSIMP	performs a Nelder-Mead simplex optimization
NEWRAP	performs a Newton-Raphson optimization combining a line-search algorithm with ridging
NRRIDG	performs a Newton-Raphson optimization with ridging
QUANew	performs a quasi-Newton optimization
TRUREG	performs a trust region optimization

The default method is METHOD=QUANew.

Plotting Options

PLOTS< (*global-plot-options*) > = *plot-request* | (*plot-requests*)

controls the display of plots. By default, the plots are displayed in panels unless the UNPACK *global-plot-option* is specified. When you specify only one *plot-request*, you can omit the parentheses around the *plot-request*.

Global Plot Options

You can specify the following *global-plot-options*:

ONLY

displays only the requested plot.

PRIOR

displays the prior predictive graph that is associated with the requested posterior predictive plot BAYESPRED. This option is available only for Bayesian analysis.

UNPACKPANEL**UNPACK**

specifies that all paneled plots be unpacked, meaning that each plot in a panel is displayed separately.

Plot Requests

You can specify the following *plot-requests*:

ALL

specifies all types of available plots.

AUTOCORR<(LAGS=*n*)>

displays the autocorrelation function plots for the parameters. This *plot-request* is available only for Bayesian analysis. The optional LAGS= suboption the number (up to lag *n*) of autocorrelations to be plotted in the AUTOCORR plot. If this suboption is not specified, autocorrelations are plotted up to lag 50.

BAYESDIAG

displays the TRACE, AUTOCORR, and DENSITY plots. This *plot-request* is available only for Bayesian analysis.

BAYESPRED

displays the predictive analysis. The predictive analysis takes into account the variability of the error term, whereas the PREDICTED *plot-request* does not. The BAYESPRED *plot-request* is available only for Bayesian analysis.

BAYESSUM

displays the posterior distribution, the prior distribution, and the maximum likelihood estimates. This *plot-request* is available only for Bayesian analysis.

CONDITIONAL

displays the conditional expected values for continuous endogenous variables. Each contributing regressor is set equal to its mean, except for the parameter that is reported on the X axis. This *plot-request* is not available for Bayesian analysis.

DENSITY<(FRINGE)>

displays the kernel density plots for the parameters. This *plot-request* is available only for Bayesian analysis. If you specify the FRINGE suboption, a fringe plot is created on the X axis of the kernel density plot. This *plot-request* is available only for Bayesian analysis.

ERRSTD

displays the error standard deviation versus observed regressors when you also specify a HETERO statement. This *plot-request* is not available for Bayesian analysis.

EXPECTED

displays the expected values for continuous endogenous variables. Each contributing regressor is set equal to its mean, except for the parameter that is reported on the X axis. This *plot-request* is not available for Bayesian analysis.

MARGINAL

displays the marginal effects. Each contributing regressor is set equal to its mean, except for the parameter that is reported on the X axis. This *plot-request* is not available for Bayesian analysis.

MILLS

displays the inverse Mills ratio. Each contributing regressor is set equal to its mean, except for the parameter that is reported on the X axis. This *plot-request* is not available for Bayesian analysis.

NONE

suppresses all diagnostic plots.

PREDICTED

displays the model predicted values. Each contributing regressor is set equal to its mean, except for the parameter that is reported on the X axis. This *plot-request* is not available for Bayesian analysis.

PROB

displays the predicted response probability. Each contributing regressor is set equal to its mean, except for the parameter that is reported on the X axis. This *plot-request* is not available for Bayesian analysis.

PROBALL

displays the predicted probabilities for each level of the response. Each contributing regressor is set equal to its mean, except for the parameter that is reported on the X axis. This *plot-request* is not available for Bayesian analysis.

PROFLIK

displays the profiled log likelihood. Each profiled graph is obtained by setting all the parameters to their maximum likelihood estimate except for the profiling parameter. The profiling parameter takes values on a predefined grid that is determined by the maximum likelihood estimate of the corresponding standard deviation.

RESIDUAL

displays the residuals versus observed regressors. This *plot-request* is not available for Bayesian analysis.

TE1

displays the technical efficiency for the stochastic frontier model as suggested by Battese and Coelli (1988). Each contributing regressor is set equal to its mean, except for the parameter that is reported on the X axis. This *plot-request* is not available for Bayesian analysis.

TE2

displays the technical efficiency for the stochastic frontier model as suggested by Jondrow et al. (1982). Each contributing regressor is set equal to its mean, except for the parameter that is reported on the X axis. This *plot-request* is not available for Bayesian analysis.

TRACE<(SMOOTH)>

displays the trace plots for the parameters. This *plot-request* is available only for Bayesian analysis. The SMOOTH suboption displays a fitted penalized B-spline curve for each TRACE plot.

XBETA

displays the structural part on the right-hand side of the model. Each contributing regressor is set equal to its mean, except for the parameter that is reported on the X axis. This is not available for Bayesian analysis.

BAYES Statement

BAYES <options> ;

The BAYES statement controls the Metropolis sampling scheme that is used to obtain samples from the posterior distribution of the underlying model and data.

AGGREGATION=WEIGHTED | NOWEIGHTED (Experimental)

specifies how multiple posterior samples should be aggregated. AGGREGATION=WEIGHTED implements a weighted resampling scheme for the aggregation of multiple posterior chains. You can use this option when the posterior distribution is characterized by several very distinct posterior modes. AGGREGATION=NOWEIGHTED aggregates multiple posterior chains without any adjustment. You can use this option when the posterior distribution is characterized by one or few relatively close posterior modes. By default, AGGREGATION=NOWEIGHTED. For more information, see the section “[Aggregation of Multiple Chains](#)” on page 1579.

AUTOMCMC <=(automcmc-options)> (Experimental)

specifies an algorithm for the auto-initialization of the MCMC sampling algorithm. For more information, see the section “[Automated Initialization of MCMC](#)” on page 1580.

ACCURACY= (accuracy-options)

customizes the behavior of the AUTOMCMC algorithm when you are searching for an accurate representation of the posterior distribution. You can specify the following *accuracy-options*:

ATTEMPTS=number

specifies the maximum number of attempts that is required in order to obtain accurate samples from the posterior distribution. By default, ATTEMPTS=10.

TARGETESS=number

requests that the accuracy search be based on the effective sample size (ESS) analysis. If you specify this option, you must also specify the minimum *number* of effective samples.

TARGETSTATS<=(targetstats-option)>

requests that the accuracy search be based on the analysis of the posterior mean and a posterior quantile of interest. You can customize the behavior of the analysis of the posterior mean by adjusting HEIDELBERGER sub-options. You can customize the behavior of the analysis of the posterior quantile by adjusting the RAFTERY sub-options. If you specify TARGETSTATS,, you can also specify how the Raftery-Lewis test should be interpreted by using the following option:

RLLIMITS=(LB=number UB=number)

specifies a region where the search for the optimal sample size depends directly on the Raftery-Lewis test. By default, RLLIMITS (LB=10000 UB=300000).

TOL=*value*

specifies the proportion of parameters that are required to be accurate. By default, TOL=0.95.

MAXNMC=*number*

specifies the maximum number of posterior samples that the AUTOMCMC option allows. By default, MAXNMC=700000.

RANDINIT <=(*randinit-options*)>

specifies random starting points for the MCMC algorithm. The starting points can be sampled around the maximum likelihood estimate and around the prior mean. You can specify the following *randinit-options*:

MULTIPLIER= (*value*)

specifies the radius of the area where the starting points are sampled. For the starting points that are sampled around the maximum likelihood estimate, the radius equals the standard deviation of the maximum likelihood estimate multiplied by the multiplier value. For the starting points that are sampled around the prior mean, the radius equals the standard deviation of the prior distribution multiplied by the multiplier value. By default, MULTIPLIER=2.

PROPORTION= (*value*)

specifies the proportion of starting points that are sampled around the maximum likelihood estimate and around the prior mean. By default, PROPORTION=0, which implies that all the initial points are sampled around the maximum likelihood estimate. If you use choose to sample starting points around the prior mean, the convergence of the MCMC algorithm could be very slow.

STATIONARITY= (*stationarity-options*)

customizes the behavior of the AUTOMCMC algorithm when you are trying to sample from the posterior distribution. You can specify the following *accuracy-options*:

ATTEMPTS=*number*

specifies the maximum number of attempts that are required in order to obtain stationary samples from the posterior distribution. By default, ATTEMPTS=10.

TOL=*value*

specifies the proportion of parameter whose samples must to be stationary. By default, TOL=0.95.

DIAGNOSTICS=ALL | NONE | (*keyword-list*)**DIAG=**ALL | NONE | (*keyword-list*)

controls which diagnostics are produced. All the following diagnostics are produced with DIAGNOSTICS=ALL. If you do not want any of these diagnostics, specify DIAGNOSTICS=NONE. If you want some but not all of the diagnostics, or if you want to change certain settings of these diagnostics, specify a subset of the following keywords. By default, DIAGNOSTICS=NONE.

AUTOCORR <(LAGS= *numeric-list*)>

computes the autocorrelations at lags that are specified in the *numeric-list*. Elements in the *numeric-list* are truncated to integers, and repeated values are removed. If the LAGS= option is not specified, autocorrelations of lags 1, 5, 10, and are computed.

ESS

computes Carlin's estimate of the effective sample size, the correlation time, and the efficiency of the chain for each parameter.

GEWEKE <(geweke-options)>

computes the Geweke spectral density diagnostics, which are essentially a two-sample t test between the first f_1 portion and the last f_2 portion of the chain. The default is $f_1 = 0.1$ and $f_2 = 0.5$, but you can choose other fractions by using the following *geweke-options*:

FRAC1=value

specifies the fraction f_1 for the first window.

FRAC2=value

specifies the fraction f_2 for the second window.

HEIDELBERGER <(heidel-options)>

computes the Heidelberg and Welch diagnostic for each variable, which consists of a stationarity test of the null hypothesis that the sample values form a stationary process. If the stationarity test is not rejected, a halfwidth test is then carried out. Optionally, you can specify one or more of the following *heidel-options*:

SALPHA=value

specifies the α level ($0 < \alpha < 1$) for the stationarity test.

HALPHA=value

specifies the α level ($0 < \alpha < 1$) for the halfwidth test.

EPS=value

specifies a positive number ϵ such that if the halfwidth is less than ϵ times the sample mean of the retained iterates, the halfwidth test is passed.

MCSE**MCERROR**

computes the Monte Carlo standard error for each parameter. The Monte Carlo standard error, which measures the simulation accuracy, is the standard error of the posterior mean estimate and is calculated as the posterior standard deviation divided by the square root of the effective sample size.

RAFTERY<(raftery-options)>

computes the Raftery and Lewis diagnostics, which evaluate the accuracy of the estimated quantile ($\hat{\theta}_Q$ for a given $Q \in (0, 1)$) of a chain. $\hat{\theta}_Q$ can achieve any degree of accuracy when the chain is allowed to run for a long time. The computation is stopped when the estimated probability $\hat{P}_Q = \Pr(\theta \leq \hat{\theta}_Q)$ reaches within $\pm R$ of the value Q with probability S ; that is, $\Pr(Q - R \leq \hat{P}_Q \leq Q + R) = S$. The following *raftery-options* enable you to specify Q , R , S , and a precision level ϵ for the test:

QUANTILE | Q=value

specifies the order (a value between 0 and 1) of the quantile of interest. The default is 0.025.

ACCURACY | R=value

specifies a small positive number as the margin of error for measuring the accuracy of estimation of the quantile. The default is 0.005.

PROBABILITY | S=value

specifies the probability of attaining the accuracy of the estimation of the quantile. The default is 0.95.

EPSILON | EPS=value

specifies the tolerance level (a small positive number) for the stationary test. The default is 0.001.

MINTUNE=number

specifies the minimum number of tuning phases. The default is 2.

MAXTUNE=number

specifies the maximum number of tuning phases. The default is 24.

NBI=number

specifies the number of burn-in iterations before the chains are saved. The default is 1,000.

NMC=number

specifies the number of iterations after the burn-in. The default is 1,000.

NMCPRIOR=number

specifies the number of samples for the prior predictive analysis when **PLOTS(PRIOR)=BAYESPRED** is requested. The default is 10,000.

NTRDS=number**THREADS=number**

specifies the number of threads to be used. The number of threads cannot exceed the number of computer cores available. Each core samples the number of iterations that is specified by the NMC option. The default is 1.

NTU=number

specifies the number of samples for each tuning phase. The default is 500.

OUTPOST=SAS-data-set

names the SAS data set to contain the posterior samples. Alternatively, you can create the output data set by specifying an ODS OUTPUT statement as follows:

ODS OUTPUT POSTERIORSAMPLE = < SAS-data-set> ;

OUTPRIOR=SAS-data-set

names the SAS data set to contain the prior samples used to generate the prior predictive analysis when you request the prior predictive plots. Alternatively, you can create the output data set by specifying an ODS OUTPUT statement as follows:

ODS OUTPUT PRIORSAMPLE = < SAS-data-set> ;

PROPCOV=*value*

specifies the method used in constructing the initial covariance matrix for the Metropolis-Hastings algorithm. The QUANEW and NMSIMP methods find numerically approximated covariance matrices at the optimum of the posterior density function with respect to all continuous parameters. The tuning phase starts at the optimized values; in some problems, this can greatly increase convergence performance. If the approximated covariance matrix is not positive definite, then an identity matrix is used instead. You can specify the following values:

CONGRA

performs a conjugate-gradient optimization.

DBLDOG

performs a version of double-dogleg optimization.

NEWRAP

performs a Newton-Raphson optimization that combines a line-search algorithm with ridging.

NMSIMP

performs a Nelder-Mead simplex optimization.

NRRIDG

performs a Newton-Raphson optimization with ridging.

QUANEW

performs a quasi-Newton optimization.

TRUREG

performs a trust-region optimization.

SAMPLING=MULTIMETROPOLIS | UNIMETROPOLIS | MODELMETROPOLIS

specifies how to sample from the posterior distribution. SAMPLING=MULTIMETROPOLIS implements a Metropolis sampling scheme on a single block that contains all the parameters of the model. SAMPLING=MODELMETROPOLIS implements a Metropolis sampling scheme on multiple blocks: one block for each model (all the parameters of the model) plus a block for all the correlation parameters across the models. SAMPLING=UNIMETROPOLIS implements a Metropolis sampling scheme on multiple blocks, one for each parameter of the model. By default, SAMPLING=MULTIMETROPOLIS.

SEED=*number*

specifies an integer seed in the range 1 to $2^{31} - 1$ for the random number generator in the simulation. Specifying a seed enables you to reproduce identical Markov chains for the same specification. If you do not specify the SEED= option, or if you specify a nonpositive seed, a random seed is derived from the time of day.

SIMTIME

prints the time required for the MCMC sampling.

STATISTICS <(global-options)> = ALL | NONE | keyword | (keyword-list)**STATS** <(global-options)> = ALL | NONE | keyword | (keyword-list)

controls the number of posterior statistics produced. Specifying STATISTICS=ALL is equivalent to specifying STATISTICS= (CORR COV INTERVAL PRIOR SUMMARY). If you do not want any posterior statistics, specify STATISTICS=NONE. The default is STATISTICS=(SUMMARY INTERVAL). You can specify the following *global-options*:

ALPHA=*numeric-list*

controls the probabilities of the credible intervals. The ALPHA= values must be between 0 and 1. Each ALPHA= value produces a pair of 100(1-ALPHA)% equal-tail and HPD intervals for each parameter. The default is ALPHA=0.05, which yields the 95% credible intervals for each parameter.

PERCENT=*numeric-list*

requests the percentile points of the posterior samples. The PERCENT= values must be between 0 and 100. The default is PERCENT=25, 50, 75, which yields the 25th, 50th, and 75th percentile points, respectively, for each parameter.

You can specify the following *keywords*:

CORR

produces the posterior correlation matrix.

COV

produces the posterior covariance matrix.

INTERVAL

produces equal-tail credible intervals and HPD intervals. The default is to produce the 95% equal-tail credible intervals and 95% HPD intervals, but you can use the ALPHA= *global-option* to request intervals of any probabilities.

NONE

suppresses printing of all summary statistics.

PRIOR

produces a summary table of the prior distributions used in the Bayesian analysis.

SUMMARY

produces the means, standard deviations, and percentile points (25th, 50th, and 75th) for the posterior samples. You can use the global PERCENT= *global-option* to request specific percentile points.

THIN=*number*

THINNING=*number*

controls the thinning of the Markov chain. Only one in every k samples is used when THIN= k , and if NBI= n_0 and NMC= n , the number of samples that are kept is

$$\left[\frac{n_0 + n}{k} \right] - \left[\frac{n_0}{k} \right]$$

where $[a]$ represents the integer part of the number a . The default is THIN=1.

BOUNDS Statement

BOUNDS *bound1* < , *bound2* ... > ;

The BOUNDS statement imposes simple boundary constraints on the parameter estimates. BOUNDS statement constraints refer to the parameters estimated by the QLIM procedure. Any number of BOUNDS statements can be specified.

Each *bound* is composed of parameters and constants and inequality operators. Parameters associated with regressor variables are referred to by the names of the corresponding regressor variables:

item operator item < operator item < operator item ... > >

Each *item* is a constant, the name of a parameter, or a list of parameter names. See the section “[Naming of Parameters](#)” on page 1595 for more details on how parameters are named in the QLIM procedure. Each *operator* is '<', '>', '<=', or '>='.

Both the BOUNDS statement and the RESTRICT statement can be used to impose boundary constraints; however, the BOUNDS statement provides a simpler syntax for specifying these kinds of constraints. See the “[RESTRICT Statement](#)” on page 1551 for more information.

The following BOUNDS statement constrains the estimates of the parameters associated with the variable *ttime* and the variables *x1* through *x10* to be between 0 and 1. This example illustrates the use of parameter lists to specify boundary constraints.

```
bounds 0 < ttime x1-x10 < 1;
```

The following BOUNDS statement constrains the estimates of the correlation (*_RHO*) and sigma (*_SIGMA*) in the bivariate model:

```
bounds _rho >= 0, _sigma.y1 > 1, _sigma.y2 < 5;
```

The BOUNDS statement is not supported if a BAYES statement is also specified. In Bayesian analysis, the restrictions on parameters are usually introduced through the prior distribution.

BY Statement

BY *variables* ;

A BY statement can be used with PROC QLIM to obtain separate analyses on observations in groups defined by the BY variables.

CLASS Statement

CLASS *variables* ;

The CLASS statement names the classification variables to be used in the analysis. Classification variables can be either character or numeric.

Class levels are determined from the formatted values of the CLASS variables. Thus, you can use formats to group values into levels. See the discussion of the FORMAT procedure in *SAS Language Reference: Dictionary* for details.

ENDOGENOUS Statement

ENDOGENOUS *variables ~ options* ;

The ENDOGENOUS statement specifies the type of dependent variables that appear on the left-hand side of the equation. Endogenous variables listed refer to the dependent variables that appear on the left-hand side of the equation.

Discrete Variable Options

DISCRETE *<(discrete-options)>*

specifies that the endogenous variables in this statement are discrete. Valid *discrete-options* are as follows:

ORDER=DATA | FORMATTED | FREQ | INTERNAL

specifies the sorting order for the levels of the discrete variables specified in the ENDOGENOUS statement. This ordering determines which parameters in the model correspond to each level in the data. The following table shows how PROC QLIM interprets values of the ORDER= option.

Value of ORDER=	Levels Sorted By
DATA	Order of appearance in the input data set
FORMATTED	Formatted value
FREQ	Descending frequency count; levels with the most observations come first in the order
INTERNAL	Unformatted value

By default, ORDER=FORMATTED. For the values FORMATTED and INTERNAL, the sort order is machine dependent. For more information about sorting order, see the chapter on the SORT procedure in the *Base SAS Procedures Guide*.

DISTRIBUTION=NORMAL | LOGISTIC

DIST=NORMAL | LOGISTIC

D=NORMAL | LOGISTIC

specifies the cumulative distribution function used to model the response probabilities. DISTRIBUTION=NORMAL specifies the normal distribution for the probit model. DISTRIBUTION=LOGISTIC specifies the logistic distribution for the logit model.

By default, DISTRIBUTION=NORMAL.

If a multivariate model is specified, logistic distribution is not allowed. Only normal distribution is supported.

Censored Variable Options

CENSORED *(censored-options)*

specifies that the endogenous variables in this statement be censored. Valid *censored-options* are as follows:

LB=*value or variable*

LOWERBOUND=*value or variable*

specifies the lower bound of the censored variables. If *value* is missing or the value in *variable* is missing, no lower bound is set. By default, no lower bound is set.

UB=*value or variable*

UPPERBOUND=*value or variable*

specifies the upper bound of the censored variables. If *value* is missing or the value in *variable* is missing, no upper bound is set. By default, no upper bound is set.

Truncated Variable Options

TRUNCATED (*truncated-options*)

specifies that the endogenous variables in this statement be truncated. Valid *truncated-options* are as follows:

LB=*value or variable*

LOWERBOUND=*value or variable*

specifies the lower bound of the truncated variables. If *value* is missing or the value in *variable* is missing, no lower bound is set. By default, no lower bound is set.

UB=*value or variable*

UPPERBOUND=*value or variable*

specifies the upper bound of the truncated variables. If *value* is missing or the value in *variable* is missing, no upper bound is set. By default, no upper bound is set.

Stochastic Frontier Variable Options

FRONTIER <(*frontier-options*)>

specifies that the endogenous variable in this statement follow a production or cost frontier. Valid *frontier-options* are as follows:

TYPE=HALF | EXPONENTIAL | TRUNCATED

specifies the model type:

HALF specifies a half-normal model.

EXPONENTIAL specifies an exponential model.

TRUNCATED specifies a truncated normal model.

PRODUCTION

specifies that the model estimated be a production function.

COST

specifies that the model estimated be a cost function.

If neither PRODUCTION nor COST option is specified, production function is estimated by default.

Selection Options

SELECT (*select-option*)

specifies selection criteria for sample selection model. The BAYES statement does not support the SELECT option. The *select-option* specifies the condition for the endogenous variable to be selected. It is written as a variable name, followed by an equality operator (=) or an inequality operator (<, >, <=, >=), followed by a number:

variable operator number

The *variable* is the endogenous variable that the selection is based on. The *operator* can be =, <, >, <=, or >=. Multiple *select-options* can be combined with the logic operators: AND, OR. The following example illustrates the use of the SELECT option:

```
endogenous y1 ~ select (z=0);
endogenous y2 ~ select (z=1 or z=2);
```

The SELECT option can be used together with the DISCRETE, CENSORED, or TRUNCATED option. For example:

```
endogenous y1 ~ select (z=0) discrete;
endogenous y2 ~ select (z=1) censored (lb=0);
endogenous y3 ~ select (z=1 or z=2) truncated (ub=10);
```

For more information about selection models with censoring or truncation, see the section “[Selection Models](#)” on page 1565.

Endogeneity and Overidentification Test Options

ENDOTEST (*regressors*)

requests the test of endogeneity for a list of regressors in the model. More specifically, this option tests the null hypothesis that the specified regressors are exogenous. Each of these regressors must also have a model of its own. The former model is considered the structural model, and the latter models are considered reduced form models.

The following example illustrates the use of the ENDOTEST option by testing whether the regressors y2 and y3 are endogenous in the model for y1:

```
proc qlim;
  model y1 = y2 y3 x1;
  model y2 = x1 x2 x3 x4 x5;
  model y3 = x1 x2 x3 x4 x5;
  endogenous y1 ~ endotest (y2 y3);
run;
```

The ENDOTEST option is not available when you specify the SELECT or FRONTIER option. You can specify the ENDOTEST option only once for each ENDOGENOUS statement.

For more information about the test for endogeneity, see the section “[Test for Endogeneity](#)” on page 1575.

OVERID (*variables*)

requests the overidentification test for a list of variables. These variables are the overidentifying instrumental variables that you provide from the reduced form models. For more information, see the section “[Overidentification Test](#)” on page 1576.

The following example illustrates the use of the OVERID option:

```
proc qlim;
  model y1 = y2 y3 x1;
  model y2 = x1 x2 x3 x4 x5;
  model y3 = x1 x2 x3 x4 x5;
  endogenous y1 ~ overid(y2.x4 y3.x5);
run;
```

The regressors y_2 and y_3 in the model for y_1 are the endogenous variables. Therefore, each of these variables has its own models, which are considered reduced form models. The overidentifying instrumental variables are x_4 and x_5 . If you specify the OVERID option as

```
endogenous y1 ~ overid(y2.x4 y2.x5);
```

then you consider only the regressor y_2 to be endogenous, and the model for y_3 is ignored during the testing process.

The OVERID option is not available when you specify the SELECT or FRONTIER option. You can specify the OVERID option only once for each ENDOGENOUS statement.

FREQ Statement

FREQ *variable* ;

The FREQ statement identifies a variable that contains the frequency of occurrence of each observation. PROC QLIM treats each observation as if it appears n times, where n is the value of the FREQ variable for the observation. If it is not an integer, the frequency value is truncated to an integer. If the frequency value is less than 1 or missing, the observation is not used in the model fitting. When the FREQ statement is not specified, each observation is assigned a frequency of 1. If you specify more than one FREQ statement, then the first FREQ statement is used.

HETERO Statement

HETERO *dependent variables* ~ *exogenous variables* </ options > ;

The HETERO statement specifies variables that are related to the heteroscedasticity of the residuals and the way these variables are used to model the error variance. The heteroscedastic regression model supported by PROC QLIM is

$$y_i = \mathbf{x}_i' \boldsymbol{\beta} + \epsilon_i$$

$$\epsilon_i \sim N(0, \sigma_i^2)$$

See the section “[Heteroscedasticity](#)” on page 1563 for more details on the specification of functional forms.

LINK=value

The functional form can be specified using the LINK= option. The following option values are allowed:

EXP specifies the exponential link function

$$\sigma_i^2 = \sigma^2(1 + \exp(\mathbf{z}_i' \boldsymbol{\gamma}))$$

LINEAR specifies the linear link function

$$\sigma_i^2 = \sigma^2(1 + \mathbf{z}_i' \boldsymbol{\gamma})$$

When the LINK= option is not specified, the exponential link function is specified by default.

NOCONST

specifies that there be no constant in the exponential heteroscedasticity model.

$$\sigma_i^2 = \sigma^2 \exp(\mathbf{z}_i' \boldsymbol{\gamma})$$

SQUARE

estimates the model by using the square of linear heteroscedasticity function. For example, you can specify the following heteroscedasticity function:

$$\sigma_i^2 = \sigma^2(1 + (\mathbf{z}_i' \boldsymbol{\gamma})^2)$$

```
model y = x1 x2 / discrete;
hetero y ~ z1 / link=linear square;
```

The option SQUARE does not apply to exponential heteroscedasticity function because the square of an exponential function of $\mathbf{z}_i' \boldsymbol{\gamma}$ is the same as the exponential of $2\mathbf{z}_i' \boldsymbol{\gamma}$. Hence the only difference is that all $\boldsymbol{\gamma}$ estimates are divided by two.

You can use the HETERO statement within a Bayesian framework, but you should do this carefully because convergence can be slower than in the homoscedastic case. For more information see “[Priors for Heteroscedastic Models](#)” on page 1585.

INIT Statement

```
INIT initvalue1 < , initvalue2 ... > ;
```

The INIT statement sets initial values for parameters in the optimization. You can specify any number of INIT statements.

Each *initvalue* is written as a parameter or parameter list, followed by an optional equality operator (=), followed by a number:

```
parameter <=> number
```

If you also specify the BAYES statement, the INIT statement also initializes the Markov chain Monte Carlo (MCMC) algorithm. In particular, the INIT statement does one of the following:

- It initializes the tuning phase (this also includes the PROPCOV option).
- It initializes the sampling phase, if there is no tuning phase.

MODEL Statement

MODEL *dependent* = *regressors* < / *options* > ;

The MODEL statement specifies the dependent variable and independent regressor variables for the regression model.

You can specify the following *options* after a slash (/).

LIMIT1=ZERO | VARYING

specifies the restriction of the threshold value of the first category when the ordinal probit or logit model is estimated. LIMIT1=ZERO is the default option. When LIMIT1=VARYING is specified, the threshold value is estimated.

NOINT

suppresses the intercept parameter.

Endogenous Variable Options

The endogenous variable options are the same as the options that you can specify in the ENDOGENOUS statement. If you specify an ENDOGENOUS statement, all endogenous options in the MODEL statement are ignored.

Endogeneity and Overidentification Test Options

The endogeneity and overidentification test options are the same as the options that you can specify in the ENDOGENOUS statement. If you specify an ENDOGENOUS statement, all endogeneity and overidentification test options in the MODEL statement are ignored.

BOXCOX Estimation Options

BOXCOX (*option-list*)

specifies options that are used for Box-Cox regression or regressor transformation. For example, the Box-Cox regression is specified as

```
model y = x1 x2 / boxcox(y=lambda,x1 x2)
```

PROC QLIM estimates the following Box-Cox regression model:

$$y_i^{(\lambda)} = \beta_0 + \beta_1 x_{1i}^{(\lambda_2)} + \beta_2 x_{2i}^{(\lambda_2)} + \epsilon_i$$

The *option-list* takes the form *variable-list* < = *varname* > separated by commas. The *variable-list* specifies that the list of variables have the same Box-Cox transformation; *varname* specifies the name of this Box-Cox coefficient. If *varname* is not specified, the coefficient is called *_Lambdai*, where *i* increments sequentially.

Variable Selection Options

SELECTVAR <=(*selectvar-option*)>

enables variable selection. The *selectvar-option* specifies a variable selection method based on an information criterion. For more information, see the section “[Variable Selection](#)” on page 1569. You can specify the following *selectvar-options*:

DIRECTION=FORWARD | BACKWARD

specifies the searching algorithm to use in the variable selection method. By default, DIRECTION=FORWARD.

CRITER=AIC | SBC

specifies the information criterion to use for the variable selection. By default, CRITER=AIC.

MAXSTEPS=*value*

specifies the maximum number of steps that are allowed in the search algorithm. The default is infinite; that is, the algorithm does not stop until the stopping criterion is satisfied.

LSTOP=*value*

specifies the stopping criterion. The *value* represents the percentage of decrease or increase in the AIC or SBC that is required for the algorithm to proceed; it must be a positive number less than 1. The default is 0.

RETAIN(*regressors*)

specifies a list of regressors that are to be retained in any model that the variable selection process considers.

The following rules apply to how regressors are handled when you specify more than one MODEL statement and use the SELECTVAR option:

- If you do not specify the SELECTVAR option in a particular MODEL statement, then all regressors in the original model are included in any model that the variable selection algorithm considers. In other words, omitting the SELECTVAR option is equivalent to providing the option: SELECTVAR=(RETAIN(*all-regressors*)).
- If you specify the SELECTVAR option without any =(*option*) clause in a MODEL statement, then all regressors in that model (other than the intercept, if present) are eligible for potential exclusion as the variable selection process is executed.

The following example specifies 10 possible regressor candidates, out of which five are selected using the AIC criterions:

```
proc qlim data=one;
  model y = x1-x10 /selectvar=(direction=forward criter=AIC maxsteps=5);
run;
```

NLOPTIONS Statement

NLOPTIONS <*options*> ;

PROC QLIM uses the nonlinear optimization (NLO) subsystem to perform nonlinear optimization tasks. For a list of all the options of the NLOPTIONS statement, see Chapter 6, “[Nonlinear Optimization Methods](#).”

OUTPUT Statement

OUTPUT <OUT=SAS-data-set> <output-options> ;

The OUTPUT statement creates a new SAS data set containing all variables in the input data set and, optionally, the estimates of $\mathbf{x}'\boldsymbol{\beta}$, predicted value, residual, marginal effects, probability, standard deviation of the error, expected value, conditional expected value, technical efficiency measures, and inverse Mills ratio. When the response values are missing for the observation, all output estimates except residual are still computed as long as none of the explanatory variables is missing. This enables you to compute these statistics for prediction. You can specify only one OUTPUT statement.

Details on the specifications in the OUTPUT statement are as follows:

CONDITIONAL

outputs estimates of conditional expected values of continuous endogenous variables.

ERRSTD

outputs estimates of σ_j , the standard deviation of the error term.

EXPECTED

outputs estimates of expected values of continuous endogenous variables.

MARGINAL

outputs marginal effects.

MILLS

outputs estimates of inverse Mills ratios of censored or truncated continuous, binary discrete, and selection endogenous variables.

OUT=SAS-data-set

names the output data set.

PREDICTED

outputs estimates of predicted endogenous variables.

PROB

outputs estimates of probability of discrete endogenous variables taking the current observed responses.

PROBALL

outputs estimates of probability of discrete endogenous variables for all possible responses.

RESIDUAL

outputs estimates of residuals of continuous endogenous variables.

XBETA

outputs estimates of $\mathbf{x}'\boldsymbol{\beta}$.

TE1

outputs estimates of technical efficiency for each producer in the stochastic frontier model suggested by Battese and Coelli (1988).

TE2

outputs estimates of technical efficiency for each producer in the stochastic frontier model suggested by Jondrow et al. (1982).

PRIOR Statement

PRIOR *_REGRESSORS* | *parameter-list* ~ *distribution* ;

The PRIOR statement specifies the prior distribution of the model parameters. You must specify a single parameter or a list of parameter, a tilde ~, and then a distribution with its parameters. Multiple PRIOR statements are allowed.

You can specify the following *distributions*:

NORMAL(MEAN= μ , VAR= σ^2)

specifies a normal distribution with parameters MEAN and VAR.

GAMMA(SHAPE= a , SCALE= b)

specifies a gamma distribution with parameters SHAPE and SCALE.

IGAMMA(SHAPE= a , SCALE= b)

specifies an inverse gamma distribution with parameters SHAPE and SCALE.

UNIFORM(MIN= m , MAX= M)

specifies a uniform distribution that is defined between MIN and MAX.

BETA(SHAPE1= a , SHAPE2= b , MIN= m , MAX= M)

specifies a beta distribution with parameters SHAPE1 and SHAPE2 and defined between MIN and MAX.

T(LOCATION= μ , DF= ν)

specifies a noncentral t distribution with DF degrees of freedom and location parameter equal to LOCATION.

See the section “[Standard Distributions](#)” on page 1588 for details about how to specify *distributions*.

You can specify the special keyword REGRESSORS to select all the parameters used in the linear regression component of the model.

RESTRICT Statement

RESTRICT *restriction1* <, *restriction2* ... > ;

The RESTRICT statement is used to impose linear restrictions on the parameter estimates. Any number of RESTRICT statements can be specified, but the number of restrictions imposed is limited by the number of regressors.

Each *restriction* is written as an expression, followed by an equality operator (=) or an inequality operator (<, >, <=, >=), followed by a second expression:

expression operator expression

The *operator* can be =, <, >, <=, or >=. The operator and second expression are optional.

Restriction expressions can be composed of parameter names, multiplication (*), addition (+) and substitution (−) operators, and constants. Parameters named in restriction expressions must be among the parameters estimated by the model. Parameters associated with a regressor variable are referred to by the name of the corresponding regressor variable. The restriction expressions must be a linear function of the parameters.

The following is an example of the use of the RESTRICT statement:

```
proc qlim data=one;
  model y = x1-x10 / discrete;
  restrict x1*2 <= x2 + x3;
run;
```

The RESTRICT statement can also be used to impose cross-equation restrictions in multivariate models. The following RESTRICT statement imposes an equality restriction on coefficients of x_1 in equation y_1 and x_1 in equation y_2 :

```
proc qlim data=one;
  model y1 = x1-x10;
  model y2 = x1-x4;
  endogenous y1 y2 ~ discrete;
  restrict y1.x1=y2.x1;
run;
```

The RESTRICT statement is not supported if a BAYES statement is also specified. In Bayesian analysis, the restrictions on parameters are usually introduced through the prior distribution.

TEST Statement

<'label':> TEST <'string':> equation [,equation...] / options ;

The TEST statement performs Wald, Lagrange multiplier, and likelihood ratio tests of linear hypotheses about the regression parameters in the preceding MODEL statement. Each equation specifies a linear hypothesis to be tested. All hypotheses in one TEST statement are tested jointly. Variable names in the equations must correspond to regressors in the preceding MODEL statement, and each name represents the coefficient of the corresponding regressor. The keyword INTERCEPT refers to the coefficient of the intercept.

The following options can be specified in the TEST statement after the slash (/):

ALL

requests Wald, Lagrange multiplier, and likelihood ratio tests.

WALD

requests the Wald test.

LM

requests the Lagrange multiplier test.

LR

requests the likelihood ratio test.

The following illustrates the use of the TEST statement:

```
proc qlim;
  model y = x1 x2 x3;
  test x1 = 0, x2 * .5 + 2 * x3 = 0;
  test _int: test intercept = 0, x3 = 0;
run;
```

The first test investigates the joint hypothesis that

$$\beta_1 = 0$$

and

$$0.5\beta_2 + 2\beta_3 = 0$$

In case there is more than one MODEL statement in one QLIM procedure, then TEST statement is capable of testing cross-equation restrictions. Each parameter reference should be preceded by the name of the dependent variable of the particular model and the dot sign. For example,

```
proc qlim;
  model y1 = x1 x2 x3;
  model y2 = x3 x5 x6;
  test y1.x1 + y2.x6 = 1;
run;
```

This cross-equation test investigates the null hypothesis that

$$\beta_{1,1} + \beta_{2,3} = 1$$

in the system of equations

$$\begin{aligned} y_{1,i} &= \alpha_1 + \beta_{1,1}x_{1,i} + \beta_{1,2}x_{2,i} + \beta_{1,3}x_{3,i} \\ y_{2,i} &= \alpha_2 + \beta_{2,1}x_{3,i} + \beta_{2,2}x_{5,i} + \beta_{2,3}x_{6,i} \end{aligned}$$

Only linear equality restrictions and tests are permitted in PROC QLIM. Tests expressions can be composed only of algebraic operations involving the addition symbol (+), subtraction symbol (-), and multiplication symbol (*).

The TEST statement accepts labels that are reproduced in the printed output. TEST statement can be labeled in two ways. A TEST statement can be preceded by a label followed by a colon. Alternatively, the keyword TEST can be followed by a quoted string. If both are present, PROC QLIM uses the label preceding the colon. In the event no label is present, PROC QLIM automatically labels the tests.

You cannot specify both the TEST statement and the BAYES statement.

WEIGHT Statement

WEIGHT *variable* *</option>* ;

The WEIGHT statement specifies a variable to supply weighting values to use for each observation in estimating parameters. The log likelihood for each observation is multiplied by the corresponding weight variable value.

If the weight of an observation is nonpositive, that observation is not used in the estimation.

The following option can be added to the WEIGHT statement after a slash (/).

NONNORMALIZE

specifies that the weights are required to be used as is. When this option is not specified, the weights are normalized so that they add up to the actual sample size. Weights w_i are normalized by multiplying them by $\frac{n}{\sum_{i=1}^n w_i}$, where n is the sample size.

Details: QLIM Procedure

Ordinal Discrete Choice Modeling

Binary Probit and Logit Model

The binary choice model is

$$y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \epsilon_i$$

where value of the latent dependent variable, y_i^* , is observed only as follows:

$$\begin{aligned} y_i &= 1 && \text{if } y_i^* > 0 \\ &= 0 && \text{otherwise} \end{aligned}$$

The disturbance, ϵ_i , of the probit model has standard normal distribution with the distribution function (CDF)

$$\Phi(x) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi}} \exp(-t^2/2) dt$$

The disturbance of the logit model has standard logistic distribution with the CDF

$$\Lambda(x) = \frac{\exp(x)}{1 + \exp(x)} = \frac{1}{1 + \exp(-x)}$$

The binary discrete choice model has the following probability that the event $\{y_i = 1\}$ occurs:

$$P(y_i = 1) = F(\mathbf{x}_i' \boldsymbol{\beta}) = \begin{cases} \Phi(\mathbf{x}_i' \boldsymbol{\beta}) & \text{(probit)} \\ \Lambda(\mathbf{x}_i' \boldsymbol{\beta}) & \text{(logit)} \end{cases}$$

The log-likelihood function is

$$\ell = \sum_{i=1}^N \{y_i \log[F(\mathbf{x}_i' \boldsymbol{\beta})] + (1 - y_i) \log[1 - F(\mathbf{x}_i' \boldsymbol{\beta})]\}$$

where the CDF $F(x)$ is defined as $\Phi(x)$ for the probit model while $F(x) = \Lambda(x)$ for logit. The first order derivatives of the logit model are

$$\frac{\partial \ell}{\partial \boldsymbol{\beta}} = \sum_{i=1}^N (y_i - \Lambda(\mathbf{x}_i' \boldsymbol{\beta})) \mathbf{x}_i$$

The probit model has more complicated derivatives

$$\frac{\partial \ell}{\partial \boldsymbol{\beta}} = \sum_{i=1}^N \left\{ \frac{(2y_i - 1)\phi[(2y_i - 1)\mathbf{x}_i' \boldsymbol{\beta}]}{\Phi[(2y_i - 1)\mathbf{x}_i' \boldsymbol{\beta}]} \right\} \mathbf{x}_i = \sum_{i=1}^N r_i \mathbf{x}_i$$

where

$$r_i = \frac{(2y_i - 1)\phi[(2y_i - 1)\mathbf{x}_i' \boldsymbol{\beta}]}{\Phi[(2y_i - 1)\mathbf{x}_i' \boldsymbol{\beta}]}$$

Note that the logit maximum likelihood estimates are $\frac{\pi}{\sqrt{3}}$ times greater than probit maximum likelihood estimates, since the probit parameter estimates, $\boldsymbol{\beta}$, are standardized, and the error term with logistic distribution has a variance of $\frac{\pi^2}{3}$.

Ordinal Probit/Logit

When the dependent variable is observed in sequence with M categories, binary discrete choice modeling is not appropriate for data analysis. McKelvey and Zavoina (1975) proposed the ordinal (or ordered) probit model.

Consider the following regression equation:

$$y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \epsilon_i$$

where error disturbances, ϵ_i , have the distribution function F . The unobserved continuous random variable, y_i^* , is identified as M categories. Suppose there are $M + 1$ real numbers, μ_0, \dots, μ_M , where $\mu_0 = -\infty$, $\mu_1 = 0$, $\mu_M = \infty$, and $\mu_0 \leq \mu_1 \leq \dots \leq \mu_M$. Define

$$R_{i,j} = \mu_j - \mathbf{x}_i' \boldsymbol{\beta}$$

The probability that the unobserved dependent variable is contained in the j th category can be written as

$$P[\mu_{j-1} < y_i^* \leq \mu_j] = F(R_{i,j}) - F(R_{i,j-1})$$

The log-likelihood function is

$$\ell = \sum_{i=1}^N \sum_{j=1}^M d_{ij} \log[F(R_{i,j}) - F(R_{i,j-1})]$$

where

$$d_{ij} = \begin{cases} 1 & \text{if } \mu_{j-1} < y_i \leq \mu_j \\ 0 & \text{otherwise} \end{cases}$$

The first derivatives are written as

$$\frac{\partial \ell}{\partial \boldsymbol{\beta}} = \sum_{i=1}^N \sum_{j=1}^M d_{ij} \left[\frac{f(R_{i,j-1}) - f(R_{i,j})}{F(R_{i,j}) - F(R_{i,j-1})} \mathbf{x}_i \right]$$

$$\frac{\partial \ell}{\partial \mu_k} = \sum_{i=1}^N \sum_{j=1}^M d_{ij} \left[\frac{\delta_{j,k} f(R_{i,j}) - \delta_{j-1,k} f(R_{i,j-1})}{F(R_{i,j}) - F(R_{i,j-1})} \right]$$

where $f(x) = \frac{dF(x)}{dx}$ and $\delta_{j,k} = 1$ if $j = k$, and $\delta_{j,k} = 0$ otherwise. When the ordinal probit is estimated, it is assumed that $F(R_{i,j}) = \Phi(R_{i,j})$. The ordinal logit model is estimated if $F(R_{i,j}) = \Lambda(R_{i,j})$. The first threshold parameter, μ_1 , is estimated when the LIMIT1=VARYING option is specified. By default (LIMIT1=ZERO), so that $M - 2$ threshold parameters (μ_2, \dots, μ_{M-1}) are estimated.

The ordered probit models are analyzed by Aitchison and Silvey (1957), and Cox (1970) discussed ordered response data by using the logit model. They defined the probability that y_i^* belongs to j th category as

$$P[\mu_{j-1} < y_i \leq \mu_j] = F(\mu_j + \mathbf{x}_i' \boldsymbol{\theta}) - F(\mu_{j-1} + \mathbf{x}_i' \boldsymbol{\theta})$$

where $\mu_0 = -\infty$ and $\mu_M = \infty$. Therefore, the ordered response model analyzed by Aitchison and Silvey can be estimated if the LIMIT1=VARYING option is specified. Note that $\boldsymbol{\theta} = -\boldsymbol{\beta}$.

Goodness-of-Fit Measures

The goodness-of-fit measures discussed in this section apply only to discrete dependent variable models.

McFadden (1974) suggested a likelihood ratio index that is analogous to the R^2 in the linear regression model:

$$R_M^2 = 1 - \frac{\ln L}{\ln L_0}$$

where L is the value of the maximum likelihood function and L_0 is the value of a likelihood function when regression coefficients except an intercept term are zero. It can be shown that L_0 can be written as

$$L_0 = \sum_{j=1}^M N_j \ln\left(\frac{N_j}{N}\right)$$

where N_j is the number of responses in category j .

Estrella (1998) proposes the following requirements for a goodness-of-fit measure to be desirable in discrete choice modeling:

- The measure must take values in $[0, 1]$, where 0 represents no fit and 1 corresponds to perfect fit.
- The measure should be directly related to the valid test statistic for significance of all slope coefficients.

- The derivative of the measure with respect to the test statistic should comply with corresponding derivatives in a linear regression.

Estrella's (1998) measure is written

$$R_{E1}^2 = 1 - \left(\frac{\ln L}{\ln L_0} \right)^{-\frac{2}{N} \ln L_0}$$

An alternative measure suggested by Estrella (1998) is

$$R_{E2}^2 = 1 - [(\ln L - K) / \ln L_0]^{-\frac{2}{N} \ln L_0}$$

where $\ln L_0$ is computed with null slope parameter values, N is the number observations used, and K represents the number of estimated parameters.

Other goodness-of-fit measures are summarized as follows:

$$R_{CU1}^2 = 1 - \left(\frac{L_0}{L} \right)^{\frac{2}{N}} \quad (\text{Cragg} - \text{Uhler1})$$

$$R_{CU2}^2 = \frac{1 - (L_0/L)^{\frac{2}{N}}}{1 - L_0^{\frac{2}{N}}} \quad (\text{Cragg} - \text{Uhler2})$$

$$R_A^2 = \frac{2(\ln L - \ln L_0)}{2(\ln L - \ln L_0) + N} \quad (\text{Aldrich} - \text{Nelson})$$

$$R_{VZ}^2 = R_A^2 \frac{2 \ln L_0 - N}{2 \ln L_0} \quad (\text{Veall} - \text{Zimmermann})$$

$$R_{MZ}^2 = \frac{\sum_{i=1}^N (\hat{y}_i - \bar{\hat{y}})^2}{N + \sum_{i=1}^N (\hat{y}_i - \bar{\hat{y}})^2} \quad (\text{McKelvey} - \text{Zavoina})$$

where $\hat{y}_i = \mathbf{x}_i' \hat{\boldsymbol{\beta}}$ and $\bar{\hat{y}} = \sum_{i=1}^N \hat{y}_i / N$.

Limited Dependent Variable Models

Censored Regression Models

When the dependent variable is censored, values in a certain range are all transformed to a single value. For example, the standard tobit model can be defined as

$$y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \epsilon_i$$

$$y_i = \begin{cases} y_i^* & \text{if } y_i^* > 0 \\ 0 & \text{if } y_i^* \leq 0 \end{cases}$$

where $\epsilon_i \sim iid N(0, \sigma^2)$. The log-likelihood function of the standard censored regression model is

$$\ell = \sum_{i \in \{y_i=0\}} \ln[1 - \Phi(\mathbf{x}_i' \boldsymbol{\beta} / \sigma)] + \sum_{i \in \{y_i>0\}} \ln \left[\phi \left(\frac{y_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma} \right) / \sigma \right]$$

where $\Phi(\cdot)$ is the cumulative density function of the standard normal distribution and $\phi(\cdot)$ is the probability density function of the standard normal distribution.

The tobit model can be generalized to handle observation-by-observation censoring. The censored model on both of the lower and upper limits can be defined as

$$y_i = \begin{cases} R_i & \text{if } y_i^* \geq R_i \\ y_i^* & \text{if } L_i < y_i^* < R_i \\ L_i & \text{if } y_i^* \leq L_i \end{cases}$$

The log-likelihood function can be written as

$$\begin{aligned} \ell = & \sum_{i \in \{L_i < y_i < R_i\}} \ln \left[\phi\left(\frac{y_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}\right) / \sigma \right] + \sum_{i \in \{y_i = R_i\}} \ln \left[\Phi\left(-\frac{R_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}\right) \right] + \\ & \sum_{i \in \{y_i = L_i\}} \ln \left[\Phi\left(\frac{L_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}\right) \right] \end{aligned}$$

Log-likelihood functions of the lower- or upper-limit censored model are easily derived from the two-limit censored model. The log-likelihood function of the lower-limit censored model is

$$\ell = \sum_{i \in \{y_i > L_i\}} \ln \left[\phi\left(\frac{y_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}\right) / \sigma \right] + \sum_{i \in \{y_i = L_i\}} \ln \left[\Phi\left(\frac{L_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}\right) \right]$$

The log-likelihood function of the upper-limit censored model is

$$\ell = \sum_{i \in \{y_i < R_i\}} \ln \left[\phi\left(\frac{y_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}\right) / \sigma \right] + \sum_{i \in \{y_i = R_i\}} \ln \left[1 - \Phi\left(\frac{R_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}\right) \right]$$

Types of Tobit Models

Amemiya (1984) classified Tobit models into five types based on characteristics of the likelihood function. For notational convenience, let P denote a distribution or density function, y_{ji}^* is assumed to be normally distributed with mean $\mathbf{x}_{ji}' \boldsymbol{\beta}_j$ and variance σ_j^2 .

Type 1 Tobit

The Type 1 Tobit model was already discussed in the preceding section.

$$\begin{aligned} y_{1i}^* &= \mathbf{x}_{1i}' \boldsymbol{\beta}_1 + u_{1i} \\ y_{1i} &= y_{1i}^* \quad \text{if } y_{1i}^* > 0 \\ &= 0 \quad \text{if } y_{1i}^* \leq 0 \end{aligned}$$

The likelihood function is characterized as $P(y_1 < 0)P(y_1)$.

Type 2 Tobit

The Type 2 Tobit model is defined as

$$\begin{aligned}
 y_{1i}^* &= \mathbf{x}_{1i}'\boldsymbol{\beta}_1 + u_{1i} \\
 y_{2i}^* &= \mathbf{x}_{2i}'\boldsymbol{\beta}_2 + u_{2i} \\
 y_{1i} &= 1 \quad \text{if } y_{1i}^* > 0 \\
 &= 0 \quad \text{if } y_{1i}^* \leq 0 \\
 y_{2i} &= y_{2i}^* \quad \text{if } y_{1i}^* > 0 \\
 &= 0 \quad \text{if } y_{1i}^* \leq 0
 \end{aligned}$$

where $(u_{1i}, u_{2i}) \sim N(0, \Sigma)$. The likelihood function is described as $P(y_1 < 0)P(y_1 > 0, y_2)$.

Type 3 Tobit

The Type 3 Tobit model is different from the Type 2 Tobit in that y_{1i}^* of the Type 3 Tobit is observed when $y_{1i}^* > 0$.

$$\begin{aligned}
 y_{1i}^* &= \mathbf{x}_{1i}'\boldsymbol{\beta}_1 + u_{1i} \\
 y_{2i}^* &= \mathbf{x}_{2i}'\boldsymbol{\beta}_2 + u_{2i} \\
 y_{1i} &= y_{1i}^* \quad \text{if } y_{1i}^* > 0 \\
 &= 0 \quad \text{if } y_{1i}^* \leq 0 \\
 y_{2i} &= y_{2i}^* \quad \text{if } y_{1i}^* > 0 \\
 &= 0 \quad \text{if } y_{1i}^* \leq 0
 \end{aligned}$$

where $(u_{1i}, u_{2i})' \sim iidN(0, \Sigma)$.

The likelihood function is characterized as $P(y_1 < 0)P(y_1, y_2)$.

Type 4 Tobit

The Type 4 Tobit model consists of three equations:

$$\begin{aligned}
 y_{1i}^* &= \mathbf{x}_{1i}'\boldsymbol{\beta}_1 + u_{1i} \\
 y_{2i}^* &= \mathbf{x}_{2i}'\boldsymbol{\beta}_2 + u_{2i} \\
 y_{3i}^* &= \mathbf{x}_{3i}'\boldsymbol{\beta}_3 + u_{3i} \\
 y_{1i} &= y_{1i}^* \quad \text{if } y_{1i}^* > 0 \\
 &= 0 \quad \text{if } y_{1i}^* \leq 0 \\
 y_{2i} &= y_{2i}^* \quad \text{if } y_{1i}^* > 0 \\
 &= 0 \quad \text{if } y_{1i}^* \leq 0 \\
 y_{3i} &= y_{3i}^* \quad \text{if } y_{1i}^* \leq 0 \\
 &= 0 \quad \text{if } y_{1i}^* > 0
 \end{aligned}$$

where $(u_{1i}, u_{2i}, u_{3i})' \sim iidN(0, \Sigma)$. The likelihood function of the Type 4 Tobit model is characterized as $P(y_1 < 0, y_3)P(y_1, y_2)$.

Type 5 Tobit

The Type 5 Tobit model is defined as follows:

$$\begin{aligned}
 y_{1i}^* &= \mathbf{x}_{1i}'\boldsymbol{\beta}_1 + u_{1i} \\
 y_{2i}^* &= \mathbf{x}_{2i}'\boldsymbol{\beta}_2 + u_{2i} \\
 y_{3i}^* &= \mathbf{x}_{3i}'\boldsymbol{\beta}_3 + u_{3i} \\
 y_{1i} &= 1 \quad \text{if } y_{1i}^* > 0 \\
 &= 0 \quad \text{if } y_{1i}^* \leq 0 \\
 y_{2i} &= y_{2i}^* \quad \text{if } y_{1i}^* > 0 \\
 &= 0 \quad \text{if } y_{1i}^* \leq 0 \\
 y_{3i} &= y_{3i}^* \quad \text{if } y_{1i}^* \leq 0 \\
 &= 0 \quad \text{if } y_{1i}^* > 0
 \end{aligned}$$

where $(u_{1i}, u_{2i}, u_{3i})'$ are from *iid* trivariate normal distribution. The likelihood function of the Type 5 Tobit model is characterized as $P(y_1 < 0, y_3)P(y_1 > 0, y_2)$.

Code examples for these models can be found in “[Example 22.6: Types of Tobit Models](#)” on page 1610.

Truncated Regression Models

In a truncated model, the observed sample is a subset of the population where the dependent variable falls in a certain range. For example, when neither a dependent variable nor exogenous variables are observed for $y_i^* < 0$, the truncated regression model can be specified.

$$\ell = \sum_{i \in \{y_i \geq 0\}} \left\{ -\ln \Phi(\mathbf{x}_i'\boldsymbol{\beta}/\sigma) + \ln \left[\frac{\phi((y_i - \mathbf{x}_i'\boldsymbol{\beta})/\sigma)}{\sigma} \right] \right\}$$

Two-limit truncation model is defined as

$$y_i = y_i^* \text{ if } L_i \leq y_i^* \leq R_i$$

The log-likelihood function of the two-limit truncated regression model is

$$\ell = \sum_{i=1}^N \left\{ \ln \left[\phi\left(\frac{y_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}\right)/\sigma \right] - \ln \left[\Phi\left(\frac{R_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}\right) - \Phi\left(\frac{L_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}\right) \right] \right\}$$

The log-likelihood functions of the lower- and upper-limit truncation model are

$$\begin{aligned}
 \ell &= \sum_{i=1}^N \left\{ \ln \left[\phi\left(\frac{y_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}\right)/\sigma \right] - \ln \left[1 - \Phi\left(\frac{L_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}\right) \right] \right\} \quad (\text{lower}) \\
 \ell &= \sum_{i=1}^N \left\{ \ln \left[\phi\left(\frac{y_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}\right)/\sigma \right] - \ln \left[\Phi\left(\frac{R_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}\right) \right] \right\} \quad (\text{upper})
 \end{aligned}$$

Stochastic Frontier Production and Cost Models

Stochastic frontier production models were first developed by Aigner, Lovell, and Schmidt (1977); Meeusen and van den Broeck (1977). Specification of these models allow for random shocks of the production or cost but also include a term for technological or cost inefficiency. Assuming that the production function takes a log-linear Cobb-Douglas form, the stochastic frontier production model can be written as

$$\ln(y_i) = \beta_0 + \sum_n \beta_n \ln(x_{ni}) + \epsilon_i$$

where $\epsilon_i = v_i - u_i$. The v_i term represents the stochastic error component and u_i is the nonnegative, technology inefficiency error component. The v_i error component is assumed to be distributed iid normal and independently from u_i . If $u_i > 0$, the error term, ϵ_i , is negatively skewed and represents technology inefficiency. If $u_i < 0$, the error term ϵ_i is positively skewed and represents cost inefficiency. PROC QLIM models the u_i error component as a half normal, exponential, or truncated normal distribution.

The Normal-Half Normal Model

In case of the normal-half normal model, v_i is iid $N(0, \sigma_v^2)$, u_i is iid $N^+(0, \sigma_u^2)$ with v_i and u_i independent of each other. Given the independence of error terms, the joint density of v and u can be written as

$$f(u, v) = \frac{2}{2\pi\sigma_u\sigma_v} \exp \left\{ -\frac{u^2}{2\sigma_u^2} - \frac{v^2}{2\sigma_v^2} \right\}$$

Substituting $v = \epsilon + u$ into the preceding equation gives

$$f(u, \epsilon) = \frac{2}{2\pi\sigma_u\sigma_v} \exp \left\{ -\frac{u^2}{2\sigma_u^2} - \frac{(\epsilon + u)^2}{2\sigma_v^2} \right\}$$

Integrating u out to obtain the marginal density function of ϵ results in the following form:

$$\begin{aligned} f(\epsilon) &= \int_0^\infty f(u, \epsilon) du \\ &= \frac{2}{\sqrt{2\pi}\sigma} \left[1 - \Phi \left(\frac{\epsilon\lambda}{\sigma} \right) \right] \exp \left\{ -\frac{\epsilon^2}{2\sigma^2} \right\} \\ &= \frac{2}{\sigma} \phi \left(\frac{\epsilon}{\sigma} \right) \Phi \left(-\frac{\epsilon\lambda}{\sigma} \right) \end{aligned}$$

where $\lambda = \sigma_u/\sigma_v$ and $\sigma = \sqrt{\sigma_u^2 + \sigma_v^2}$.

In the case of a stochastic frontier cost model, $v = \epsilon - u$ and

$$f(\epsilon) = \frac{2}{\sigma} \phi \left(\frac{\epsilon}{\sigma} \right) \Phi \left(\frac{\epsilon\lambda}{\sigma} \right)$$

The log-likelihood function for the production model with N producers is written as

$$\ln L = \text{constant} - N \ln \sigma + \sum_i \ln \Phi \left(-\frac{\epsilon_i \lambda}{\sigma} \right) - \frac{1}{2\sigma^2} \sum_i \epsilon_i^2$$

The Normal-Exponential Model

Under the normal-exponential model, v_i is iid $N(0, \sigma_v^2)$ and u_i is iid exponential with scale parameter σ_u . Given the independence of error term components u_i and v_i , the joint density of v and u can be written as

$$f(u, v) = \frac{1}{\sqrt{2\pi}\sigma_u\sigma_v} \exp \left\{ -\frac{u}{\sigma_u} - \frac{v^2}{2\sigma_v^2} \right\}$$

The marginal density function of ϵ for the production function is

$$\begin{aligned} f(\epsilon) &= \int_0^\infty f(u, \epsilon) du \\ &= \left(\frac{1}{\sigma_u} \right) \Phi \left(-\frac{\epsilon}{\sigma_v} - \frac{\sigma_v}{\sigma_u} \right) \exp \left\{ \frac{\epsilon}{\sigma_u} + \frac{\sigma_v^2}{2\sigma_u^2} \right\} \end{aligned}$$

and the marginal density function for the cost function is equal to

$$f(\epsilon) = \left(\frac{1}{\sigma_u} \right) \Phi \left(\frac{\epsilon}{\sigma_v} - \frac{\sigma_v}{\sigma_u} \right) \exp \left\{ -\frac{\epsilon}{\sigma_u} + \frac{\sigma_v^2}{2\sigma_u^2} \right\}$$

The log-likelihood function for the normal-exponential production model with N producers is

$$\ln L = \text{constant} - N \ln \sigma_u + N \left(\frac{\sigma_v^2}{2\sigma_u^2} \right) + \sum_i \frac{\epsilon_i}{\sigma_u} + \sum_i \ln \Phi \left(-\frac{\epsilon_i}{\sigma_v} - \frac{\sigma_v}{\sigma_u} \right)$$

The Normal-Truncated Normal Model

The normal-truncated normal model is a generalization of the normal-half normal model by allowing the mean of u_i to differ from zero. Under the normal-truncated normal model, the error term component v_i is iid $N(0, \sigma_v^2)$ and u_i is iid $N^+(\mu, \sigma_u^2)$. The joint density of v_i and u_i can be written as

$$f(u, v) = \frac{1}{2\pi\sigma_u\sigma_v\Phi(\mu/\sigma_u)} \exp \left\{ -\frac{(u-\mu)^2}{2\sigma_u^2} - \frac{v^2}{2\sigma_v^2} \right\}$$

The marginal density function of ϵ for the production function is

$$\begin{aligned} f(\epsilon) &= \int_0^\infty f(u, \epsilon) du \\ &= \frac{1}{\sqrt{2\pi}\sigma\Phi(\mu/\sigma_u)} \Phi \left(\frac{\mu}{\sigma\lambda} - \frac{\epsilon\lambda}{\sigma} \right) \exp \left\{ -\frac{(\epsilon + \mu)^2}{2\sigma^2} \right\} \\ &= \frac{1}{\sigma} \phi \left(\frac{\epsilon + \mu}{\sigma} \right) \Phi \left(\frac{\mu}{\sigma\lambda} - \frac{\epsilon\lambda}{\sigma} \right) \left[\Phi \left(\frac{\mu}{\sigma_u} \right) \right]^{-1} \end{aligned}$$

and the marginal density function for the cost function is

$$f(\epsilon) = \frac{1}{\sigma} \phi \left(\frac{\epsilon - \mu}{\sigma} \right) \Phi \left(\frac{\mu}{\sigma\lambda} + \frac{\epsilon\lambda}{\sigma} \right) \left[\Phi \left(\frac{\mu}{\sigma_u} \right) \right]^{-1}$$

The log-likelihood function for the normal-truncated normal production model with N producers is

$$\ln L = \text{constant} - N \ln \sigma - N \ln \Phi \left(\frac{\mu}{\sigma_u} \right) + \sum_i \ln \Phi \left(\frac{\mu}{\sigma \lambda} - \frac{\epsilon_i \lambda}{\sigma} \right) - \frac{1}{2} \sum_i \left(\frac{\epsilon_i + \mu}{\sigma} \right)^2$$

For more detail on normal-half normal, normal-exponential, and normal-truncated models, see Kumbhakar and Lovell (2000); Coelli, Prasada Rao, and Battese (1998).

Heteroscedasticity and Box-Cox Transformation

Heteroscedasticity

If the variance of regression disturbance, (ϵ_i) , is heteroscedastic, the variance can be specified as a function of variables

$$E(\epsilon_i^2) = \sigma_i^2 = f(\mathbf{z}_i' \boldsymbol{\gamma})$$

The following table shows various functional forms of heteroscedasticity and the corresponding options to request each model.

No.	Model	Options
1	$f(\mathbf{z}_i' \boldsymbol{\gamma}) = \sigma^2(1 + \exp(\mathbf{z}_i' \boldsymbol{\gamma}))$	LINK=EXP (default)
2	$f(\mathbf{z}_i' \boldsymbol{\gamma}) = \sigma^2 \exp(\mathbf{z}_i' \boldsymbol{\gamma})$	LINK=EXP NOCONST
3	$f(\mathbf{z}_i' \boldsymbol{\gamma}) = \sigma^2(1 + \sum_{l=1}^L \gamma_l z_{li})$	LINK=LINEAR
4	$f(\mathbf{z}_i' \boldsymbol{\gamma}) = \sigma^2(1 + (\sum_{l=1}^L \gamma_l z_{li})^2)$	LINK=LINEAR SQUARE

For discrete choice models, σ^2 is normalized ($\sigma^2 = 1$) since this parameter is not identified. Note that in models 3 and 5, it may be possible that variances of some observations are negative. Although the QLIM procedure assigns a large penalty to move the optimization away from such region, it is possible that the optimization cannot improve the objective function value and gets locked in the region. Signs of such outcome include extremely small likelihood values or missing standard errors in the estimates. In models 2 and 6, variances are guaranteed to be greater or equal to zero, but it may be possible that variances of some observations are very close to zero. In these scenarios, standard errors may be missing. Models 1 and 4 do not have such problems. Variances in these models are always positive and never close to zero.

The heteroscedastic regression model is estimated using the following log-likelihood function:

$$\ell = -\frac{N}{2} \ln(2\pi) - \sum_{i=1}^N \frac{1}{2} \ln(\sigma_i^2) - \frac{1}{2} \sum_{i=1}^N \left(\frac{e_i}{\sigma_i} \right)^2$$

where $e_i = y_i - \mathbf{x}_i' \boldsymbol{\beta}$.

Box-Cox Modeling

The Box-Cox transformation on x is defined as

$$x^{(\lambda)} = \begin{cases} \frac{x^\lambda - 1}{\lambda} & \text{if } \lambda \neq 0 \\ \ln(x) & \text{if } \lambda = 0 \end{cases}$$

The Box-Cox regression model with heteroscedasticity is written as

$$\begin{aligned} y_i^{(\lambda_0)} &= \beta_0 + \sum_{k=1}^K \beta_k x_{ki}^{(\lambda_k)} + \epsilon_i \\ &= \mu_i + \epsilon_i \end{aligned}$$

where $\epsilon_i \sim N(0, \sigma_i^2)$ and transformed variables must be positive. In practice, too many transformation parameters cause numerical problems in model fitting. It is common to have the same Box-Cox transformation performed on all the variables — that is, $\lambda_0 = \lambda_1 = \dots = \lambda_K$. It is required for the magnitude of transformed variables to be in the tolerable range if the corresponding transformation parameters are $|\lambda| > 1$.

The log-likelihood function of the Box-Cox regression model is written as

$$\ell = -\frac{N}{2} \ln(2\pi) - \sum_{i=1}^N \ln(\sigma_i) - \frac{1}{2\sigma_i^2} \sum_{i=1}^N e_i^2 + (\lambda_0 - 1) \sum_{i=1}^N \ln(y_i)$$

where $e_i = y_i^{(\lambda_0)} - \mu_i$.

When the dependent variable is discrete, censored, or truncated, the Box-Cox transformation can be applied only to explanatory variables.

Bivariate Limited Dependent Variable Modeling

The generic form of a bivariate limited dependent variable model is

$$\begin{aligned} y_{1i}^* &= \mathbf{x}_{1i}' \boldsymbol{\beta}_1 + \epsilon_{1i} \\ y_{2i}^* &= \mathbf{x}_{2i}' \boldsymbol{\beta}_2 + \epsilon_{2i} \end{aligned}$$

where the disturbances, ϵ_{1i} and ϵ_{2i} , have joint normal distribution with zero mean, standard deviations σ_1 and σ_2 , and correlation of ρ . y_1^* and y_2^* are latent variables. The dependent variables y_1 and y_2 are observed if the latent variables y_1^* and y_2^* fall in certain ranges:

$$\begin{aligned} y_1 &= y_{1i} \text{ if } y_{1i}^* \in D_1(y_{1i}) \\ y_2 &= y_{2i} \text{ if } y_{2i}^* \in D_2(y_{2i}) \end{aligned}$$

D is a transformation from (y_{1i}^*, y_{2i}^*) to (y_{1i}, y_{2i}) . For example, if y_1 and y_2 are censored variables with lower bound 0, then

$$\begin{aligned} y_1 &= y_{1i} \text{ if } y_{1i}^* > 0, & y_1 &= 0 \text{ if } y_{1i}^* \leq 0 \\ y_2 &= y_{2i} \text{ if } y_{2i}^* > 0, & y_2 &= 0 \text{ if } y_{2i}^* \leq 0 \end{aligned}$$

There are three cases for the log likelihood of (y_{1i}, y_{2i}) . The first case is that $y_{1i} = y_{1i}^*$ and $y_{2i} = y_{2i}^*$. That is, this observation is mapped to one point in the space of latent variables. The log likelihood is computed from a bivariate normal density,

$$\ell_i = \ln \left[\phi_2 \left(\frac{y_1 - \mathbf{x}_1' \boldsymbol{\beta}_1}{\sigma_1}, \frac{y_2 - \mathbf{x}_2' \boldsymbol{\beta}_2}{\sigma_2}, \rho \right) \right] - \ln \sigma_1 - \ln \sigma_2$$

where $\phi_2(u, v, \rho)$ is the density function for standardized bivariate normal distribution with correlation ρ ,

$$\phi_2(u, v, \rho) = \frac{e^{-(1/2)(u^2 + v^2 - 2\rho uv)/(1 - \rho^2)}}{2\pi(1 - \rho^2)^{1/2}}$$

The second case is that one observed dependent variable is mapped to a point of its latent variable and the other dependent variable is mapped to a segment in the space of its latent variable. For example, in the bivariate censored model specified, if observed $y_1 > 0$ and $y_2 = 0$, then $y_1^* = y_1$ and $y_2^* \in (-\infty, 0]$. In general, the log likelihood for one observation can be written as follows (the subscript i is dropped for simplicity): If one set is a single point and the other set is a range, without loss of generality, let $D_1(y_1) = \{y_1\}$ and $D_2(y_2) = [L_2, R_2]$,

$$\begin{aligned} \ell_i = & \ln \left[\phi \left(\frac{y_1 - \mathbf{x}_1' \boldsymbol{\beta}_1}{\sigma_1} \right) \right] - \ln \sigma_1 \\ & + \ln \left[\Phi \left(\frac{R_2 - \mathbf{x}_2' \boldsymbol{\beta}_2 - \rho \frac{y_1 - \mathbf{x}_1' \boldsymbol{\beta}_1}{\sigma_1}}{\sigma_2} \right) - \Phi \left(\frac{L_2 - \mathbf{x}_2' \boldsymbol{\beta}_2 - \rho \frac{y_1 - \mathbf{x}_1' \boldsymbol{\beta}_1}{\sigma_1}}{\sigma_2} \right) \right] \end{aligned}$$

where ϕ and Φ are the density function and the cumulative probability function for standardized univariate normal distribution.

The third case is that both dependent variables are mapped to segments in the space of latent variables. For example, in the bivariate censored model specified, if observed $y_1 = 0$ and $y_2 = 0$, then $y_1^* \in (-\infty, 0]$ and $y_2^* \in (-\infty, 0]$. In general, if $D_1(y_1) = [L_1, R_1]$ and $D_2(y_2) = [L_2, R_2]$, the log likelihood is

$$\ell_i = \ln \int_{\frac{L_1 - \mathbf{x}_1' \boldsymbol{\beta}_1}{\sigma_1}}^{\frac{R_1 - \mathbf{x}_1' \boldsymbol{\beta}_1}{\sigma_1}} \int_{\frac{L_2 - \mathbf{x}_2' \boldsymbol{\beta}_2}{\sigma_2}}^{\frac{R_2 - \mathbf{x}_2' \boldsymbol{\beta}_2}{\sigma_2}} \phi_2(u, v, \rho) du dv$$

Selection Models

In sample selection models, one or several dependent variables are observed when another variable takes certain values. For example, the standard Heckman selection model can be defined as

$$\begin{aligned} z_i^* &= \mathbf{w}_i' \boldsymbol{\gamma} + u_i \\ z_i &= \begin{cases} 1 & \text{if } z_i^* > 0 \\ 0 & \text{if } z_i^* \leq 0 \end{cases} \\ y_i &= \mathbf{x}_i' \boldsymbol{\beta} + \epsilon_i \quad \text{if } z_i = 1 \end{aligned}$$

where u_i and ϵ_i are jointly normal with 0 mean, standard deviations of 1 and σ , respectively, and correlation of ρ . Selection is based on the variable z , and y is observed when z has a value of 1. Least squares regression

that uses the observed data of y produces inconsistent estimates of β . The maximum likelihood method is used to estimate selection models. It is also possible to estimate these models by using Heckman's method, which is more computationally efficient. But it can be shown that the resulting estimates, although consistent, are not asymptotically efficient under a normality assumption. Moreover, this method often violates the constraint on the correlation coefficient $|\rho| \leq 1$.

The log-likelihood function of the Heckman selection model is written as

$$\begin{aligned} \ell &= \sum_{i \in \{z_i=0\}} \ln[1 - \Phi(\mathbf{w}_i' \boldsymbol{\gamma})] \\ &+ \sum_{i \in \{z_i=1\}} \left\{ \ln \phi\left(\frac{y_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}\right) - \ln \sigma + \ln \Phi\left(\frac{\mathbf{w}_i' \boldsymbol{\gamma} + \rho \frac{y_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}}{\sqrt{1 - \rho^2}}\right) \right\} \end{aligned}$$

The selection can be based on only one variable, but the selection can lead to several variables. For example, selection is based on the variable z in the following switching regression model:

$$z_i^* = \mathbf{w}_i' \boldsymbol{\gamma} + u_i$$

$$z_i = \begin{cases} 1 & \text{if } z_i^* > 0 \\ 0 & \text{if } z_i^* \leq 0 \end{cases}$$

$$y_{1i} = \mathbf{x}_{1i}' \boldsymbol{\beta}_1 + \epsilon_{1i} \quad \text{if } z_i = 0$$

$$y_{2i} = \mathbf{x}_{2i}' \boldsymbol{\beta}_2 + \epsilon_{2i} \quad \text{if } z_i = 1$$

If $z = 0$, then y_1 is observed. If $z = 1$, then y_2 is observed. Because y_1 and y_2 are never observed at the same time, the correlation between y_1 and y_2 cannot be estimated. Only the correlation between z and y_1 and the correlation between z and y_2 can be estimated. This estimation uses the maximum likelihood method.

A brief example of the SAS statements for this model can be found in “[Example 22.4: Sample Selection Model](#)” on page 1606.

The Heckman selection model can include censoring or truncation. For a brief example of the SAS statements for these models see “[Example 22.5: Sample Selection Model with Truncation and Censoring](#)” on page 1607. The following example shows a variable y_i that is censored from below at zero:

$$z_i^* = \mathbf{w}_i' \boldsymbol{\gamma} + u_i$$

$$z_i = \begin{cases} 1 & \text{if } z_i^* > 0 \\ 0 & \text{if } z_i^* \leq 0 \end{cases}$$

$$y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \epsilon_i \quad \text{if } z_i = 1$$

$$y_i = \begin{cases} y_i^* & \text{if } y_i^* > 0 \\ 0 & \text{if } y_i^* \leq 0 \end{cases}$$

In this case, the log-likelihood function of the Heckman selection model needs to be modified as follows to include the censored region:

$$\begin{aligned}\ell &= \sum_{\{i|z_i=0\}} \ln[1 - \Phi(\mathbf{w}_i'\boldsymbol{\gamma})] \\ &+ \sum_{\{i|z_i=1, y_i=y_i^*\}} \left\{ \ln \left[\phi\left(\frac{y_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}\right) \right] - \ln \sigma + \ln \left[\Phi \left(\frac{\mathbf{w}_i'\boldsymbol{\gamma} + \rho \frac{y_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}}{\sqrt{1 - \rho^2}} \right) \right] \right\} \\ &+ \sum_{\{i|z_i=1, y_i=0\}} \ln \int_{-\infty}^{\frac{-\mathbf{x}_i'\boldsymbol{\beta}}{\sigma}} \int_{-\mathbf{w}_i'\boldsymbol{\gamma}}^{\infty} \phi_2(u, v, \rho) du dv\end{aligned}$$

In case y_i is truncated from below at 0 instead of censored, the likelihood function can be written as

$$\begin{aligned}\ell &= \sum_{\{i|z_i=0\}} \ln[1 - \Phi(\mathbf{w}_i'\boldsymbol{\gamma})] \\ &+ \sum_{\{i|z_i=1\}} \left\{ \ln \left[\phi\left(\frac{y_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}\right) \right] - \ln \sigma + \ln \left[\Phi \left(\frac{\mathbf{w}_i'\boldsymbol{\gamma} + \rho \frac{y_i - \mathbf{x}_i'\boldsymbol{\beta}}{\sigma}}{\sqrt{1 - \rho^2}} \right) \right] - \ln [\Phi(\mathbf{x}_i'\boldsymbol{\beta}/\sigma)] \right\}\end{aligned}$$

Heckman's Two-Step Selection Method

Sample selection bias arises from nonrandom selection of the sample from the population. A classic example is using a sample of market wages for working women to estimate female labor supply function. This sample is nonrandom because it includes only the wages of women whose market wage exceeds their home wage at zero hours of work.

A simple selection model can be written as the latent model

$$\begin{aligned}z_i^* &= \mathbf{w}_i'\boldsymbol{\gamma} + u_i \\ z_i &= \begin{cases} 1 & \text{if } z_i^* > 0 \\ 0 & \text{if } z_i^* \leq 0 \end{cases} \\ y_i &= \mathbf{x}_i'\boldsymbol{\beta} + \epsilon_i \quad \text{if } z_i = 1\end{aligned}$$

where u_i and ϵ_i are jointly normal with 0 mean, standard deviations of 1 and σ , respectively, and correlation of ρ . The dependent variable y_i (wage) is observed if the latent variable z_i^* (the difference between market wage and reservation wage) is positive or if the indicator variable z_i (labor force participation) is 1.

The model of interest that applies to the observations in the selected sample can be written as

$$E(y_i | \mathbf{x}_i, z_i = 1) = \mathbf{x}_i'\boldsymbol{\beta} + \rho\sigma\lambda(\mathbf{w}_i'\boldsymbol{\gamma})$$

where $\lambda(\mathbf{w}_i'\boldsymbol{\gamma}) = \phi(\mathbf{w}_i'\boldsymbol{\gamma}) / \Phi(\mathbf{w}_i'\boldsymbol{\gamma})$. Hence, the following regression equation is valid for the observations for which $z_i = 1$:

$$y_i = \mathbf{x}_i'\boldsymbol{\beta} + \rho\sigma\lambda(\mathbf{w}_i'\boldsymbol{\gamma}) + v_i$$

Therefore, estimates of β that are obtained from the OLS regression of y on x by using the selected sample (that is, the sample for which $z_i = 1$) suffer from omitted variable bias if selection bias is really the case. Although maximum likelihood estimation of β is consistent and efficient, Heckman's two-step method is more frequently used. Heckman's two-step method can be requested by specifying the HECKIT option of the QLIM statement.

Heckman's two-step method is as follows:

1. Obtain $\hat{\gamma}$, the estimate of the parameters of the probability that $z_i^* > 0$, by using regressors w_i and the binary dependent variable z_i by probit analysis for the full sample. Compute $\hat{\lambda}_i = \lambda(w_i' \hat{\gamma})$.
2. Obtain $\hat{\beta}$ and $\hat{\beta}_\lambda$, the estimates of β and $\rho\sigma$, by least squares regression of y_i on x_i and $\hat{\lambda}_i$ by using observations on the selected subsample.

The standard least squares estimators of the population variance σ^2 and the variances of the estimated coefficients are incorrect. To test hypotheses, the correct ones need to be calculated. An estimator of σ^2 is

$$\hat{\sigma}^2 = \frac{1}{N_1} \sum_{i=1}^{N_1} e_i^2 + \hat{\beta}_\lambda^2 \frac{1}{N_1} \sum_{i=1}^{N_1} \hat{\delta}_i$$

where N_1 is the selected subsample size, e_i is the residual for the i th observation obtained from step 2, and $\hat{\delta}_i = \hat{\lambda}_i^2 + \hat{\lambda}_i w_i' \hat{\gamma}$. Let X_* be an $N_1 \times (K + 1)$ matrix with i th row $[x_i' \ \lambda_i]$, and define W similarly with i th row w_i' . Then the estimator of the asymptotic covariance of $[\hat{\beta}, \hat{\beta}_\lambda]$ is

$$\text{EstAsyVar}[\hat{\beta}, \hat{\beta}_\lambda] = \hat{\sigma}^2 [X_*' X_*]^{-1} [X_*' (I - \hat{\rho}^2 \hat{\Delta}) X_* + Q] [X_*' X_*]^{-1}$$

where $\hat{\rho}^2 = \hat{\beta}_\lambda^2 / \hat{\sigma}^2$, $\hat{\Delta} = \text{diag}(\hat{\delta}_i)$, and

$$Q = \hat{\sigma}^2 (X_*' \hat{\Delta} W) \text{Est.Asy.Var}(\hat{\gamma}) (W' \hat{\Delta} X_*)$$

where $\text{Est.Asy.Var}(\hat{\gamma})$ is the estimator of the asymptotic covariance of the probit coefficients that are obtained in step 1. When you specify the HECKIT option, PROC QLIM uses a numerical estimated asymptotic variance.

When the HECKIT option is specified, PROC QLIM reports the corrected standard errors for $[\hat{\beta}, \hat{\beta}_\lambda]$ automatically. However, if you need the conventional OLS standard errors, you can specify the HECKIT(UNCORRECTED) option.

In the selected regression model, when the coefficient of $\lambda(w_i' \gamma)$ is 0, you do not need Heckman's two-step estimation method; a simple regression of y on x produces consistent estimates for β , and the OLS standard errors are correct. Thus, a standard t test on $\hat{\beta}_\lambda$ (which uses the estimate from step 2 and the uncorrected standard errors) is a valid test of the null hypothesis of no selection bias.

Although Heckman's two-step method uses the OLS method in the second stage, you can request the ML method by specifying the HECKIT(SECONDSTAGE=ML) option. When the second-stage method is the ML method, the model for y_i can be nonlinear.

Multivariate Limited Dependent Models

The multivariate model is similar to bivariate models. The generic form of the multivariate limited dependent variable model is

$$\begin{aligned} y_{1i}^* &= \mathbf{x}_{1i}' \boldsymbol{\beta}_1 + \epsilon_{1i} \\ y_{2i}^* &= \mathbf{x}_{2i}' \boldsymbol{\beta}_2 + \epsilon_{2i} \\ &\dots \\ y_{mi}^* &= \mathbf{x}_{mi}' \boldsymbol{\beta}_m + \epsilon_{mi} \end{aligned}$$

where m is the number of models to be estimated. The vector ϵ has multivariate normal distribution with mean 0 and variance-covariance matrix Σ . Similar to bivariate models, the likelihood may involve computing multivariate normal integrations. This is done using Monte Carlo integration. (See Genz (1992); Hajivassiliou and McFadden (1998).)

When the number of equations, N , increases in a system, the number of parameters increases at the rate of N^2 because of the correlation matrix. When the number of parameters is large, sometimes the optimization converges but some of the standard deviations are missing. This usually means that the model is over-parameterized. The default method for computing the covariance is to use the inverse Hessian matrix. The Hessian is computed by finite differences, and in over-parameterized cases, the inverse cannot be computed. It is recommended that you reduce the number of parameters in such cases. Sometimes using the outer product covariance matrix (COVEST=OP option) may also help.

Variable Selection

Variable Selection

Variable selection uses either Akaike's information criterion (AIC) or the Schwartz Bayesian criterion (SBC) and either a forward selection method or a backward elimination method.

Forward selection starts from a small subset of variables. In each step, the variable that gives the largest decrease in the value of the information criterion specified in the CRITER= option (AIC or SBC) is added. The process stops when the next candidate to be added does not reduce the value of the information criterion by more than the amount specified in the LSTOP= option in the MODEL statement.

Backward elimination starts from a larger subset of variables. In each step, one variable is dropped based on the information criterion that is chosen.

Tests on Parameters

Tests on Parameters

In general, the hypothesis tested can be written as

$$H_0 : \mathbf{h}(\theta) = 0$$

where $\mathbf{h}(\theta)$ is an r by 1 vector valued function of the parameters θ given by the r expressions specified in the TEST statement.

Let \hat{V} be the estimate of the covariance matrix of $\hat{\theta}$. Let $\hat{\theta}$ be the unconstrained estimate of θ and $\tilde{\theta}$ be the constrained estimate of θ such that $h(\tilde{\theta}) = 0$. Let

$$A(\theta) = \partial h(\theta) / \partial \theta \mid_{\hat{\theta}}$$

Using this notation, the test statistics for the three kinds of tests are computed as follows.

The Wald test statistic is defined as

$$W = h'(\hat{\theta}) \left(A(\hat{\theta}) \hat{V} A'(\hat{\theta}) \right)^{-1} h(\hat{\theta})$$

The Wald test is not invariant to reparameterization of the model (Gregory and Veall 1985, Gallant 1987, p. 219). For more information about the theoretical properties of the Wald test, see Phillips and Park (1988).

The Lagrange multiplier test statistic is

$$LM = \lambda' A(\tilde{\theta}) \tilde{V} A'(\tilde{\theta}) \lambda$$

where λ is the vector of Lagrange multipliers from the computation of the restricted estimate $\tilde{\theta}$.

The likelihood ratio test statistic is

$$LR = 2 \left(L(\hat{\theta}) - L(\tilde{\theta}) \right)$$

where $\tilde{\theta}$ represents the constrained estimate of θ and L is the concentrated log-likelihood value.

For each kind of test, under the null hypothesis the test statistic is asymptotically distributed as a χ^2 random variable with r degrees of freedom, where r is the number of expressions in the TEST statement. The p -values reported for the tests are computed from the $\chi^2(r)$ distribution and are only asymptotically valid.

Monte Carlo simulations suggest that the asymptotic distribution of the Wald test is a poorer approximation to its small sample distribution than that of the other two tests. However, the Wald test has the lowest computational cost, since it does not require computation of the constrained estimate $\tilde{\theta}$.

The following is an example of using the TEST statement to perform a likelihood ratio test:

```
proc qlim;
  model y = x1 x2 x3;
  test x1 = 0, x2 * .5 + 2 * x3 = 0 /lr;
run;
```

Endogeneity and Instrumental Variables

The PROC QLIM models such as qualitative response or limited dependent variable models assume that the errors are independent of the explanatory variables. If this assumption fails to hold, the distributional form that the likelihood is based on is misspecified and the obtained coefficients are inconsistent.

To begin, consider a linear model

$$y_i = y_i^* = \beta_0 + \beta_1 x_{1i} + \cdots + \beta_k x_{ki} + u_i$$

Assume that $E(u) = 0$, $\text{Cov}(x_j, u) = 0$ for $j = 1, \dots, k-1$, and $\text{Cov}(x_k, u) = \rho \neq 0$. Therefore, x_k is endogenous. The endogeneity comes from many sources, such as x_k having measurement error or omitting a variable that is correlated with x_k . If you ignore the endogeneity, you can estimate this model in PROC QLIM as follows (assuming $k = 4$):

```
proc qlim data=a;
  model y = x1 x2 x3 x4;
run;
```

However, this approach produces inconsistent maximum likelihood estimates. To obtain consistent maximum likelihood estimates, you should consider the joint density of the dependent variable and the endogenous variables. To do this in PROC QLIM, you need at least one instrument—that is, an observable variable, z_1 —that is not in the [structural equation](#) and that satisfies two conditions: z_1 is exogenous (that is, $\text{Cov}(z_1, u) = 0$), and z_1 must be correlated with the endogenous regressor x_k . Then, you can model x_k as

$$x_{ki} = \pi_0 + \pi_1 x_{1i} + \cdots + \pi_{k-1} x_{(k-1)i} + \theta z_{1i} + \epsilon_i$$

You can now write this reduced form equation along with the [structural equation](#) to obtain the consistent maximum likelihood estimates as follows:

```
proc qlim data=a;
  model y = x1 x2 x3 x4;
  model x4 = x1 x2 x3 z1;
run;
```

Estimating the structural model together with the reduced form models for the endogenous explanatory variables gives you the full information maximum likelihood (FIML) estimates. Because of the linearity of the model, you can estimate it efficiently and more simply by using the two-stage least squares estimator. However, PROC QLIM handles nonlinear models such as qualitative response and limited dependent variable models, and in their estimation it maximizes the corresponding joint likelihood function. In the case of endogeneity, when the reduced form models for the endogenous explanatory variables are written along with the structural model, PROC QLIM maximizes the likelihood function that is obtained from the joint density of the response variable and the endogenous explanatory variables. For example, consider the following censored regression model in which one of the explanatory variables is a continuous endogenous variable:

$$\begin{aligned} y_{1i}^* &= \alpha y_{2i} + \mathbf{z}_{1i}' \boldsymbol{\beta} + u_i \\ y_{2i} &= \mathbf{z}_i' \boldsymbol{\pi} + \epsilon_i \\ y_{1i} &= \begin{cases} y_{1i}^* & \text{if } y_{1i}^* > 0 \\ 0 & \text{if } y_{1i}^* \leq 0 \end{cases} \end{aligned}$$

The exogenous explanatory variables are \mathbf{z}_{1i} , and the continuous endogenous explanatory variable is y_{2i} .

The likelihood function to maximize is

$$L = \prod_{i \in \{y_{1i} > 0\}} f(y_{1i}, y_{2i}) \cdot \prod_{i \in \{y_{1i} = 0\}} \int_{-\infty}^0 f(y_{1i}^*, y_{2i}) dy_{1i}^*$$

where $f(y_{1i}^*, y_{2i})$ is the joint density of y_{1i}^* and y_{2i} . Note that y_{1i} is substituted for y_{1i}^* when $y_{1i} > 0$. If you assume $(u_i, \epsilon_i) \stackrel{iid}{\sim} N(\mathbf{0}, \Sigma)$ with $\Sigma = \begin{bmatrix} \sigma_u^2 & \eta \\ \eta & \sigma_\epsilon^2 \end{bmatrix}$, then, by using $f(y_{1i}^*, y_{2i}) = f(y_{1i}^* | y_{2i}) \cdot f(y_{2i})$, you can write the likelihood function for each i as a multiplication of two parts. The first part is the probability density function of the normal distribution with mean $\mathbf{z}_i' \boldsymbol{\pi}$ and variance σ_ϵ^2 , and the second part follows a Tobit model that has latent mean $\alpha y_{2i} + \mathbf{z}_{1i}' \boldsymbol{\pi} + (\eta/\sigma_\epsilon^2)(y_{2i} - \mathbf{z}_{1i}' \boldsymbol{\pi})$ and variance $\sigma_u^2 - (\eta^2/\sigma_\epsilon^2)$. Then, you can obtain the log-likelihood function by taking the log of this multiplication and summing over i (for more information, see Wooldridge (2002, chapter 16)). This is the log-likelihood function that PROC QLIM maximizes. The parameters $(\hat{\alpha}, \hat{\boldsymbol{\beta}}, \hat{\boldsymbol{\pi}}, \hat{\sigma}_u^2, \hat{\sigma}_\epsilon^2, \hat{\eta})$ that are obtained from this maximization are the FIML estimators. Assuming that the latent model includes two instrumental variables and two exogenous explanatory variables, you can estimate this model in PROC QLIM as follows:

```
proc qlim data=a;
  model y1 = y2 z11 z12 / censored(lb=0);
  model y2 = z11 z12 z21 z22;
run;
```

For simple examples like the preceding ones, you can derive the likelihood function easily. However, as the number of endogenous explanatory variables increases, if these variables have a discontinuous nature, if simultaneity among equations exists, or if a combination of these occurs, then the derivation of the likelihood function becomes cumbersome, or, in some cases, the likelihood function does not even have a closed analytical form.

PROC QLIM can handle endogeneity regardless of the nature of the endogenous explanatory variables for a single structural model. In the case of one endogenous explanatory variable, PROC QLIM reports the FIML estimates that are calculated by using the analytical likelihood function that is obtained from the joint distribution of the dependent variable and the endogenous variable. When there is more than one endogenous explanatory variable, the analytical form of the likelihood function is usually not available; in this case PROC QLIM reports the simulated maximum likelihood estimates. For the simulated maximum likelihood estimation method, PROC QLIM uses the Geweke-Hajivassiliou-Keane (GHK) simulator (see, among others, Hajivassiliou, McFadden, and Ruud (1996)) to simulate the joint distribution of the dependent variable and the endogenous variables. The simulation is facilitated by assuming that the error terms in the latent models for the dependent variable and the endogenous explanatory variables are distributed as multivariate normal.

When you estimate a model in PROC QLIM, you can take the endogeneity into account by writing the structural model along with the reduced form models for each endogenous variable. Examples are provided in the following sections.

Probit Model with a Continuous Endogenous Explanatory Variable

Consider a probit model that contains a single endogenous explanatory variable in addition to two instruments and two exogenous explanatory variables. The model is

$$\begin{aligned} y_{1i}^* &= \alpha_1 y_{2i} + \beta_1 z_{1i} + \beta_2 z_{2i} + u_i \\ y_{2i}^* &= \pi_1 z_{1i} + \pi_2 z_{2i} + \pi_3 z_{3i} + \pi_4 z_{4i} + \epsilon_i \\ y_{1i} &= \begin{cases} 1 & \text{if } y_{1i}^* > 0 \\ 0 & \text{if } y_{1i}^* \leq 0 \end{cases} \\ y_{2i} &= y_{2i}^* \end{aligned}$$

where $\text{Cov}(u, \epsilon) = \eta$. You can estimate this model by using the following statements:

```
proc qlim data=a;
  model y1 = y2 z1 z2 / discrete;
  model y2 = z1 z2 z3 z4;
run;
```

Probit Model with a Binary Endogenous Explanatory Variable

Consider a probit model that contains a single binary endogenous explanatory variable in addition to two instruments and two exogenous explanatory variables. The model is

$$\begin{aligned}
 y_{1i}^* &= \alpha_1 y_{2i} + \beta_1 z_{1i} + \beta_2 z_{2i} + u_i \\
 y_{2i}^* &= \pi_1 z_{1i} + \pi_2 z_{2i} + \pi_3 z_{3i} + \pi_4 z_{4i} + \epsilon_i \\
 y_{1i} &= \begin{cases} 1 & \text{if } y_{1i}^* > 0 \\ 0 & \text{if } y_{1i}^* \leq 0 \end{cases} \\
 y_{2i} &= \begin{cases} 1 & \text{if } y_{2i}^* > 0 \\ 0 & \text{if } y_{2i}^* \leq 0 \end{cases}
 \end{aligned}$$

where $\text{Cov}(u, \epsilon) = \eta$. You can estimate this model by using the following statements:

```
proc qlim data=a;
  model y1 = y2 z1 z2 / discrete;
  model y2 = z1 z2 z3 z4 / discrete;
run;
```

Probit Model with a Censored Endogenous Explanatory Variable

Consider a probit model that contains a single censored (below zero) endogenous explanatory variable in addition to two instruments and two exogenous explanatory variables. The model is

$$\begin{aligned}
 y_{1i}^* &= \alpha_1 y_{2i} + \beta_1 z_{1i} + \beta_2 z_{2i} + u_i \\
 y_{2i}^* &= \pi_1 z_{1i} + \pi_2 z_{2i} + \pi_3 z_{3i} + \pi_4 z_{4i} + \epsilon_i \\
 y_{1i} &= \begin{cases} 1 & \text{if } y_{1i}^* > 0 \\ 0 & \text{if } y_{1i}^* \leq 0 \end{cases} \\
 y_{2i} &= \begin{cases} y_{2i}^* & \text{if } y_{2i}^* > 0 \\ 0 & \text{if } y_{2i}^* \leq 0 \end{cases}
 \end{aligned}$$

where $\text{Cov}(u, \epsilon) = \eta$. You can estimate this model by using the following statements:

```
proc qlim data=a;
  model y1 = y2 z1 z2 / discrete;
  model y2 = z1 z2 z3 z4 / censored(lb=0);
run;
```

Censored Regression Model with a Binary Endogenous Explanatory Variable

Consider a Type 1 Tobit model that contains a single binary endogenous explanatory variable in addition to two instruments and two exogenous explanatory variables. The model is

$$\begin{aligned}
 y_{1i}^* &= \alpha_1 y_{2i} + \beta_1 z_{1i} + \beta_2 z_{2i} + u_i \\
 y_{2i}^* &= \pi_1 z_{1i} + \pi_2 z_{2i} + \pi_3 z_{3i} + \pi_4 z_{4i} + \epsilon_i \\
 y_{1i} &= \begin{cases} y_{1i}^* & \text{if } y_{1i}^* > 0 \\ 0 & \text{if } y_{1i}^* \leq 0 \end{cases} \\
 y_{2i} &= \begin{cases} 1 & \text{if } y_{2i}^* > 0 \\ 0 & \text{if } y_{2i}^* \leq 0 \end{cases}
 \end{aligned}$$

where $\text{Cov}(u, \epsilon) = \eta$. You can estimate this model by using the following statements:

```
proc qlim data=a;
  model y1 = y2 z1 z2 / censored(lb=0);
  model y2 = z1 z2 z3 z4 / discrete;
run;
```

Censored Regression Model with Binary and Continuous Endogenous Explanatory Variables

Consider a Type 1 Tobit model that contain binary and continuous endogenous explanatory variables in addition to two instruments and two exogenous explanatory variables. The model is

$$\begin{aligned}
 y_{1i}^* &= \alpha_1 y_{21i} + \alpha_2 y_{22i} + \beta_1 z_{1i} + \beta_2 z_{2i} + u_i \\
 y_{21i}^* &= \pi_{11} z_{1i} + \pi_{12} z_{2i} + \pi_{13} z_{3i} + \pi_{14} z_{4i} + \epsilon_{1i} \\
 y_{22i}^* &= \pi_{21} z_{1i} + \pi_{22} z_{2i} + \pi_{23} z_{3i} + \pi_{24} z_{4i} + \epsilon_{2i} \\
 y_{1i} &= \begin{cases} y_{1i}^* & \text{if } y_{1i}^* > 0 \\ 0 & \text{if } y_{1i}^* \leq 0 \end{cases} \\
 y_{21i} &= \begin{cases} 1 & \text{if } y_{21i}^* > 0 \\ 0 & \text{if } y_{21i}^* \leq 0 \end{cases} \\
 y_{22i} &= y_{22i}^*
 \end{aligned}$$

where $\text{Cov}(u, \epsilon_1, \epsilon_2) = \eta$. You can estimate this model by using the following statements:

```
proc qlim data=a;
  model y1 = y21 y22 z1 z2 / censored(lb=0);
  model y21 = z1 z2 z3 z4 / discrete;
  model y22 = z1 z2 z3 z4;
run;
```

Probit Model with Binary, Censored, and Truncated Endogenous Explanatory Variables

Consider a probit model that contains binary, censored (below zero), and truncated (below zero) endogenous explanatory variables. The model is

$$\begin{aligned}
 y_{1i}^* &= \alpha_1 y_{21i} + \alpha_2 y_{22i} + \alpha_3 y_{23i} + u_i \\
 y_{21i}^* &= \pi_{11} z_{1i} + \pi_{12} z_{2i} + \pi_{13} z_{3i} + \pi_{14} z_{4i} + \epsilon_{1i} \\
 y_{22i}^* &= \pi_{21} z_{1i} + \pi_{22} z_{2i} + \pi_{23} z_{3i} + \pi_{24} z_{4i} + \epsilon_{2i} \\
 y_{23i}^* &= \pi_{31} z_{1i} + \pi_{32} z_{2i} + \pi_{33} z_{3i} + \pi_{34} z_{4i} + \epsilon_{3i} \\
 y_{1i} &= \begin{cases} 1 & \text{if } y_{1i}^* > 0 \\ 0 & \text{if } y_{1i}^* \leq 0 \end{cases} \\
 y_{21i} &= \begin{cases} 1 & \text{if } y_{21i}^* > 0 \\ 0 & \text{if } y_{21i}^* \leq 0 \end{cases} \\
 y_{22i} &= \begin{cases} y_{22i}^* & \text{if } y_{22i}^* > 0 \\ 0 & \text{if } y_{22i}^* \leq 0 \end{cases} \\
 y_{23i} &= y_{23i}^* \text{ if } y_{23i}^* > 0
 \end{aligned}$$

where z_1, \dots, z_4 are the instrumental variables that are independent of the errors. You can estimate this model by using the following statements:

```

proc qlim data=a;
  model y1 = y21 y22 y23 / discrete;
  model y21 = z1 z2 z3 z4 / discrete;
  model y22 = z1 z2 z3 z4 / censored(lb=0);
  model y23 = z1 z2 z3 z4 / truncated(lb=0);
run;

```

Note that the dependent variable y_1 should not occur in the models for the endogenous explanatory variables, because this causes inconsistent coefficient estimates. In other words, you should write the models for the endogenous explanatory variables as reduced form models. PROC QLIM does not handle simultaneous equations models.

Test for Endogeneity

PROC QLIM has two ways to test the null hypothesis that an endogenous explanatory variable (EEV) is in fact exogenous. In the case of a single EEV, the first testing method involves a likelihood ratio test of $H_0 : \rho = 0$. For example, consider the probit model with a binary endogenous explanatory variable that was considered earlier; y_2 is exogenous if the error term in the model for y_1^* is uncorrelated with the error term in the model for y_2^* . Therefore, testing to determine whether this correlation is 0 or not provides an endogeneity test for y_2 . You can do this in PROC QLIM as follows:

```

proc qlim data=a;
  model y1 = y2 z1 z2 / discrete;
  model y2 = z1 z2 z3 z4 / discrete;
  test _rho = 0 / LR;
run;

```

Failing to reject the null hypothesis favors the decision that y_2 is exogenous in the model for y_1 .

When there are two or more EEVs, the test becomes the joint likelihood ratio test of whether corresponding correlations are 0 or not.

The second testing method is similar to the approach of Rivers and Vuong (1988). Considering the same model, you can write

$$u_i = \theta\epsilon_i + e_i$$

where $\theta = \eta/\sigma_\epsilon^2$ and e is independent of z s and ϵ . You can now write

$$y_{1i}^* = \alpha_1 y_{2i} + \beta_1 z_{1i} + \beta_2 z_{2i} + \theta\epsilon_i + e_i$$

Testing $H_0 : \theta = 0$ is the same as testing whether u_i is correlated with ϵ_i or testing whether y_{2i} is endogenous or not. Because ϵ_i are unobserved, you can replace them with the OLS residuals from the model for y_{2i}^* and apply a robust t test. Note that even though y_{2i} is binary (or censored), the test is still correct under H_0 .

This approach can be summarized as a two-step procedure. In the first step, generated regressors—that is, the OLS residuals from the models for each of the EEVs—are obtained. In the second step, the structural model that includes the generated regressors as additional explanatory variables is estimated by the maximum likelihood method and the joint significance of these generated regressors is tested by the Wald test.

In PROC QLIM, you can apply the second method for the same test that was considered previously as follows:

```
proc qlim data=a;
  model y1 = y2 z1 z2 / discrete endotest(y2);
  model y2 = z1 z2 z3 z4 / discrete;
run;
```

Overidentification Test

In PROC QLIM you can test the validity of instrumental variables (IVs) by specifying the OVERID option in the ENDOGENOUS or MODEL statement. The OVERID test is a maximum likelihood version of the overidentifying restrictions test in the IV framework. If you have more IVs than are necessary for identification—that is, overidentifying IVs—you can use them to test the validity of your IVs. When you use the OVERID option to specify the overidentifying IVs, it applies the likelihood ratio test of the joint significance of these IVs, included as additional explanatory variables in the structural model that it estimates by the MLE jointly with the reduced form models. In effect, you test whether the overidentifying IVs are correlated with the error term in the structural model. You specify the reduced form models through the overidentifying IVs. The structural model is the model that includes the OVERID option. For example, consider the probit model that contains a continuous endogenous explanatory variable. You can consider z_3 or z_4 in the model for y_2 as an overidentifying IV; therefore, you can specify the OVERID test as follows:

```
proc qlim data=a;
  model y1 = y2 z1 z2 / discrete overid(y2.z4);
  model y2 = z1 z2 z3 z4;
run;
```

In this case, PROC QLIM estimates the structural model y_1 , including the overidentifying IV z_4 as an additional explanatory variable in this model, jointly with the reduced form model y_2 . Then it uses the likelihood ratio test to test the hypothesis that the overidentifying IV is insignificant. Rejecting this hypothesis raises doubts about the validity of the instruments z_3 and z_4 .

Note that, as long as you have continuous endogenous explanatory variables, the test result is invariant to which overidentifying IVs you specify in the test.

Bayesian Analysis

To perform Bayesian analysis, you must specify a BAYES statement. Unless otherwise stated, all options in this section are options in the BAYES statement.

By default, PROC QLIM uses the random walk Metropolis algorithm to obtain posterior samples. For the implementation details of the Metropolis algorithm in PROC QLIM, such as the blocking of the parameters and tuning of the covariance matrices, see the sections “[Blocking of Parameters](#)” on page 1577 and “[Tuning the Proposal Distribution](#)” on page 1577.

The Bayes theorem states that

$$p(\theta|y) \propto \pi(\theta)L(y|\theta)$$

where θ is a parameter or a vector of parameters and $\pi(\theta)$ is the product of the prior densities that are specified in the [PRIOR](#) statement. The term $L(y|\theta)$ is the likelihood associated with the [MODEL](#) statement.

Blocking of Parameters

In a multivariate parameter model, all the parameters are updated in one single block (by default or when you specify the `SAMPLING=MULTIMETROPOLIS` option). This could be inefficient, especially when parameters have vastly different scales. As an alternative, you could update the parameters one at the time (by specifying `SAMPLING=UNIMETROPOLIS`).

Tuning the Proposal Distribution

One key factor in achieving high efficiency of a Metropolis-based Markov chain is finding a good proposal distribution for each block of parameters. This process is called tuning. The tuning phase consists of a number of loops controlled by the options `MINTUNE` and `MAXTUNE`. The `MINTUNE=` option controls the minimum number of tuning loops and has a default value of 2. The `MAXTUNE=` option controls the maximum number of tuning loops and has a default value of 24. Each loop is iterated the number of times specified by the `NTU=` option, which has a default of 500. At the end of every loop, PROC QLIM examines the acceptance probability for each block. The acceptance probability is the percentage of NTU proposed values that have been accepted. If this probability does not fall within the acceptance tolerance range (see the following section), the proposal distribution is modified before the next tuning loop.

A good proposal distribution should resemble the actual posterior distribution of the parameters. Large sample theory states that the posterior distribution of the parameters approaches a multivariate normal distribution (see Gelman et al. 2004, Appendix B; Schervish 1995, Section 7.4). That is why a normal proposal distribution often works well in practice. The default proposal distribution in PROC QLIM is the normal distribution.

Scale Tuning

The acceptance rate is closely related to the sampling efficiency of a Metropolis chain. For a random walk Metropolis, a high acceptance rate means that most new samples occur right around the current data point. Their frequent acceptance means that the Markov chain is moving rather slowly and not exploring the parameter space fully. A low acceptance rate means that the proposed samples are often rejected; hence the chain is not moving much. An efficient Metropolis sampler has an acceptance rate that is neither too high nor too low. The scale c in the proposal distribution $q(\cdot|\cdot)$ effectively controls this acceptance probability. Roberts, Gelman, and Gilks (1997) show that if both the target and proposal densities are normal, the optimal acceptance probability for the Markov chain should be around 0.45 in a one-dimension problem and should asymptotically approach 0.234 in higher-dimension problems. The corresponding optimal scale is 2.38, which is the initial scale that is set for each block.

Because of the nature of stochastic simulations, it is impossible to fine-tune a set of variables so that the Metropolis chain has exactly the desired acceptance rate that you want. In addition, Roberts and Rosenthal (2001) empirically demonstrate that an acceptance rate between 0.15 and 0.5 is at least 80% efficient, so there is really no need to fine-tune the algorithms to reach an acceptance probability that is within a small tolerance of the optimal values. PROC QLIM works with a probability range, determined by `TargetAcceptance` ± 0.075 . If the observed acceptance rate in a given tuning loop is less than the lower bound of the range, the scale is reduced; if the observed acceptance rate is greater than the upper bound of the range, the scale is increased. During the tuning phase, a scale parameter in the normal distribution is adjusted as a function of the observed acceptance rate and the target acceptance rate. PROC QLIM uses the following updating scheme: ¹

$$c_{\text{new}} = \frac{c_{\text{cur}} \cdot \Phi^{-1}(p_{\text{opt}}/2)}{\Phi^{-1}(p_{\text{cur}}/2)}$$

where c_{cur} is the current scale, p_{cur} is the current acceptance rate, and p_{opt} is the optimal acceptance probability.

Covariance Tuning

To tune a covariance matrix, PROC QLIM takes a weighted average of the old proposal covariance matrix and the recent observed covariance matrix, based on the number samples (as specified by the `NTU=` option) NTU samples in the current loop. The formula to update the covariance matrix is:

$$\text{COV}_{\text{new}} = 0.75 \text{COV}_{\text{cur}} + 0.25 \text{COV}_{\text{old}}$$

There are two ways to initialize the covariance matrix:

- The default is an identity matrix that is multiplied by the initial scale of 2.38 and divided by the square root of the number of estimated parameters in the model. A number of tuning phases might be required before the proposal distribution is tuned to its optimal stage, because the Markov chain needs to spend time to learn about the posterior covariance structure. If the posterior variances of your parameters vary by more than a few orders of magnitude, if the variances of your parameters are much different from 1, or if the posterior correlations are high, then the proposal tuning algorithm might have difficulty forming an acceptable proposal distribution.

¹ Roberts and associates demonstrate that the relationship between acceptance probability and scale in a random walk Metropolis scheme is $p = 2\Phi(-\sqrt{I}c/2)$, where c is the scale, p is the acceptance rate, Φ is the CDF of a standard normal, and $I \equiv E_f[(f'(x)/f(x))^2]$, $f(x)$ is the density function of samples (Roberts, Gelman, and Gilks 1997; Roberts and Rosenthal 2001). This relationship determines the updating scheme, with I replaced by the identity matrix to simplify calculation.

- Alternatively, you can use a numerical optimization routine, such as the quasi-Newton method, to find a starting covariance matrix. The optimization is performed on the joint posterior distribution, and the covariance matrix is a quadratic approximation at the posterior mode. In some cases this is a better and more efficient way of initializing the covariance matrix. However, there are cases, such as when the number of parameters is large, where the optimization could fail to find a matrix that is positive definite. In those cases, the tuning covariance matrix is reset to the identity matrix.

A by-product of the optimization routine is that it also finds the maximum a posteriori (MAP) estimates with respect to the posterior distribution. The MAP estimates are used as the initial values of the Markov chain.

For more information, see the section “[INIT Statement](#)” on page 1547.

Initial Values of the Markov Chains

You can assign initial values to any parameters. (For more information, see the [INIT](#) statement.) If you use the optimization option [PROPCOV=](#), then PROC QLIM starts the tuning at the optimized values. This option overwrites the provided initial values. If you specify the [RANDINIT](#) option, the information that the [INIT](#) statement provides is overwritten.

Aggregation of Multiple Chains

When you want to exploit the possibility of running several MCMC instances at the same time ([NTRDS=n>1](#)), you face the problem of aggregating the chains. In ordinary applications, each MCMC instance can easily obtain stationary samples from the entire posterior distribution. In these applications, you can use the option [AGGREGATION=NOWEIGHTED](#). This option piles up one chain on top of another and makes no particular adjustment. However, when the posterior distribution is characterized by multiple distinct posterior modes, some of the MCMC instances fail to obtain stationary samples from the entire posterior distribution. You can use the option [AGGREGATION=WEIGHTED](#) when the posterior samples from each MCMC instance approximate well only a part of the posterior distribution.

The main idea behind the option [AGGREGATION=WEIGHTED](#) is to consider the entire posterior distribution to be similar to a mixture distribution. When you are sampling with multiple threads, each MCMC instance samples from one of the mixture components. Then the samples from each mixture component are aggregated together using a resampling scheme in which weights are proportional to the nonnormalized posterior distribution.

Description of the Algorithm

The preliminary step of the aggregation that is implied by the option [AGGREGATION=WEIGHTED](#) is to run several (K) independent instances of the MCMC algorithm. Each instance searches for a set of stationary samples. Notice that the concept of stationarity is weaker: each instance might be able to explore not the entire posterior but only portions of it. In the next equation, each column represents the output from one MCMC instance,

$$\begin{pmatrix} x_{11} \\ x_{21} \\ \dots \\ x_{n1} \end{pmatrix} \begin{pmatrix} x_{12} \\ x_{22} \\ \dots \\ x_{n2} \end{pmatrix} \dots \begin{pmatrix} x_{1K} \\ x_{2K} \\ \dots \\ x_{nK} \end{pmatrix} \sim \text{globally/locally sampled from the posterior}$$

If the length of each chain is less than n , you can augment the the corresponding chain by subsampling the chain itself. Each chain is then sorted with respect to the nonnormalized posterior density: $\pi(x_{[1].}) \leq \pi(x_{[2].}) \leq \dots \leq \pi(x_{[n].})$. Therefore,

$$\begin{pmatrix} x_{11} \\ x_{21} \\ \dots \\ x_{n1} \end{pmatrix} \begin{pmatrix} x_{12} \\ x_{22} \\ \dots \\ x_{n2} \end{pmatrix} \dots \begin{pmatrix} x_{1K} \\ x_{2K} \\ \dots \\ x_{nK} \end{pmatrix} \rightarrow \begin{pmatrix} x_{[1]1} \\ x_{[2]1} \\ \dots \\ x_{[n]1} \end{pmatrix} \begin{pmatrix} x_{[1]2} \\ x_{[2]2} \\ \dots \\ x_{[n]2} \end{pmatrix} \dots \begin{pmatrix} x_{[1]K} \\ x_{[2]K} \\ \dots \\ x_{[n]K} \end{pmatrix}$$

The final step is to use a multinomial sampler to resample each row i with weights proportional to the nonnormalized posterior densities:

$$\tilde{x}_{(i-1)K+1}, \tilde{x}_{(i-1)K+2}, \dots, \tilde{x}_{(i-1)K+K} \sim \text{Multinom} [x_{[i]1}, x_{[i]2}, \dots, x_{[i]K}; \pi(x_{[i]1}), \pi(x_{[i]2}), \dots, \pi(x_{[i]K})]$$

The resulting posterior sample

$$\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_K, \dots, \tilde{x}_{(i-1)K+1}, \tilde{x}_{(i-1)K+2}, \dots, \tilde{x}_{(i-1)K+K}, \dots, \tilde{x}_{(n-1)K+1}, \tilde{x}_{(n-1)K+2}, \dots, \tilde{x}_{nK}$$

is a good approximation of the posterior distribution that is characterized by multiple modes.

Automated Initialization of MCMC

The MCMC methods can generate samples from the posterior distribution. The correct implementation of these methods often requires the stationarity analysis, the convergence analysis and the accuracy analysis of the posterior samples. These analyses usually imply the following:

- initialization of the proposal distribution
- initialization of the chains (starting values)
- determination of the burn-in
- determination of the length of the chains.

In more general terms, this determination is equivalent to deciding whether the samples are drawn from the posterior distribution (stationarity analysis), and whether the number of samples is large enough to accurately approximate the posterior distribution (accuracy analysis). You can use the AUTOMCMC option to automate and facilitate the stationary analysis and the accuracy analysis.

Description of the Algorithm

The algorithm consists of two phases. In the first phase, the stationarity phase, the algorithm tries to generate stationary samples from the posterior distribution. In the second phase, the accuracy phase, the algorithm searches for an accurate representation of the posterior distribution. The algorithm implements the following tools:

- Geweke test to check stationarity
- Heidelberger-Welch test to check stationarity and provide a proxy for the burn-in
- Heidelberger-Welch half-test to check the accuracy of the posterior mean


- Raftery-Lewis test to check the accuracy of a given percentile (indirectly proving a proxy for the number of required samples)
- effective sample size analysis to determine a proxy of the number of required samples

During the stationarity phase, the algorithm searches for stationarity. The number of attempts that the algorithm makes is determined by the option `ATTEMPTS=number`. During each attempt, a preliminary tuning stage chooses a proposal distribution for the MCMC sampler. At the end of the preliminary tuning phase, the algorithm analyzes tests for the stationarity of the samples. If the percentage of successful stationary tests is equal to or greater than the percentage that is indicated by the option `TOL=value`, then the posterior sample is considered to be stationary. If the sample cannot be considered stationary, then the algorithm attempts to achieve stationarity by changing some of the initialization parameters as follows:

- increasing the number of tuning samples (NTU)
- increasing the number of posterior samples (NMC)
- increasing the burn-in (NBI)

Figure 22.8 shows a flowchart of the algorithm as it searches for stationarity.

Figure 22.8 Flowchart of the AUTOMCMC Algorithm: Stationarity Analysis



You can initialize NMC=M, NBI=B, and NTU=T during the stationarity phase by specifying NMC, NBI, and NTU as options in the BAYES statement. You can also change the minimum stationarity acceptance ratio of successful stationarity tests that are needed to exit the stationarity phase. By default, TOL=0.95. For example:

```
proc qlim data=dataset;
  ...;
  bayes nmc=M nbi=B ntu=T automcmc=( stationarity=(tol=0.95) );
  ...;
run;
```


During the accuracy phase, the algorithm attempts to determine how many posterior samples are needed. The number of attempts is determined by the option ATTEMPTS=*number*. You can choose between two different approaches to study the accuracy:

- accuracy analysis based on the effective sample size (ESS)
- accuracy analysis based on the Heidelberger-Welch half-test and the Raftery-Lewis test

If you choose the effective sample size approach, you must provide the minimum number of effective samples that are needed. You can also change the tolerance for the ESS accuracy analysis (by default, TOL=0.95). For example:

```
proc qlim data=dataset;
  ...;
  bayes automcmc=(targetess=N accuracy=(tol=0.95));
  ...;
run;
```

Figure 22.9 shows a flowchart of the algorithm based on the effective sample size approach to determine whether the samples provide an accurate representation of the posterior distribution.

Figure 22.9 Flowchart of the AUTOMCMC Algorithm: Accuracy Analysis Based on the ESS

If you choose the accuracy analysis based on the Heidelberg-Welch half-test and the Raftery-Lewis test (the default option), then you might want to choose a posterior quantile of interest for the Raftery-Lewis test (by default, 0.025). You can also change the tolerance for the accuracy analysis (by default, $TOL=0.95$). Notice that the Raftery-Lewis test produces a proxy of the number of posterior sample required. In each attempt, the current number of posterior samples is compared to this proxy. If the proxy is greater than the current nmc, then the algorithm reinitializes itself. To control this reinitialization, you can use the option `RLLIMITS=(LB=lb UB=ub)`. In particular, there are three cases

- if the proxy is greater than `ub`, then NMC is set equal to `ub`,
- if the proxy is less than `lb`, then NMC is set equal to `lb`,
- if `lb` is less than the proxy, which is less than `ub`, then NMC is set equal to the proxy.

For example:


```

proc qlim data=dataset;
  ...;
  bayes raftery(q=0.025) automcmc=( accuracy=(tol=0.95 targetstats=(rllimits=(lb=k1 ub=k2))
  ...;
run;

```

Figure 22.10 shows a flowchart of the algorithm based on the Heidelberger-Welch half-test and the Raftery-Lewis test approach to determine whether the posterior samples provide an accurate representation of the posterior distribution.

Figure 22.10 Flowchart of the AUTOMCMC Algorithm: Accuracy Analysis Based on the Heidelberger-Welch Half-test and the Raftery-Lewis Test



Prior Distributions

The PRIOR statement is used to specify the prior distribution of the model parameters. You must specify a list of parameters, a tilde \sim , and then a distribution with its parameters. You can specify multiple PRIOR statements to define independent priors. Parameters that are associated with a regressor variable are referred to by the name of the corresponding regressor variable.

You can specify the special keyword `_REGRESSORS` to consider all the regressors of a model. If multiple prior statements affect the same parameter, the prior that is specified is used. For example, in a regression with three regressors (X1, X2, X3) the following statements imply that the prior on X1 is `NORMAL(MEAN=0, VAR=1)`, the prior on X2 is `GAMMA(SHAPE=3, SCALE=4)`, and the prior on X3 is `UNIFORM(MIN=0, MAX=1)`:

```
...
prior _Regressors ~ uniform(min=0, max=1);
```

```
prior X1 X2 ~ gamma(shape=3, scale=4);
prior X1 ~ normal(mean=0, var=1);
...
```

If a parameter is not associated with a PRIOR statement or if some of the prior hyperparameters are missing, then the following default choices are considered:

Table 22.2 Default values for prior distributions.

PRIOR distribution	Hyperparameter ₁	Hyperparameter ₂	Min	Max	Parameters Default Choice
NORMAL	MEAN=0	VAR=1E6	$-\infty$	∞	Regression-Location-Threshold Scale
IGAMMA	SHAPE=2.000001	SCALE=1	> 0	∞	
GAMMA	SHAPE=1	SCALE=1	0	∞	
UNIFORM			$-\infty$	∞	Cross-correlation
UNIFORM			> -1	< 1	
BETA	SHAPE1=1	SHAPE2=1	$-\infty$	∞	
T	LOCATION=0	DF=3	$-\infty$	∞	

See the section “[Standard Distributions](#)” on page 1588 for density specification.

Priors for Heteroscedastic Models

The choice of the prior distribution for a heteroscedastic model is particularly interesting. Based on the notation provided in section “[HETERO Statement](#)” on page 1546, you need to provide a prior for γ . This prior is enough to induce different σ_i^2 into the analysis. The resulting inference is a compromise between two cases: the inference based on the entire sample and the inference based on a single unit \mathbf{z}_i . The degree of compromise is determined by $\pi(\gamma)$.

This type of modeling is similar to a method called “hierarchical Bayes,” in which the prior is characterized by two levels: one for each individual $\pi(\sigma_i^2|\gamma)$ and one for the entire population $\pi(\gamma)$. In this scenario the degree of compromise between the information provided by a unit and the information provided by the entire sample is determined by the data.

The choice of the prior might not be straightforward, and it can heavily affect sampling performance. Depending on how the heteroscedastic effects are modeled, the default priors are

$$\begin{aligned}
 &\text{if } [1 + \exp(\mathbf{z}_i' \gamma)], \quad \pi(\gamma_j) = \mathbf{normal} \left\{ \text{mean} = \frac{1}{\bar{z}_j J} \left[\log \left(\frac{\varepsilon^4}{1 + \varepsilon^2} \right) \right], \text{var} = \frac{1}{\bar{z}_j^2 J} \left[\log \left(\frac{1 + \varepsilon^2}{\varepsilon^2} \right) \right] \right\} \\
 &\text{if } [\exp(\mathbf{z}_i' \gamma)], \quad \pi(\gamma_j) = \mathbf{normal} \left\{ \text{mean} = \frac{1}{\bar{z}_j J} \left[\log \left(\frac{1}{2} \right) \right], \text{var} = \frac{1}{\bar{z}_j^2 J} [\log(2)] \right\} \\
 &\text{if } (1 + \mathbf{z}_i' \gamma), \quad \pi(\gamma_j) = \mathbf{normal} \left\{ \text{mean} = 0, \text{var} = \frac{1}{\bar{z}_j^2 J} \right\} \\
 &\text{if } [1 + (\mathbf{z}_i' \gamma)^2], \quad \pi(\gamma_j) = \mathbf{normal} \left\{ \text{mean} = \frac{(\varepsilon^2 - 1/2)^{1/4}}{\bar{z}_j J}, \text{var} = \frac{\varepsilon - (\varepsilon^2 - 1/2)^{1/2}}{\bar{z}_j^2 J} \right\}
 \end{aligned}$$

where $\bar{z}_j = \frac{1}{n} \sum_{i=1}^n z_{ij}$, $\forall j$, and ε is a small number (by default, $\varepsilon = 0.1$ for the EXPONENTIAL link function and $\varepsilon = 0.71$ for the QUADRATIC link function).

The priors for the EXPONENTIAL and QUADRATIC link functions are not straightforward. To understand the choices, do the following:

1. Assume that

$$\mathbf{z}_i' \boldsymbol{\gamma} = z_{i1}\gamma_1 + \dots + z_{iJ}\gamma_J \approx \bar{z}_1\gamma_1 + \dots + \bar{z}_J\gamma_J, \quad \forall i$$

2. Set the priors according to the link function type:

- For the EXPONENTIAL link function, set

$$\begin{aligned} \mathrm{E} \left[\exp(\mathbf{z}_i' \boldsymbol{\gamma}) \right] &\approx \mathrm{E} [\exp(\bar{z}_1\gamma_1)] \times \dots \times \mathrm{E} [\exp(\bar{z}_J\gamma_J)] = \varepsilon \\ \mathrm{V} \left[\exp(\mathbf{z}_i' \boldsymbol{\gamma}) \right] &\approx \mathrm{E} [\exp(2\bar{z}_1\gamma_1)] \times \dots \times \mathrm{E} [\exp(2\bar{z}_J\gamma_J)] - \varepsilon^2 = 1 \end{aligned}$$

Assume a normal prior for $\pi(\gamma_j)$, and set

$$\begin{aligned} \mathrm{E} [\exp(\bar{z}_j\gamma_j)] &= \varepsilon^{\frac{1}{J}}, \forall j \\ \mathrm{E} [\exp(2\bar{z}_j\gamma_j)] &= (1 + \varepsilon^2)^{\frac{1}{J}}, \forall j \end{aligned}$$

Based on the properties of the lognormal distribution, the prior hyperparameters for γ_j can be derived. Notice that J is the number of regressors that are used in the heterogeneous regression. If the intercept is excluded, then $\varepsilon = 1$.

- For the QUADRATIC link function, set

$$\begin{aligned} \mathrm{E} \left[(\mathbf{z}_i' \boldsymbol{\gamma})^2 \right] &\approx [\mathrm{E} (\bar{z}_1\gamma_1 + \dots + \bar{z}_J\gamma_J)]^2 + \mathrm{V} [\bar{z}_1\gamma_1 + \dots + \bar{z}_J\gamma_J] = \varepsilon \\ \mathrm{V} \left[(\mathbf{z}_i' \boldsymbol{\gamma})^2 \right] &\approx \mathrm{E} [(\bar{z}_1\gamma_1 + \dots + \bar{z}_J\gamma_J)^4] - \varepsilon^2 = 1 \end{aligned}$$

Assume a normal prior for $\pi(\gamma_j)$. Based on the properties of the normal distribution, the preceding expressions return

$$\begin{aligned} \mathrm{E} [\bar{z}_1\gamma_1 + \dots + \bar{z}_J\gamma_J] &= (\varepsilon^2 - 1/2)^{1/4} \\ \mathrm{V} [\bar{z}_1\gamma_1 + \dots + \bar{z}_J\gamma_J] &= \varepsilon - (\varepsilon^2 - 1/2)^{1/2} \\ \varepsilon &> (1/2)^{1/2} \end{aligned}$$

The prior hyperparameters for γ_j can be derived by setting

$$\begin{aligned} \mathrm{E} [\bar{z}_j\gamma_j] &= \frac{(\varepsilon^2 - 1/2)^{1/4}}{J}, \forall j \\ \mathrm{V} [\bar{z}_j\gamma_j] &= \frac{\varepsilon - (\varepsilon^2 - 1/2)^{1/2}}{J}, \forall j \end{aligned}$$

Notice that J is the number of regressors that are used in the heterogeneous regression. It is important to emphasize that the restriction $\varepsilon > (1/2)^{1/2} \approx 0.71$ is likely to introduce some distortion because ε cannot be any “small” number.

Automated MCMC

The main purpose is to provide the user with the opportunity of obtaining a good approximation of the posterior distribution without initializing the MCMC algorithm: initial values, proposal distributions, burn-in and number of samples.

The automated algorithm is composed of two phases: tuning and sampling. In the tuning phase, there are two main concerns: the choice of a good proposal distribution and the search for the stationary region of the posterior distribution. In the sampling phase, the algorithm will decide how many samples are necessary to obtain good approximations of the posterior mean and some quantiles of interest.

Tuning Phase

During the tuning phase, the algorithm tries to search for a good proposal distribution and, at the same time, to reach the stationary region of the posterior. The choice of the proposal distribution is based on the analysis of the acceptance rates. This is similar to what is done in PROC MCMC: for more details see Chapter 61.10, “Tuning the Proposal Distribution” (*SAS/STAT User’s Guide*). For the stationarity analysis, the main idea is to run two tests, Geweke (Ge) and Heidelberger-Welch (HW), on the posterior chains at the end of each attempt. For more details, see Chapter 7.4, “Geweke Diagnostics” (*SAS/STAT User’s Guide*), and Chapter 7.4, “Heidelberger and Welch Diagnostics” (*SAS/STAT User’s Guide*). If the stationarity hypothesis is rejected, then the tuning samples are increased and the tests repeated in the next attempt. After 10 attempts, the tuning phase will be ended regardless of the results. The tuning parameters for the first attempt are fixed:

0 burn-in (nbi),
1000 tuning samples (ntu),
10000 MCMC samples (nmc).

For the remaining attempts, the tuning parameters will be adjusted dynamically. More specifically, each parameter will be assigned an acceptance ratio (AR) of the stationarity hypothesis:

$AR_i = 0$ if both tests reject the stationarity hypothesis,
 $AR_i = 0.5$ if one tests rejects and the other does not,
 $AR_i = 1$ if both tests do not reject the stationarity hypothesis,

for $i = 1, \dots, k$. Then, an overall stationarity average (SA) for all parameters ratios is evaluated,

$$SA = \sum_{i=1}^k \frac{AR_i}{k}, \quad (22.1)$$

and the number of tuning samples is updated accordingly:

$ntu = ntu + 2000$ if $SA < 70\%$,
 $ntu = ntu + 1000$ if $70\% \leq SA < 100\%$,
 $ntu = ntu$ if $SA = 100\%$.

The Heidelberger-Welch test also provides an indications of how much burn-in should be used. The algorithm requires this burn-in to be: $nbi(HW) = 0$. If that is not the case, the burn-in will updated accordingly, $nbi = nbi + nbi(HW)$, and a new tuning attempt will be implemented. This choice is motivated by the fact that the burn-in must be discarded in order to reach the stationary region of the posterior distribution.

The number of samples predicted by the Raftery-Lewis $nmc(RL)$ diagnostic tool will be also taken into consideration: $nmc = nmc + nmc(RL)$. For more details see Chapter 7.4, “Raftery and Lewis Diagnostics” (*SAS/STAT User’s Guide*). However, in order to exit the tuning phase, it will not be required $nmc(RL) = 0$.

Sampling Phase

The main idea of the sampling phase is to make sure that the mean and a quantile of interest are evaluated accurately. This can be tested by implementing the half-width test by Heidelberger-Welch and by analyzing the Raftery-Lewis diagnostic tool. In addition, the requirements defined in the tuning phase will also be checked: the Geweke and the Heidelberger-Welch tests must not reject the stationary hypothesis and the burn-in predicted by the Heidelberger-Welch test must be zero.

The sampling phase is characterized by a maximum of 10 attempts. If the algorithm exceeds this limit, the sampling phase will end and indications on how to improve sampling will be given. The sampling phase will first update the burn-in with the information provided by the HW test: $nbi = nbi + nbi(HW)$. Then, it determines the difference between the actual number of samples and the number of samples predicted by the RL test: $\Delta[nmc(RL)] = nmc(RL) - nmc$. The new number of samples will be updated accordingly:

$$\begin{aligned} nmc &= nmc + 1000 && \text{if } 0 < \Delta[nmc(RL)] \leq 10000, \\ nmc &= nmc + \Delta[nmc(RL)] && \text{if } 10000 < \Delta[nmc(RL)] \leq 300000, \\ nmc &= nmc + 300000 && \text{if } 300000 < \Delta[nmc(RL)]. \end{aligned}$$

Finally, the HW half-width test checks whether the sample mean is accurate. If the mean of any parameters is not considered to be accurate, the number of sampling is increased:

$$\begin{aligned} nmc &= nmc + \{10000 - \Delta[nmc(RL)]\} && \text{if } 10000 - \Delta[nmc(RL)] \geq 0, \\ nmc &= nmc && \text{if } 10000 - \Delta[nmc(RL)] < 0. \end{aligned}$$

Standard Distributions

Table 22.3 through Table 22.8 show all the distribution density functions that PROC QLIM recognizes. You specify these distribution densities in the **PRIOR** statement.

Table 22.3 Beta Distribution

PRIOR statement	BETA(SHAPE1= a , SHAPE2= b , MIN= m , MAX= M)
	Note: Commonly $m = 0$ and $M = 1$.
Density	$\frac{(\theta-m)^{a-1}(M-\theta)^{b-1}}{B(a,b)(M-m)^{a+b-1}}$
Parameter restriction	$a > 0, \quad b > 0, \quad -\infty < m < M < \infty$
Range	$\begin{cases} [m, M] & \text{when } a = 1, b = 1 \\ [m, M) & \text{when } a = 1, b \neq 1 \\ (m, M] & \text{when } a \neq 1, b = 1 \\ (m, M) & \text{otherwise} \end{cases}$
Mean	$\frac{a}{a+b} \times (M - m) + m$
Variance	$\frac{ab}{(a+b)^2(a+b+1)} \times (M - m)^2$

Mode	$\begin{cases} \frac{a-1}{a+b-2} \times M + \frac{b-1}{a+b-2} \times m & a > 1, b > 1 \\ m \text{ and } M & a < 1, b < 1 \\ m & \begin{cases} a < 1, b \geq 1 \\ a = 1, b > 1 \end{cases} \\ M & \begin{cases} a \geq 1, b < 1 \\ a > 1, b = 1 \end{cases} \\ \text{not unique} & a = b = 1 \end{cases}$
Defaults	SHAPE1=SHAPE2=1, MIN $\rightarrow -\infty$, MAX $\rightarrow \infty$

Table 22.4 Gamma Distribution

PRIOR statement	GAMMA(SHAPE= a , SCALE= b)
Density	$\frac{1}{b^a \Gamma(a)} \theta^{a-1} e^{-\theta/b}$
Parameter restriction	$a > 0, b > 0$
Range	$[0, \infty)$
Mean	ab
Variance	ab^2
Mode	$(a - 1)b$
Defaults	SHAPE=SCALE=1

Table 22.5 Inverse-Gamma Distribution

PRIOR statement	IGAMMA(SHAPE= a , SCALE= b)
Density	$\frac{b^a}{\Gamma(a)} \theta^{-(a+1)} e^{-b/\theta}$
Parameter restriction	$a > 0, b > 0$
Range	$0 < \theta < \infty$
Mean	$\frac{b}{a-1}, \quad a > 1$
Variance	$\frac{b^2}{(a-1)^2(a-2)}, \quad a > 2$
Mode	$\frac{b}{a+1}$
Defaults	SHAPE=2.000001, SCALE=1

Table 22.6 Normal Distribution

PRIOR statement	NORMAL(MEAN= μ , VAR= σ^2)
Density	$\frac{1}{\sigma \sqrt{2\pi}} \exp\left(-\frac{(\theta-\mu)^2}{2\sigma^2}\right)$
Parameter restriction	$\sigma^2 > 0$

Range	$-\infty < \theta < \infty$
Mean	μ
Variance	σ^2
Mode	μ
Defaults	MEAN=0, VAR=1000000

Table 22.7 t Distribution

PRIOR statement	T(LOCATION= μ , DF= ν)
Density	$\frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})\sqrt{\pi\nu}} \left[1 + \frac{(\theta-\mu)^2}{\nu} \right]^{-\frac{\nu+1}{2}}$
Parameter restriction	$\nu > 0$
Range	$-\infty < \theta < \infty$
Mean	μ , for $\nu > 1$
Variance	$\frac{\nu}{\nu-2}$, for $\nu > 2$
Mode	μ
Defaults	LOCATION=0, DF=3

Table 22.8 Uniform Distribution

PRIOR statement	UNIFORM(MIN= m , MAX= M)
Density	$\frac{1}{M-m}$
Parameter restriction	$-\infty < m < M < \infty$
Range	$\theta \in [m, M]$
Mean	$\frac{m+M}{2}$
Variance	$\frac{(M-m)^2}{12}$
Mode	Not unique
Defaults	MIN $\rightarrow -\infty$, MAX $\rightarrow \infty$

Output to SAS Data Set

XBeta, Predicted, Residual

XBeta is the structural part on the right-hand side of the model. Predicted value is the predicted dependent variable value. For censored variables, if the predicted value is outside the boundaries, it is reported as the

closest boundary. For discrete variables, it is the level whose boundaries $X\beta$ falls between. Residual is defined only for continuous variables and is defined as

$$Residual = Observed - Predicted$$

Error Standard Deviation

Error standard deviation is σ_i in the model. It varies only when the HETERO statement is used.

Marginal Effects

Marginal effect is defined as a contribution of one control variable to the response variable. For the binary choice model with two response categories, $\mu_0 = -\infty$, $\mu_1 = 0$, $\mu_2 = \infty$; and ordinal response model with M response categories, μ_0, \dots, μ_M , define

$$R_{i,j} = \mu_j - \mathbf{x}_i' \boldsymbol{\beta}$$

The probability that the unobserved dependent variable is contained in the j th category can be written as

$$P[\mu_{j-1} < y_i^* \leq \mu_j] = F(R_{i,j}) - F(R_{i,j-1})$$

The marginal effect of changes in the regressors on the probability of $y_i = j$ is then

$$\frac{\partial Prob[y_i = j]}{\partial \mathbf{x}} = [f(\mu_{j-1} - \mathbf{x}_i' \boldsymbol{\beta}) - f(\mu_j - \mathbf{x}_i' \boldsymbol{\beta})] \boldsymbol{\beta}$$

where $f(x) = \frac{dF(x)}{dx}$. In particular,

$$f(x) = \frac{dF(x)}{dx} = \begin{cases} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} & (\text{probit}) \\ \frac{e^{-x}}{[1+e^{(-x)}]^2} & (\text{logit}) \end{cases}$$

The marginal effects in the Box-Cox regression model are

$$\frac{\partial E[y_i]}{\partial \mathbf{x}} = \boldsymbol{\beta} \frac{x^{\lambda_k-1}}{y^{\lambda_0-1}}$$

The marginal effects in the truncated regression model are

$$\frac{\partial E[y_i | L_i < y_i^* < R_i]}{\partial \mathbf{x}} = \boldsymbol{\beta} \left[1 - \frac{(\phi(a_i) - \phi(b_i))^2}{(\Phi(b_i) - \Phi(a_i))^2} + \frac{a_i \phi(a_i) - b_i \phi(b_i)}{\Phi(b_i) - \Phi(a_i)} \right]$$

where $a_i = \frac{L_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma_i}$ and $b_i = \frac{R_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma_i}$.

The marginal effects in the censored regression model are

$$\frac{\partial E[y | \mathbf{x}_i]}{\partial \mathbf{x}} = \boldsymbol{\beta} \times Prob[L_i < y_i^* < R_i]$$

Inverse Mills Ratio, Expected and Conditionally Expected Values

Expected and conditionally expected values are computed only for continuous variables. The inverse Mills ratio is computed for censored or truncated continuous, binary discrete, and selection endogenous variables.

Let L_i and R_i be the lower boundary and upper boundary, respectively, for the y_i . Define $a_i = \frac{L_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma_i}$ and $b_i = \frac{R_i - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma_i}$. Then the inverse Mills ratio is defined as

$$\lambda = \frac{(\phi(a_i) - \phi(b_i))}{(\Phi(b_i) - \Phi(a_i))}$$

for a continuous variable and defined as

$$\lambda = \frac{\phi(\mathbf{x}_i' \boldsymbol{\beta})}{\Phi(\mathbf{x}_i' \boldsymbol{\beta})}$$

for a binary discrete variable.

The expected value is the unconditional expectation of the dependent variable. For a censored variable, it is

$$E[y_i] = \Phi(a_i)L_i + (\mathbf{x}_i' \boldsymbol{\beta} + \lambda\sigma_i)(\Phi(b_i) - \Phi(a_i)) + (1 - \Phi(b_i))R_i$$

For a left-censored variable ($R_i = \infty$), this formula is

$$E[y_i] = \Phi(a_i)L_i + (\mathbf{x}_i' \boldsymbol{\beta} + \lambda\sigma_i)(1 - \Phi(a_i))$$

where $\lambda = \frac{\phi(a_i)}{1 - \Phi(a_i)}$.

For a right-censored variable ($L_i = -\infty$), this formula is

$$E[y_i] = (\mathbf{x}_i' \boldsymbol{\beta} + \lambda\sigma_i)\Phi(b_i) + (1 - \Phi(b_i))R_i$$

where $\lambda = -\frac{\phi(b_i)}{\Phi(b_i)}$.

For a uncensored variable, this formula is

$$E[y_i] = \mathbf{x}_i' \boldsymbol{\beta}$$

The conditional expected value is the expectation given that the variable is inside the boundaries:

$$E[y_i | L_i < y_i < R_i] = \mathbf{x}_i' \boldsymbol{\beta} + \lambda\sigma_i$$

Probability

Probability applies only to discrete responses. It is the marginal probability that the discrete response is taking the value of the observation. If the PROBALL option is specified, then the probability for all of the possible responses of the discrete variables is computed.

Technical Efficiency

Technical efficiency for each producer is computed only for stochastic frontier models.

In general, the stochastic production frontier can be written as

$$y_i = f(x_i; \beta) \exp\{v_i\} TE_i$$

where y_i denotes producer i 's actual output, $f(\cdot)$ is the deterministic part of production frontier, $\exp\{v_i\}$ is a producer-specific error term, and TE_i is the technical efficiency coefficient, which can be written as

$$TE_i = \frac{y_i}{f(x_i; \beta) \exp\{v_i\}}.$$

In the case of a Cobb-Douglas production function, $TE_i = \exp\{-u_i\}$. See the section “[Stochastic Frontier Production and Cost Models](#)” on page 1561.

Cost frontier can be written in general as

$$E_i = c(y_i, w_i; \beta) \exp\{v_i\} / CE_i$$

where w_i denotes producer i 's input prices, $c(\cdot)$ is the deterministic part of cost frontier, $\exp\{v_i\}$ is a producer-specific error term, and CE_i is the cost efficiency coefficient, which can be written as

$$CE_i = \frac{c(x_i, w_i; \beta) \exp\{v_i\}}{E_i}$$

In the case of a Cobb-Douglas cost function, $CE_i = \exp\{-u_i\}$. See the section “[Stochastic Frontier Production and Cost Models](#)” on page 1561. Hence, both technical and cost efficiency coefficients are the same. The estimates of technical efficiency are provided in the following subsections.

Normal-Half Normal Model

Define $\mu_* = -\epsilon\sigma_u^2/\sigma^2$ and $\sigma_*^2 = \sigma_u^2\sigma_v^2/\sigma^2$. Then, as it is shown by Jondrow et al. (1982), conditional density is as follows:

$$f(u|\epsilon) = \frac{f(u, \epsilon)}{f(\epsilon)} = \frac{1}{\sqrt{2\pi}\sigma_*} \exp\left\{-\frac{(u - \mu_*)^2}{2\sigma_*^2}\right\} \Bigg/ \left[1 - \Phi\left(-\frac{\mu_*}{\sigma_*}\right)\right]$$

Hence, $f(u|\epsilon)$ is the density for $N^+(\mu_*, \sigma_*^2)$.

Using this result, it follows that the estimate of technical efficiency (Battese and Coelli 1988) is

$$TE1_i = E(\exp\{-u_i\}|\epsilon_i) = \left[\frac{1 - \Phi(\sigma_* - \mu_{*i}/\sigma_*)}{1 - \Phi(-\mu_{*i}/\sigma_*)}\right] \exp\left\{-\mu_{*i} + \frac{1}{2}\sigma_*^2\right\}$$

The second version of the estimate (Jondrow et al. 1982) is

$$TE2_i = \exp\{-E(u_i|\epsilon_i)\}$$

where

$$E(u_i|\epsilon_i) = \mu_{*i} + \sigma_* \left[\frac{\phi(-\mu_{*i}/\sigma_*)}{1 - \Phi(-\mu_{*i}/\sigma_*)} \right] = \sigma_* \left[\frac{\phi(\epsilon_i \lambda / \sigma)}{1 - \Phi(\epsilon_i \lambda / \sigma)} - \left(\frac{\epsilon_i \lambda}{\sigma} \right) \right]$$

Normal-Exponential Model

Define $A = -\tilde{\mu}/\sigma_v$ and $\tilde{\mu} = -\epsilon - \sigma_v^2/\sigma_u$. Then, as it is shown by Kumbhakar and Lovell (2000), conditional density is as follows:

$$f(u|\epsilon) = \frac{1}{\sqrt{2\pi}\sigma_v\Phi(-\tilde{\mu}/\sigma_v)} \exp\left\{-\frac{(u - \tilde{\mu})^2}{2\sigma^2}\right\}$$

Hence, $f(u|\epsilon)$ is the density for $N^+(\tilde{\mu}, \sigma_v^2)$.

Using this result, it follows that the estimate of technical efficiency is

$$TE1_i = E(\exp\{-u_i\}|\epsilon_i) = \left[\frac{1 - \Phi(\sigma_v - \tilde{\mu}_i/\sigma_v)}{1 - \Phi(-\tilde{\mu}_i/\sigma_v)} \right] \exp\left\{-\tilde{\mu}_i + \frac{1}{2}\sigma_v^2\right\}$$

The second version of the estimate is

$$TE2_i = \exp\{-E(u_i|\epsilon_i)\}$$

where

$$E(u_i|\epsilon_i) = \tilde{\mu}_i + \sigma_v \left[\frac{\phi(-\tilde{\mu}_i/\sigma_v)}{1 - \Phi(-\tilde{\mu}_i/\sigma_v)} \right] = \sigma_v \left[\frac{\phi(A)}{\Phi(-A)} - A \right]$$

Normal-Truncated Normal Model

Define $\tilde{\mu} = (-\sigma_u^2\epsilon_i + \mu\sigma_v^2)/\sigma^2$ and $\sigma_*^2 = \sigma_u^2\sigma_v^2/\sigma^2$. Then, as it is shown by Kumbhakar and Lovell (2000), conditional density is as follows:

$$f(u|\epsilon) = \frac{1}{\sqrt{2\pi}\sigma_*[1 - \Phi(-\tilde{\mu}/\sigma_*)]} \exp\left\{-\frac{(u - \tilde{\mu})^2}{2\sigma_*^2}\right\}$$

Hence, $f(u|\epsilon)$ is the density for $N^+(\tilde{\mu}, \sigma_*^2)$.

Using this result, it follows that the estimate of technical efficiency is

$$TE1_i = E(\exp\{-u_i\}|\epsilon_i) = \frac{1 - \Phi(\sigma_* - \tilde{\mu}_i/\sigma_*)}{1 - \Phi(-\tilde{\mu}_i/\sigma_*)} \exp\left\{-\tilde{\mu}_i + \frac{1}{2}\sigma_*^2\right\}$$

The second version of the estimate is

$$TE2_i = \exp\{-E(u_i|\epsilon_i)\}$$

where

$$E(u_i|\epsilon_i) = \tilde{\mu}_i + \sigma_* \left[\frac{\phi(\tilde{\mu}_i/\sigma_*)}{1 - \Phi(-\tilde{\mu}_i/\sigma_*)} \right]$$

OUTEST= Data Set

The OUTEST= data set contains all the parameters estimated in a MODEL statement. The OUTEST= option can be used when the PROC QLIM call contains one MODEL statement:

```
proc qlim data=a outest=e;
  model y = x1 x2 x3;
  endogenous y ~ censored(lb=0);
run;
```

Each parameter contains the estimate for the corresponding parameter in the corresponding model. In addition, the OUTEST= data set contains the following variables:

<code>_NAME_</code>	the name of the independent variable
<code>_TYPE_</code>	type of observation. PARM indicates the row of coefficients; STD indicates the row of standard deviations of the corresponding coefficients.
<code>_STATUS_</code>	convergence status for optimization

The rest of the columns correspond to the explanatory variables.

The OUTEST= data set contains one observation for the MODEL statement, giving the parameter estimates for that model. If the COVOUT option is specified, the OUTEST= data set includes additional observations for the MODEL statement, giving the rows of the covariance matrix of parameter estimates. For covariance observations, the value of the `_TYPE_` variable is COV, and the `_NAME_` variable identifies the parameter associated with that row of the covariance matrix. If the CORROUT option is specified, the OUTEST= data set includes additional observations for the MODEL statement, giving the rows of the correlation matrix of parameter estimates. For correlation observations, the value of the `_TYPE_` variable is CORR, and the `_NAME_` variable identifies the parameter associated with that row of the correlation matrix.

Naming

Naming of Parameters

When there is only one equation in the estimation, parameters are named in the same way as in other SAS procedures such as REG, PROBIT, and so on. The constant in the regression equation is called Intercept. The coefficients on independent variables are named by the independent variables. The standard deviation of the errors is called `_Sigma`. If there are Box-Cox transformations, the coefficients are named `_Lambdai`, where i increments from 1, or as specified by the user. The limits for the discrete dependent variable are named `_Limiti`. If the LIMIT=varying option is specified, then `_Limiti` starts from 1. If the LIMIT=varying option is not specified, then `_Limit1` is set to 0 and the limit parameters start from $i = 2$. If the HETERO statement is included, the coefficients of the independent variables in the hetero equation are called `_H.x`, where x is the name of the independent variable. If the parameter name includes interaction terms, it needs to be enclosed in quotation marks followed by N . The following example restricts the parameter that includes the interaction term to be greater than zero:

```

proc qlim data=a;
  model y = x1|x2;
  endogenous y ~ discrete;
  restrict "x1*x2">0;
run;

```

When there are multiple equations in the estimation, the parameters in the main equation are named in the format of $y.x$, where y is the name of the dependent variable and x is the name of the independent variable. The standard deviation of the errors is called $_Sigma.y$. The correlation of the errors is called $_Rho$ for bivariate model. For the model with three variables it is $_Rho.y1.y2$, $_Rho.y1.y3$, $_Rho.y2.y3$. The construction of correlation names for multivariate models is analogous. Box-Cox parameters are called $_Lambda.i.y$ and limit variables are called $_Limit.i.y$. Parameters in the HETERO statement are named as $_H.y.x$. In the OUTEST= data set, all variables are changed from $_.$ to $_'$.

Naming of Output Variables

The following table shows the option in the OUTPUT statement, with the corresponding variable names and their explanation.

Option	Name	Explanation
PREDICTED	P_y	Predicted value of y
RESIDUAL	RESID_y	Residual of y , $(y - \text{Predicted}Y)$
XBETA	XBETA_y	Structure part $(x'\beta)$ of y equation
ERRSTD	ERRSTD_y	Standard deviation of error term
PROB	PROB_y	Probability that y is taking the observed value in this observation (discrete y only)
PROBALL	PROB_i_y	Probability that y is taking the i th value (discrete y only)
MILLS	MILLS_y	Inverse Mills ratio for y
EXPECTED	EXPCT_y	Unconditional expected value of y
CONDITIONAL	CEXPCT_y	Conditional expected value of y , condition on the truncation.
MARGINAL	MEFF_x	Marginal effect of x on y ($\frac{\partial y}{\partial x}$) with single equation
	MEFF_y_x	Marginal effect of x on y ($\frac{\partial y}{\partial x}$) with multiple equations
	MEFF_Pi_x	Marginal effect of x on y ($\frac{\partial \text{Prob}(y=i)}{\partial x}$) with single equation and discrete y
	MEFF_Pi_y_x	Marginal effect of x on y ($\frac{\partial \text{Prob}(y=i)}{\partial x}$) with multiple equations and discrete y
TE1	TE1	Technical efficiency estimate for each producer proposed by Battese and Coelli (1988)
TE2	TE2	Technical efficiency estimate for each producer proposed by Jondrow et al. (1982)

If you prefer to name the output variables differently, you can use the RENAME option in the data set. For example, the following statements rename the residual of *y* as *Resid*:

```
proc qlim data=one;
  model y = x1-x10 / censored;
  output out=outds(rename=(resid_y=resid)) residual;
run;
```

ODS Table Names

PROC QLIM assigns a name to each table it creates. You can use these names to denote the table when using the Output Delivery System (ODS) to select tables and create output data sets. These names are listed in the Table 22.9.

Table 22.9 ODS Tables Produced in PROC QLIM by the MODEL Statement and TEST Statement

ODS Table Name	Description	Option
ODS Tables Created by the MODEL Statement and TEST Statement		
ResponseProfile	Response profile	default
ClassLevels	Class levels	default
FitSummary	Summary of nonlinear estimation	default
GoodnessOfFit	Pseudo-R-square measures	default
ConvergenceStatus	Convergence status	default
ParameterEstimates	Parameter estimates	default
SummaryContResponse	Summary of continuous response	default
CovB	Covariance of parameter estimates	COVB
CorrB	Correlation of parameter estimates	CORRB
FitSummaryHeckman1	Heckman First Step Model Fit Summary	HECKIT
FitSummaryHeckman2	Heckman Second Model Fit Summary	HECKIT
LinCon	Linear constraints	ITPRINT
InputOptions	Input options	ITPRINT
ProblemDescription	Problem description	ITPRINT
IterStart	Optimization start summary	ITPRINT
IterHist	Iteration history	ITPRINT
IterStop	Optimization results	ITPRINT
ConvergenceStatus	Convergence status	ITPRINT
ParameterEstimatesStart	Optimization start	ITPRINT
ParameterEstimatesResults	Resulting parameters	ITPRINT
LinConSol	Linear constraints evaluated at solution	ITPRINT
VariableSelection	Variable selection summary	SELECTVAR
ODS Tables Created by the TEST Statement		
TestResults	Test results	default

Table 22.9 (continued)

ODS Table Name	Description	Option
ODS Tables Created by the BAYES Statement		
AutoCorr	Autocorrelation statistics for each parameter	default
Corr	Correlation matrix of the posterior samples	STATS=COR
Cov	Covariance matrix of the posterior samples	STATS=COV
ESS	Effective sample size for each parameter	Default
MCSE	Monte Carlo standard error for each parameter	Default
Geweke	Geweke diagnostics for each parameter	Default
Heidelberger	Heidelberger-Welch diagnostics for each parameter	DIAGNOSTICS=HEIDEL
PostIntervals	Equal-tail and HPD intervals for each parameter	Default
PosteriorSample	Posterior samples	(ODS output data set only)
PostSummaries	Posterior summaries	default
PriorSample	Prior samples used for prior predictive analysis	(ODS output data set only)
PriorSummaries	Prior summaries	STATS=PRIOR
Raftery	Raftery-Lewis diagnostics for each parameter	DIAGNOSTICS=RAFTER

ODS Graphics

You can reference every graph that is produced through ODS Graphics with a name. The names of the graphs that PROC QLIM generates are listed in [Table 22.10](#) for the frequentist approach and in [Table 22.11](#) for the Bayesian approach.

Table 22.10 Graphs Produced by PROC QLIM without a BAYES Statement

ODS Graph Name	Plot Description	Statement & Option
Frequentist Output Plots		
ResidPlot	Frequentist analysis of residuals	PLOTS=RESIDUAL
XbetaPlot	Frequentist analysis of xbeta	PLOTS=XBETA
PredPlot	Frequentist analysis of Predictions	PLOTS=PREDICTED
MargnPlot	Frequentist analysis of marginal effects	PLOTS=MARGINAL
ErrStdPlot	Frequentist analysis of the error standard deviation (meaningful only with a HETERO statement)	PLOTS=ERRSTD
MillsPlot	Frequentist analysis of Mills ratio	PLOTS=MILLS
ExpctPlot	Frequentist analysis of expected values for continuous endogenous variables	PLOTS=EXPECTED
TE1Plot	Frequentist analysis of technical efficiency (only in stochastic frontier model) suggested by Battese and Coelli (1988)	PLOTS=TE1

Table 22.10 *continued*

ODS Graph Name	Plot Description	Statement and Option
TE2Plot	Frequentist analysis of technical efficiency (only in stochastic frontier model) suggested by Jondrow et al. (1982)	PLOTS=TE2
CExpctPlot	Frequentist analysis of conditional expected values for continuous endogenous variables	PLOTS=CONDITIONAL
ProbPlot	Frequentist analysis of probability of discrete endogenous variables that take the current observed responses	PLOTS=PROB
ProbAllPlot	Frequentist analysis of probability of discrete endogenous variables for all responses	PLOTS=PROBALL
ProfLikPlot	Profile log-likelihood plot	PLOTS=PROFLIK

Table 22.11 Graphs Produced by PROC QLIM When a BAYES Statement Is Included

ODS Graph Name	Plot Description	Statement and Option
Bayesian Diagnostic Plots		
ADPanel	Autocorrelation function and density panel	PLOTS=(AUTOCORR DENSITY)
AutocorrPanel	Autocorrelation function panel	PLOTS=AUTOCORR
AutocorrPlot	Autocorrelation function plot	PLOTS(UNPACK)=AUTOCORR
DensityPanel	Density panel	PLOTS=DENSITY
DensityPlot	Density plot	PLOTS(UNPACK)=DENSITY
ProfLikPlot	Profile log-likelihood plot	PLOTS=PROFLIK
TAPanel	Trace and autocorrelation function panel	PLOTS=(TRACE AUTOCORR)
TADPanel	Trace, density, and autocorrelation function panel	PLOTS=(TRACE AUTOCORR DENSITY)
		PLOTS=BAYESDIAG
TDPanel	Trace and density panel	PLOTS=(TRACE DENSITY)
TracePanel	Trace panel	PLOTS=TRACE
TracePlot	Trace plot	PLOTS(UNPACK)=TRACE
Bayesian Summary Plots		
BayesSumPlot	Prior/posterior densities and MLE	PLOTS=BAYESSUM
Bayesian Output Plots		
PredictiveByObsNumPlot	Predictive analysis by observation number	PLOTS(PRIOR)=BAYESPRED
PredictivePlot	Predictive analysis by regressor	PLOTS(PRIOR)=BAYESPRED

Examples: QLIM Procedure

Example 22.1: Ordered Data Modeling

Cameron and Trivedi (1986, 1998) studied the number of doctor visits from the Australian Health Survey 1977-78. In the following data set, the dependent variable, DVISITS, contains the number of doctor visits in the past 2 weeks (0, 1, or more than 2). The explanatory variables are: SEX indicates if the patient is female; AGE is the age in years divided by 100; INCOME is the annual income (\$10,000); LEVYPLUS indicates if the patient has private health insurance; FREEPOOR indicates free government health insurance due to low income; FREEREPA indicates free government health insurance for other reasons; ILLNESS is the number of illnesses in the past 2 weeks; ACTDAYS is the number of days the illness caused reduced activity; HSCORE is a questionnaire score; CHCOND1 indicates a chronic condition that does not limit activity; and CHCOND2 indicates a chronic condition that limits activity.

```
data docvisit;
  input sex age agesq income levyplus freepoor freerepa
        illness actdays hscore chcond1 chcond2 dvisits;
  y = (dvisits > 0);
  if ( dvisits > 8 ) then dvisits = 8;
datalines;
1 0.19 0.0361 0.55 1 0 0 1 4 1 0 0 1
1 0.19 0.0361 0.45 1 0 0 1 2 1 0 0 1

... more lines ...

1 0.37 0.1369 0.25 0 0 1 1 0 1 0 0 0
1 0.52 0.2704 0.65 0 0 0 0 0 0 0 0 0
0 0.72 0.5184 0.25 0 0 1 0 0 0 0 0 0
;
```

The dependent variable, dvisits, has nine ordered values. The following SAS statements estimate the ordinal probit model:

```
/*-- Ordered Discrete Responses --*/
proc qlim data=docvisit;
  model dvisits = sex age agesq income levyplus
                freepoor freerepa illness actdays hscore
                chcond1 chcond2 / discrete;
run;
```

The output of the QLIM procedure for ordered data modeling is shown in [Output 22.1.1](#).

Output 22.1.1 Ordered Data Modeling**Binary Data****The QLIM Procedure**

Discrete Response Profile of dvisits		
Index	Value	Total Frequency
1	0	4141
2	1	782
3	2	174
4	3	30
5	4	24
6	5	9
7	6	12
8	7	12
9	8	6

Output 22.1.1 *continued*

Model Fit Summary	
Number of Endogenous Variables	1
Endogenous Variable	dvisits
Number of Observations	5190
Log Likelihood	-3138
Maximum Absolute Gradient	0.0003675
Number of Iterations	82
Optimization Method	Quasi-Newton
AIC	6316
Schwarz Criterion	6447

Goodness-of-Fit Measures		
Measure	Value	Formula
Likelihood Ratio (R)	789.73	$2 * (\text{LogL} - \text{LogL0})$
Upper Bound of R (U)	7065.9	$-2 * \text{LogL0}$
Aldrich-Nelson	0.1321	$R / (R+N)$
Cragg-Uhler 1	0.1412	$1 - \exp(-R/N)$
Cragg-Uhler 2	0.1898	$(1 - \exp(-R/N)) / (1 - \exp(-U/N))$
Estrella	0.149	$1 - (1 - R/U)^{(U/N)}$
Adjusted Estrella	0.1416	$1 - ((\text{LogL} - K) / \text{LogL0})^{(-2/N * \text{LogL0})}$
McFadden's LRI	0.1118	R / U
Veall-Zimmermann	0.2291	$(R * (U+N)) / (U * (R+N))$
McKelvey-Zavoina	0.2036	
N = # of observations, K = # of regressors		

Output 22.1.1 continued

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
Intercept	1	-1.378705	0.147413	-9.35	<.0001
sex	1	0.131885	0.043785	3.01	0.0026
age	1	-0.534190	0.815907	-0.65	0.5126
agesq	1	0.857308	0.898364	0.95	0.3399
income	1	-0.062211	0.068017	-0.91	0.3604
levyplus	1	0.137030	0.053262	2.57	0.0101
freepoor	1	-0.346045	0.129638	-2.67	0.0076
freerepa	1	0.178382	0.074348	2.40	0.0164
illness	1	0.150485	0.015747	9.56	<.0001
actdays	1	0.100575	0.005850	17.19	<.0001
hscore	1	0.031862	0.009201	3.46	0.0005
chcond1	1	0.061601	0.049024	1.26	0.2089
chcond2	1	0.135321	0.067711	2.00	0.0457
_Limit2	1	0.938884	0.031219	30.07	<.0001
_Limit3	1	1.514288	0.049329	30.70	<.0001
_Limit4	1	1.711660	0.058151	29.43	<.0001
_Limit5	1	1.952860	0.072014	27.12	<.0001
_Limit6	1	2.087422	0.081655	25.56	<.0001
_Limit7	1	2.333786	0.101760	22.93	<.0001
_Limit8	1	2.789796	0.156189	17.86	<.0001

By default, ordinal probit/logit models are estimated assuming that the first threshold or limit parameter (μ_1) is 0. However, this parameter can also be estimated when the LIMIT1=VARYING option is specified. The probability that y_i^* belongs to the j th category is defined as

$$P[\mu_{j-1} < y_i^* < \mu_j] = F(\mu_j - \mathbf{x}_i' \boldsymbol{\beta}) - F(\mu_{j-1} - \mathbf{x}_i' \boldsymbol{\beta})$$

where $F(\cdot)$ is the logistic or standard normal CDF, $\mu_0 = -\infty$ and $\mu_9 = \infty$. Output 22.1.2 lists ordinal probit estimates computed in the following program. Note that the intercept term is suppressed for model identification when μ_1 is estimated.

```

/*-- Ordered Probit --*/
proc qlim data=docvisit;
  model dvisits = sex age agesq income levyplus
               freepoor freerepa illness actdays hscore
               chcond1 chcond2 / discrete(d=normal) limit1=varying;
run;

```


Output 22.1.2 Ordinal Probit Parameter Estimates with LIMIT1=VARYING**Binary Data****The QLIM Procedure**

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
sex	1	0.131885	0.043785	3.01	0.0026
age	1	-0.534181	0.815915	-0.65	0.5127
agesq	1	0.857298	0.898371	0.95	0.3399
income	1	-0.062211	0.068017	-0.91	0.3604
levyplus	1	0.137031	0.053262	2.57	0.0101
freepoor	1	-0.346045	0.129638	-2.67	0.0076
freerepa	1	0.178382	0.074348	2.40	0.0164
illness	1	0.150485	0.015747	9.56	<.0001
actdays	1	0.100575	0.005850	17.19	<.0001
hscore	1	0.031862	0.009201	3.46	0.0005
chcond1	1	0.061602	0.049024	1.26	0.2089
chcond2	1	0.135322	0.067711	2.00	0.0457
_Limit1	1	1.378706	0.147415	9.35	<.0001
_Limit2	1	2.317590	0.150206	15.43	<.0001
_Limit3	1	2.892994	0.155198	18.64	<.0001
_Limit4	1	3.090367	0.158263	19.53	<.0001
_Limit5	1	3.331566	0.164065	20.31	<.0001
_Limit6	1	3.466128	0.168799	20.53	<.0001
_Limit7	1	3.712493	0.179756	20.65	<.0001
_Limit8	1	4.168502	0.215738	19.32	<.0001

Example 22.2: Tobit Analysis

The following statements show a subset of the Mroz (1987) data set. In these data, Hours is the number of hours the wife worked outside the household in a given year, Yrs_Ed is the years of education, and Yrs_Exp is the years of work experience. A Tobit model will be fit to the hours worked with years of education and experience as covariates.

By the nature of the data it is clear that there are a number of women who committed some positive number of hours to outside work ($y_i > 0$ is observed). There are also a number of women who did not work at all ($y_i = 0$ is observed). This gives us the following model:

$$y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \epsilon_i$$

$$y_i = \begin{cases} y_i^* & \text{if } y_i^* > 0 \\ 0 & \text{if } y_i^* \leq 0 \end{cases}$$

where $\epsilon_i \sim iidN(0, \sigma^2)$. The set of explanatory variables is denoted by \mathbf{x}_i .

```

title1 'Estimating a Tobit model';

data subset;
    input Hours Yrs_Ed Yrs_Exp @@;
    if Hours eq 0 then Lower=.;
    else
        Lower=Hours;
datalines;
0 8 9 0 8 12 0 9 10 0 10 15 0 11 4 0 11 6
1000 12 1 1960 12 29 0 13 3 2100 13 36
3686 14 11 1920 14 38 0 15 14 1728 16 3
1568 16 19 1316 17 7 0 17 15
;

/*-- Tobit Model --*/
proc qlim data=subset;
    model hours = yrs_ed yrs_exp;
    endogenous hours ~ censored(lb=0);
run;

```

The output of the QLIM procedure is shown in [Output 22.2.1](#).

Output 22.2.1 Tobit Analysis Results

Estimating a Tobit model

The QLIM Procedure

Model Fit Summary					
Number of Endogenous Variables		1			
Endogenous Variable		Hours			
Number of Observations		17			
Log Likelihood		-74.93700			
Maximum Absolute Gradient		1.18953E-6			
Number of Iterations		23			
Optimization Method		Quasi-Newton			
AIC		157.87400			
Schwarz Criterion		161.20685			

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
Intercept	1	-5598.295129	27.692220	-202.16	<.0001
Yrs_Ed	1	373.123254	53.988877	6.91	<.0001
Yrs_Exp	1	63.336247	36.551299	1.73	0.0831
_Sigma	1	1582.859635	390.076480	4.06	<.0001

In the “Parameter Estimates” table there are four rows. The first three of these rows correspond to the vector estimate of the regression coefficients β . The last one is called `_Sigma`, which corresponds to the estimate of the error variance σ .

Example 22.3: Bivariate Probit Analysis

This example shows how to estimate a bivariate probit model. Note the `INIT` statement in the following program, which sets the initial values for some parameters in the optimization:

```
data a;
  keep y1 y2 x1 x2;
  do i = 1 to 500;
    x1 = rannor( 19283 );
    x2 = rannor( 19283 );
    u1 = rannor( 19283 );
    u2 = rannor( 19283 );
    y1l = 1 + 2 * x1 + 3 * x2 + u1;
    y2l = 3 + 4 * x1 - 2 * x2 + u1*.2 + u2;
    if ( y1l > 0 ) then y1 = 1;
    else y1 = 0;
    if ( y2l > 0 ) then y2 = 1;
    else y2 = 0;
    output;
  end;
run;

/*-- Bivariate Probit --*/
proc qlim data=a method=qn;
  init y1.x1 2.8, y1.x2 2.1, _rho .1;
  model y1 = x1 x2;
  model y2 = x1 x2;
  endogenous y1 y2 ~ discrete;
run;
```

The output of the QLIM procedure is shown in [Output 22.3.1](#).

Output 22.3.1 Bivariate Probit Analysis Results**Estimating a Tobit model****The QLIM Procedure**

Model Fit Summary					
Number of Endogenous Variables		2			
Endogenous Variable		y1 y2			
Number of Observations		500			
Log Likelihood		-134.90796			
Maximum Absolute Gradient		3.23363E-7			
Number of Iterations		17			
Optimization Method		Quasi-Newton			
AIC		283.81592			
Schwarz Criterion		313.31817			

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
y1.Intercept	1	1.003639	0.153678	6.53	<.0001
y1.x1	1	2.244374	0.256062	8.76	<.0001
y1.x2	1	3.273441	0.341581	9.58	<.0001
y2.Intercept	1	3.621164	0.457173	7.92	<.0001
y2.x1	1	4.551525	0.576547	7.89	<.0001
y2.x2	1	-2.442769	0.332295	-7.35	<.0001
_Rho	1	0.144097	0.336459	0.43	0.6685

Example 22.4: Sample Selection Model

This example illustrates the use of PROC QLIM for sample selection models. The data set is the same one from Mroz (1987). The goal is to estimate a wage offer function for married women, accounting for potential selection bias. Of the 753 women, the wage is observed for 428 working women. The labor force participation equation estimated in the introductory example is used for selection. The wage equation uses log wage (`lwage`) as the dependent variable. The explanatory variables in the wage equation are the woman's years of schooling (`educ`), wife's labor experience (`exper`), and square of experience (`expersq`). The program is as follows:

```

/*-- Sample Selection --*/
proc qlim data=mroz;
    model inlf = nwifeinc educ exper expersq
               age kidslt6 kidsge6 /discrete;
    model lwage = educ exper expersq / select(inlf=1);
run;

```

The output of the QLIM procedure is shown in [Output 22.4.1](#).

Output 22.4.1 Sample Selection**Binary Data****The QLIM Procedure**

Model Fit Summary					
Number of Endogenous Variables					2
Endogenous Variable					lnlf lwage
Number of Observations					753
Log Likelihood					-832.88509
Maximum Absolute Gradient					0.00502
Number of Iterations					78
Optimization Method					Quasi-Newton
AIC					1694
Schwarz Criterion					1759

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
lwage.Intercept	1	-0.552716	0.260371	-2.12	0.0338
lwage.educ	1	0.108351	0.014861	7.29	<.0001
lwage.exper	1	0.042837	0.014878	2.88	0.0040
lwage.expersq	1	-0.000837	0.000417	-2.01	0.0449
_Sigma.lwage	1	0.663397	0.022706	29.22	<.0001
lnlf.Intercept	1	0.266459	0.508954	0.52	0.6006
lnlf.nwifeinc	1	-0.012132	0.004877	-2.49	0.0129
lnlf.educ	1	0.131341	0.025383	5.17	<.0001
lnlf.exper	1	0.123282	0.018728	6.58	<.0001
lnlf.expersq	1	-0.001886	0.000601	-3.14	0.0017
lnlf.age	1	-0.052829	0.008479	-6.23	<.0001
lnlf.kidslt6	1	-0.867398	0.118647	-7.31	<.0001
lnlf.kidsge6	1	0.035872	0.043476	0.83	0.4093
_Rho	1	0.026617	0.147073	0.18	0.8564

Note the correlation estimate is insignificant. This indicates that selection bias is not a big problem in the estimation of wage equation.

Example 22.5: Sample Selection Model with Truncation and Censoring

In this example the data are generated such that the selection variable is discrete and the variable Y is truncated from below by zero. The program follows, with the results shown in [Output 22.5.1](#):

```

data trunc;
  keep z y x1 x2;
  do i = 1 to 500;
    x1 = rannor( 19283 );
    x2 = rannor( 19283 );
    u1 = rannor( 19283 );
    u2 = rannor( 19283 );
    z1 = 1 + 2 * x1 + 3 * x2 + u1;
    y = 3 + 4 * x1 - 2 * x2 + u1*.2 + u2;
    if ( z1 > 0 ) then z = 1;
    else z = 0;
    if y>=0 then output;
  end;
run;

/*-- Sample Selection with Truncation --*/
proc qlim data=trunc method=qn;
  model z = x1 x2 / discrete;
  model y = x1 x2 / select(z=1) truncated(lb=0);
run;

```

Output 22.5.1 Sample Selection with Truncation**Binary Data****The QLIM Procedure**

Model Fit Summary						
Number of Endogenous Variables			2			
Endogenous Variable			z y			
Number of Observations			379			
Log Likelihood			-344.10722			
Maximum Absolute Gradient			4.95535E-6			
Number of Iterations			17			
Optimization Method			Quasi-Newton			
AIC			704.21444			
Schwarz Criterion			735.71473			

Parameter Estimates						
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t	
y.Intercept	1	3.014158	0.128548	23.45	<.0001	
y.x1	1	3.995671	0.099599	40.12	<.0001	
y.x2	1	-1.972697	0.096385	-20.47	<.0001	
_Sigma.y	1	0.923428	0.047233	19.55	<.0001	
z.Intercept	1	0.949444	0.190265	4.99	<.0001	
z.x1	1	2.163928	0.288384	7.50	<.0001	
z.x2	1	3.134213	0.379251	8.26	<.0001	
_Rho	1	0.494356	0.176542	2.80	0.0051	

In the following statements the data are generated such that the selection variable is discrete and the variable *Y* is censored from below by zero. The results are shown in [Output 22.5.2](#).

```

data cens;
  keep z y x1 x2;
  do i = 1 to 500;
    x1 = rannor( 19283 );
    x2 = rannor( 19283 );
    u1 = rannor( 19283 );
    u2 = rannor( 19283 );
    z1 = 1 + 2 * x1 + 3 * x2 + u1;
    y1 = 3 + 4 * x1 - 2 * x2 + u1*.2 + u2;
    if ( z1 > 0 ) then z = 1;
    else z = 0;
    if ( y1 > 0 ) then y = y1;
    else y = 0;
    output;
  end;
run;

/*-- Sample Selection with Censoring --*/
proc qlim data=cens method=qn;
  model z = x1 x2 / discrete;
  model y = x1 x2 / select(z=1) censored(lb=0);
run;

```

Output 22.5.2 Sample Selection with Censoring

Binary Data

The QLIM Procedure

Model Fit Summary						
Number of Endogenous Variables			2			
Endogenous Variable			z y			
Number of Observations			500			
Log Likelihood			-399.78508			
Maximum Absolute Gradient			2.30443E-6			
Number of Iterations			19			
Optimization Method			Quasi-Newton			
AIC			815.57015			
Schwarz Criterion			849.28702			

Parameter Estimates						
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t	
y.Intercept	1	3.074276	0.111617	27.54	<.0001	
y.x1	1	3.963619	0.085796	46.20	<.0001	
y.x2	1	-2.023548	0.088714	-22.81	<.0001	
_Sigma.y	1	0.920860	0.043278	21.28	<.0001	
z.Intercept	1	1.013610	0.154081	6.58	<.0001	
z.x1	1	2.256922	0.255999	8.82	<.0001	
z.x2	1	3.302692	0.342168	9.65	<.0001	
_Rho	1	0.350776	0.197093	1.78	0.0751	

Example 22.6: Types of Tobit Models

The following five examples show how to estimate different types of Tobit models (see “Types of Tobit Models” on page 1558). [Output 22.6.1](#) through [Output 22.6.5](#) show the results of the corresponding programs.

Type 1 Tobit

```
data a1;
  keep y x;
  do i = 1 to 500;
    x = rannor( 19283 );
    u = rannor( 19283 );
    y1 = 1 + 2 * x + u;
    if ( y1 > 0 ) then y = y1;
    else          y = 0;
    output;
  end;
run;

/*-- Type 1 Tobit --*/
proc qlim data=a1 method=qn;
  model y = x;
  endogenous y ~ censored(lb=0);
run;
```

Output 22.6.1 Type 1 Tobit

Binary Data

The QLIM Procedure

Model Fit Summary					
Number of Endogenous Variables		1			
Endogenous Variable		y			
Number of Observations		500			
Log Likelihood		-554.17696			
Maximum Absolute Gradient		4.65556E-7			
Number of Iterations		9			
Optimization Method		Quasi-Newton			
AIC		1114			
Schwarz Criterion		1127			

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
Intercept	1	0.942734	0.056784	16.60	<.0001
x	1	2.049571	0.066979	30.60	<.0001
_Sigma	1	1.016571	0.039035	26.04	<.0001

Type 2 Tobit

```

data a2;
  keep y1 y2 x1 x2;
  do i = 1 to 500;
    x1 = rannor( 19283 );
    x2 = rannor( 19283 );
    u1 = rannor( 19283 );
    u2 = rannor( 19283 );
    y1l = 1 + 2 * x1 + 3 * x2 + u1;
    y2l = 3 + 4 * x1 - 2 * x2 + u1*.2 + u2;
    if ( y1l > 0 ) then y1 = 1;
    else y1 = 0;
    if ( y1l > 0 ) then y2 = y2l;
    else y2 = 0;
    output;
  end;
run;

/*-- Type 2 Tobit --*/
proc qlim data=a2 method=qn;
  model y1 = x1 x2 / discrete;
  model y2 = x1 x2 / select(y1=1);
run;

```

Output 22.6.2 Type 2 Tobit**Binary Data****The QLIM Procedure**

Model Fit Summary					
Number of Endogenous Variables		2			
Endogenous Variable		y1 y2			
Number of Observations		500			
Log Likelihood		-476.12328			
Maximum Absolute Gradient		8.30075E-7			
Number of Iterations		17			
Optimization Method		Quasi-Newton			
AIC		968.24655			
Schwarz Criterion		1002			

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
y2.Intercept	1	3.066992	0.106903	28.69	<.0001
y2.x1	1	4.004874	0.072043	55.59	<.0001
y2.x2	1	-2.079352	0.087544	-23.75	<.0001
_Sigma.y2	1	0.940559	0.039321	23.92	<.0001
y1.Intercept	1	1.017140	0.154975	6.56	<.0001
y1.x1	1	2.253080	0.256097	8.80	<.0001
y1.x2	1	3.305140	0.343695	9.62	<.0001
_Rho	1	0.292992	0.210073	1.39	0.1631

Type 3 Tobit

```

data a3;
  keep y1 y2 x1 x2;
  do i = 1 to 500;
    x1 = rannor( 19283 );
    x2 = rannor( 19283 );
    u1 = rannor( 19283 );
    u2 = rannor( 19283 );
    y1l = 1 + 2 * x1 + 3 * x2 + u1;
    y2l = 3 + 4 * x1 - 2 * x2 + u1*.2 + u2;
    if ( y1l > 0 ) then y1 = y1l;
    else y1 = 0;
    if ( y1l > 0 ) then y2 = y2l;
    else y2 = 0;
    output;
  end;
run;

/*-- Type 3 Tobit --*/
proc qlim data=a3 method=qn;
  model y1 = x1 x2 / censored(lb=0);
  model y2 = x1 x2 / select(y1>0);
run;

```

Output 22.6.3 Type 3 Tobit**Binary Data****The QLIM Procedure**

Model Fit Summary	
Number of Endogenous Variables	2
Endogenous Variable	y1 y2
Number of Observations	500
Log Likelihood	-838.94087
Maximum Absolute Gradient	9.71691E-6
Number of Iterations	16
Optimization Method	Quasi-Newton
AIC	1696
Schwarz Criterion	1734

Output 22.6.3 continued

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
y2.Intercept	1	3.081206	0.080121	38.46	<.0001
y2.x1	1	3.998361	0.063734	62.73	<.0001
y2.x2	1	-2.088280	0.072876	-28.66	<.0001
_Sigma.y2	1	0.939799	0.039047	24.07	<.0001
y1.Intercept	1	0.981975	0.067351	14.58	<.0001
y1.x1	1	2.032675	0.059363	34.24	<.0001
y1.x2	1	2.976609	0.065584	45.39	<.0001
_Sigma.y1	1	0.969968	0.039795	24.37	<.0001
_Rho	1	0.226281	0.057672	3.92	<.0001

Type 4 Tobit

```

data a4;
  keep y1 y2 y3 x1 x2;
  do i = 1 to 500;
    x1 = rannor( 19283 );
    x2 = rannor( 19283 );
    u1 = rannor( 19283 );
    u2 = rannor( 19283 );
    u3 = rannor( 19283 );
    y1l = 1 + 2 * x1 + 3 * x2 + u1;
    y2l = 3 + 4 * x1 - 2 * x2 + u1*.2 + u2;
    y3l = 0 - 1 * x1 + 1 * x2 + u1*.1 - u2*.5 + u3*.5;
    if ( y1l > 0 ) then y1 = y1l;
    else y1 = 0;
    if ( y1l > 0 ) then y2 = y2l;
    else y2 = 0;
    if ( y1l <= 0 ) then y3 = y3l;
    else y3 = 0;
    output;
  end;
run;

/*-- Type 4 Tobit --*/
proc qlim data=a4 method=qn;
  model y1 = x1 x2 / censored(lb=0);
  model y2 = x1 x2 / select(y1>0);
  model y3 = x1 x2 / select(y1<=0);
run;

```

Output 22.6.4 Type 4 Tobit**Binary Data****The QLIM Procedure**

Model Fit Summary					
Number of Endogenous Variables		3			
Endogenous Variable		y1 y2 y3			
Number of Observations		500			
Log Likelihood		-1128			
Maximum Absolute Gradient		0.0000161			
Number of Iterations		21			
Optimization Method		Quasi-Newton			
AIC		2285			
Schwarz Criterion		2344			

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
y2.Intercept	1	2.894656	0.076079	38.05	<.0001
y2.x1	1	4.072704	0.062675	64.98	<.0001
y2.x2	1	-1.901163	0.076874	-24.73	<.0001
_Sigma.y2	1	0.981655	0.039564	24.81	<.0001
y3.Intercept	1	0.064594	0.179441	0.36	0.7189
y3.x1	1	-0.938384	0.096570	-9.72	<.0001
y3.x2	1	1.035798	0.123104	8.41	<.0001
_Sigma.y3	1	0.743124	0.038240	19.43	<.0001
y1.Intercept	1	0.987370	0.067861	14.55	<.0001
y1.x1	1	2.050408	0.060819	33.71	<.0001
y1.x2	1	2.982190	0.072552	41.10	<.0001
_Sigma.y1	1	1.032473	0.040971	25.20	<.0001
_Rho.y1.y2	1	0.291587	0.053436	5.46	<.0001
_Rho.y1.y3	1	-0.031665	0.260057	-0.12	0.9031

Type 5 Tobit

```

data a5;
  keep y1 y2 y3 x1 x2;
  do i = 1 to 500;
    x1 = rannor( 19283 );
    x2 = rannor( 19283 );
    u1 = rannor( 19283 );
    u2 = rannor( 19283 );
    u3 = rannor( 19283 );
    y1l = 1 + 2 * x1 + 3 * x2 + u1;
    y2l = 3 + 4 * x1 - 2 * x2 + u1*.2 + u2;
    y3l = 0 - 1 * x1 + 1 * x2 + u1*.1 - u2*.5 + u3*.5;
    if ( y1l > 0 ) then y1 = 1;
    else y1 = 0;
    if ( y1l > 0 ) then y2 = y2l;
    else y2 = 0;
    if ( y1l <= 0 ) then y3 = y3l;
    else y3 = 0;
    output;
  end;
run;

/*-- Type 5 Tobit --*/
proc qlim data=a5 method=qn;
  model y1 = x1 x2 / discrete;
  model y2 = x1 x2 / select (y1>0);
  model y3 = x1 x2 / select (y1<=0);
run;

```

Output 22.6.5 Type 5 Tobit**Binary Data****The QLIM Procedure**

Model Fit Summary	
Number of Endogenous Variables	3
Endogenous Variable	y1 y2 y3
Number of Observations	500
Log Likelihood	-734.50612
Maximum Absolute Gradient	3.57134E-7
Number of Iterations	20
Optimization Method	Quasi-Newton
AIC	1495
Schwarz Criterion	1550

Output 22.6.5 continued

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
y2.Intercept	1	2.887523	0.095193	30.33	<.0001
y2.x1	1	4.078926	0.069623	58.59	<.0001
y2.x2	1	-1.898898	0.086578	-21.93	<.0001
_Sigma.y2	1	0.983059	0.039987	24.58	<.0001
y3.Intercept	1	0.071764	0.171522	0.42	0.6757
y3.x1	1	-0.935299	0.092843	-10.07	<.0001
y3.x2	1	1.039954	0.120697	8.62	<.0001
_Sigma.y3	1	0.743083	0.038225	19.44	<.0001
y1.Intercept	1	1.067578	0.142789	7.48	<.0001
y1.x1	1	2.068376	0.226020	9.15	<.0001
y1.x2	1	3.157385	0.314743	10.03	<.0001
_Rho.y1.y2	1	0.312369	0.177010	1.76	0.0776
_Rho.y1.y3	1	-0.018225	0.234886	-0.08	0.9382

Example 22.7: Stochastic Frontier Models

This example illustrates the estimation of stochastic frontier production and cost models.

First, a production function model is estimated. The data for this example were collected by Christensen Associates; they represent a sample of 125 observations on inputs and output for 10 airlines between 1970 and 1984. The explanatory variables (inputs) are fuel (LF), materials (LM), equipment (LE), labor (LL), and property (LP), and (LQ) is an index that represents passengers, charter, mail, and freight transported.

The following statements create the dataset:

```

title1 'Stochastic Frontier Production Model';
data airlines;
    input TS FIRM NI LQ LF LM LE LL LP;
datalines;
1 1 15 -0.0484 0.2473 0.2335 0.2294 0.2246 0.2124
1 1 15 -0.0133 0.2603 0.2492 0.241 0.2216 0.1069
2 1 15 0.088 0.2666 0.3273 0.3365 0.2039 0.0865
3 1 15 0.1619 0.3019 0.4573 0.3532 0.2346 0.0242
... more lines ...

```

The following statements estimate a stochastic frontier exponential production model that uses Christensen Associates data:

```

/*-- Stochastic Frontier Production Model --*/
proc qlim data=airlines;
    model LQ=LF LM LE LL LP;
    endogenous LQ ~ frontier (type=exponential production);
run;

```

Figure 22.7.1 shows the results from this production model.

Output 22.7.1 Stochastic Frontier Production Model**Stochastic Frontier Production Model****The QLIM Procedure**

Model Fit Summary					
Number of Endogenous Variables		1			
Endogenous Variable		LQ			
Number of Observations		125			
Log Likelihood		83.27815			
Maximum Absolute Gradient		9.83602E-6			
Number of Iterations		19			
Optimization Method		Quasi-Newton			
AIC		-150.55630			
Schwarz Criterion		-127.92979			
Sigma		0.12445			
Lambda		0.55766			

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
Intercept	1	-0.085048	0.024528	-3.47	0.0005
LF	1	-0.115802	0.124178	-0.93	0.3511
LM	1	0.756253	0.078755	9.60	<.0001
LE	1	0.424916	0.081893	5.19	<.0001
LL	1	-0.136421	0.089702	-1.52	0.1283
LP	1	0.098967	0.042776	2.31	0.0207
_Sigma_v	1	0.108688	0.010063	10.80	<.0001
_Sigma_u	1	0.060611	0.017603	3.44	0.0006

Similarly, the stochastic frontier production function can be estimated with (type=half) or (type=truncated) options that represent half-normal and truncated normal production models.

In the next step, stochastic frontier cost function is estimated. The data for the cost model are provided by Christensen and Greene (1976). The data describe costs and production inputs of 145 U.S. electricity producers in 1955. The model being estimated follows the nonhomogeneous version of the Cobb-Douglas cost function:

$$\log\left(\frac{\text{Cost}}{\text{FPrice}}\right) = \beta_0 + \beta_1 \log\left(\frac{\text{KPrice}}{\text{FPrice}}\right) + \beta_2 \log\left(\frac{\text{LPrice}}{\text{FPrice}}\right) + \beta_3 \log(\text{Output}) + \beta_4 \frac{1}{2} \log(\text{Output})^2 + \epsilon$$

All dollar values are normalized by fuel price. The quadratic log of the output is added to capture nonlinearities due to scale effects in cost functions. New variables, `log_C_PF`, `log_PK_PF`, `log_PL_PF`, `log_y`, and `log_y_sq`, are created to reflect transformations. The following statements create the data set and transformed variables:

```

title1 'Stochastic Frontier Cost Model';
data electricity;
    input Firm Year Cost Output LPrice LShare KPrice KShare FPrice FShare;
datalines;
1 1955 .0820 2.0 2.090 .3164 183.000 .4521 17.9000 .2315
2 1955 .6610 3.0 2.050 .2073 174.000 .6676 35.1000 .1251
3 1955 .9900 4.0 2.050 .2349 171.000 .5799 35.1000 .1852
4 1955 .3150 4.0 1.830 .1152 166.000 .7857 32.2000 .0990

... more lines ...

/* Data transformations */
data electricity;
    set electricity;
    label Firm="firm index"
           Year="1955 for all observations"
           Cost="Total cost"
           Output="Total output"
           LPrice="Wage rate"
           LShare="Cost share for labor"
           KPrice="Capital price index"
           KShare="Cost share for capital"
           FPrice="Fuel price"
           FShare="Cost share for fuel";
    log_C_PF=log(Cost/FPrice);
    log_PK_PF=log(KPrice/FPrice);
    log_PL_PF=log(LPrice/FPrice);
    log_y=log(Output);
    log_y_sq=log_y**2/2;
run;

```

The following statements estimate a stochastic frontier exponential cost model that uses Christensen and Greene (1976) data:

```

/*-- Stochastic Frontier Cost Model --*/
proc qlim data=electricity;
    model log_C_PF = log_PK_PF log_PL_PF log_y log_y_sq;
    endogenous log_C_PF ~ frontier (type=exponential cost);
run;

```

Output 22.7.2 shows the results.

Output 22.7.2 Exponential Distribution

Stochastic Frontier Cost Model

The QLIM Procedure

Model Fit Summary						
Number of Endogenous Variables		1				
Endogenous Variable		log_C_PF				
Number of Observations		159				
Log Likelihood		-23.30430				
Maximum Absolute Gradient		3.0458E-6				
Number of Iterations		21				
Optimization Method		Quasi-Newton				
AIC		60.60860				
Schwarz Criterion		82.09093				
Sigma		0.30750				
Lambda		1.71345				

Parameter Estimates						
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t	
Intercept	1	-4.983211	0.543328	-9.17	<.0001	
log_PK_PF	1	0.090242	0.109202	0.83	0.4086	
log_PL_PF	1	0.504299	0.118263	4.26	<.0001	
log_y	1	0.427182	0.066680	6.41	<.0001	
log_y_sq	1	0.066120	0.010079	6.56	<.0001	
_Sigma_v	1	0.154998	0.020271	7.65	<.0001	
_Sigma_u	1	0.265581	0.033614	7.90	<.0001	

Similarly, the stochastic frontier cost model can be estimated with (type=half) or (type=truncated) options that represent half-normal and truncated normal errors.

The following statements illustrate the half-normal option:

```

/*-- Stochastic Frontier Cost Model --*/
proc qlim data=electricity;
  model log_C_PF = log_PK_PF log_PL_PF log_y log_y_sq;
  endogenous log_C_PF ~ frontier (type=half cost);
run;

```

Output 22.7.3 shows the result.

Output 22.7.3 Half-Normal Distribution Stochastic Frontier Cost Model

The QLIM Procedure

Model Fit Summary						
Number of Endogenous Variables		1				
Endogenous Variable		log_C_PF				
Number of Observations		159				
Log Likelihood		-34.95304				
Maximum Absolute Gradient		0.0001150				
Number of Iterations		22				
Optimization Method		Quasi-Newton				
AIC		83.90607				
Schwarz Criterion		105.38840				
Sigma		0.42761				
Lambda		1.80031				

Parameter Estimates						
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t	
Intercept	1	-4.434634	0.690197	-6.43	<.0001	
log_PK_PF	1	0.069624	0.136250	0.51	0.6093	
log_PL_PF	1	0.474578	0.146812	3.23	0.0012	
log_y	1	0.256874	0.080777	3.18	0.0015	
log_y_sq	1	0.088051	0.011817	7.45	<.0001	
_Sigma_v	1	0.207637	0.039222	5.29	<.0001	
_Sigma_u	1	0.373810	0.073605	5.08	<.0001	

The following statements illustrate the truncated normal option:

```

/*-- Stochastic Frontier Cost Model --*/
proc qlim data=electricity;
  model log_C_PF = log_PK_PF log_PL_PF log_y log_y_sq;
  endogenous log_C_PF ~ frontier (type=truncated cost);
run;

```

Output 22.7.4 shows the results.

Output 22.7.4 Truncated Normal Distribution**Stochastic Frontier Cost Model****The QLIM Procedure**

Model Fit Summary						
Number of Endogenous Variables			1			
Endogenous Variable			log_C_PF			
Number of Observations			159			
Log Likelihood			-60.32110			
Maximum Absolute Gradient			4225			
Number of Iterations			4			
Optimization Method			Quasi-Newton			
AIC			136.64220			
Schwarz Criterion			161.19343			
Sigma			0.37350			
Lambda			0.70753			

Parameter Estimates						
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t	
Intercept	1	-3.770440	0.839388	-4.49	<.0001	
log_PK_PF	1	-0.045852	0.176682	-0.26	0.7952	
log_PL_PF	1	0.602961	0.191454	3.15	0.0016	
log_y	1	0.094966	0.071124	1.34	0.1818	
log_y_sq	1	0.113010	0.012225	9.24	<.0001	
_Sigma_v	1	0.304905	0.047868	6.37	<.0001	
_Sigma_u	1	0.215728	0.068725	3.14	0.0017	
_Mu	1	0.477097	0.116295	4.10	<.0001	

If no (Production) or (Cost) option is specified, the stochastic frontier production model is estimated by default.

Example 22.8: Bayesian Modeling

This example illustrates how to use the QLIM procedure to perform Bayesian analysis. The generated data mimic a hypothetical scenario in which you study the number of tickets sold for a sports event given the probability of the hosting team winning and the price of the tickets. The following statements create the dataset:

```

title1 'Bayesian Analysis';

ods graphics on;

data test;
  do i=1 to 200;
    e1 = rannor(8726)*2000;
    WinChance = ranuni(8772);
    Price = 10+ranexp(8773)*4;
    y = 48000 + 5000*WinChance - 100 * price + e1;
    if y>50000 then TicketSales = 50000;
    if y<=50000 then TicketSales = y;
    output;
  end;
  keep WinChance price y TicketSales;
run;

```

The following statements perform Bayesian analysis of a Tobit model:

```

proc qlim data=test plots(prior)=all;
  model TicketSales = WinChance price;
  endogenous TicketSales ~ censored(lb=0 ub= 50000);
  prior intercept~normal(mean=48000);
  prior WinChance~normal(mean=5000);
  prior Price~normal(mean=-100);
  bayes NBI=10000 NMC=30000 THIN=1 ntrds=1 DIAG=ALL STATS=ALL seed=2;
run;

```

Output 22.8.1 shows the results from the maximum likelihood estimation and the Bayesian analysis with diffuse prior of this Tobit model.

Output 22.8.1 Bayesian Tobit Model

Bayesian Analysis


The QLIM Procedure

Parameter Estimates					
Parameter	DF	Estimate	Standard Error	t Value	Approx Pr > t
Intercept	1	48119	623.565045	77.17	<.0001
WinChance	1	5242.083501	559.151222	9.38	<.0001
Price	1	-106.731665	40.660795	-2.62	0.0087
_Sigma	1	1939.607206	134.348772	14.44	<.0001



Output 22.8.1 *continued*


Posterior Summaries				Percentiles		
Parameter	N	Mean	Standard Deviation	25%	50%	75%
Intercept	30000	48109.4	535.0	47750.5	48102.6	48460.1
WinChance	30000	5212.9	483.4	4878.8	5205.2	5533.0
Price	30000	-104.7	36.5224	-128.6	-104.2	-79.4191
_Sigma	30000	1950.9	132.9	1858.4	1945.0	2034.0

Output 22.8.2 depicts a graphical representation of MLE, prior, and posterior distributions.

Output 22.8.2 Predictive Analysis by Observation Number


Output 22.8.2 continued





Output 22.8.2 *continued*


The validity of the MCMC sampling phase can be monitored with [Output 22.8.3](#).

Output 22.8.3 Predictive Analysis by Observation Number




Output 22.8.3 continued



Output 22.8.3 continued

Finally the prior and the posterior predictive analyses are represented in [Output 22.8.4](#)

Output 22.8.4 Predictive Analysis by Observation Number

References

- Abramowitz, M. and Stegun, A. (1970), *Handbook of Mathematical Functions*, New York: Dover Press.
- Aigner, C., Lovell, C. A. K., and Schmidt, P. (1977), "Formulation and Estimation of Stochastic Frontier Production Function Models," *Journal of Econometrics*, 6, 21–37.
- Aitchison, J. and Silvey, S. (1957), "The Generalization of Probit Analysis to the Case of Multiple Responses," *Biometrika*, 44, 131–140.
- Amemiya, T. (1978a), "The Estimation of a Simultaneous Equation Generalized Probit Model," *Econometrica*, 46, 1193–1205.
- Amemiya, T. (1978b), "On a Two-Step Estimate of a Multivariate Logit Model," *Journal of Econometrics*, 8, 13–21.
- Amemiya, T. (1981), "Qualitative Response Models: A Survey," *Journal of Economic Literature*, 19, 483–536.
- Amemiya, T. (1984), "Tobit Models: A Survey," *Journal of Econometrics*, 24, 3–61.
- Amemiya, T. (1985), *Advanced Econometrics*, Cambridge, MA: Harvard University Press.
- Battese, G. E. and Coelli, T. J. (1988), "Prediction of Firm-Level Technical Efficiencies with a Generalized Frontier Production Function and Panel Data," *Journal of Econometrics*, 38, 387–399.
- Ben-Akiva, M. and Lerman, S. R. (1985), *Discrete Choice Analysis: Theory and Application to Travel Demand*, MIT Press Series in Transportation Studies, Cambridge, MA: MIT Press.
- Bera, A. K., Jarque, C. M., and Lee, L.-F. (1984), "Testing the Normality Assumption in Limited Dependent Variable Models," *International Economic Review*, 25, 563–578.
- Berger, J. O. (1985), *Statistical Decision Theory and Bayesian Analysis*, 2nd Edition, New York: Springer-Verlag.
- Bloom, D. E. and Killingsworth, M. R. (1985), "Correcting for Truncation Bias Caused by a Latent Truncation Variable," *Journal of Econometrics*, 27, 131–135.
- Box, G. E. P. and Cox, D. R. (1964), "An Analysis of Transformations," *Journal of the Royal Statistical Society, Series B*, 26, 211–234.
- Cameron, A. C. and Trivedi, P. K. (1986), "Econometric Models Based on Count Data: Comparisons and Applications of Some Estimators and Tests," *Journal of Applied Econometrics*, 1, 29–53.
- Cameron, A. C. and Trivedi, P. K. (1998), *Regression Analysis of Count Data*, Cambridge: Cambridge University Press.
- Christensen, L. R. and Greene, W. H. (1976), "Economies of Scale in U.S. Electric Power Generation," *Journal of Political Economy*, 84, 655–676.
- Coelli, T. J., Prasada Rao, D. S., and Battese, G. E. (1998), *An Introduction to Efficiency and Productivity Analysis*, London: Kluwer Academic.

- Copley, P. A., Doucet, M. S., and Gaver, K. M. (1994), "A Simultaneous Equations Analysis of Quality Control Review Outcomes and Engagement Fees for Audits of Recipients of Federal Financial Assistance," *Accounting Review*, 69, 244–256.
- Cox, D. R. (1970), *Analysis of Binary Data*, London: Methuen.
- Cox, D. R. (1972), "Regression Models and Life Tables," *Journal of the Royal Statistical Society, Series B*, 20, 187–220, with discussion.
- Cox, D. R. (1975), "Partial Likelihood," *Biometrika*, 62, 269–276.
- Deis, D. R. and Hill, R. C. (1998), "An Application of the Bootstrap Method to the Simultaneous Equations Model of the Demand and Supply of Audit Services," *Contemporary Accounting Research*, 15, 83–99.
- Estrella, A. (1998), "A New Measure of Fit for Equations with Dichotomous Dependent Variables," *Journal of Business and Economic Statistics*, 16, 198–205.
- Gallant, A. R. (1987), *Nonlinear Statistical Models*, New York: John Wiley & Sons.
- Gelman, A., Carlin, J. B., Stern, H. S., and Rubin, D. B. (2004), *Bayesian Data Analysis*, 2nd Edition, London: Chapman & Hall.
- Genz, A. (1992), "Numerical Computation of Multivariate Normal Probabilities," *Journal of Computational and Graphical Statistics*, 1, 141–150.
- Godfrey, L. G. (1988), *Misspecification Tests in Econometrics*, Cambridge: Cambridge University Press.
- Gourieroux, C., Monfort, A., Renault, E., and Trognon, A. (1987), "Generalized Residuals," *Journal of Econometrics*, 34, 5–32.
- Greene, W. H. (1997), *Econometric Analysis*, 3rd Edition, Upper Saddle River, NJ: Prentice-Hall.
- Gregory, A. W. and Veall, M. R. (1985), "On Formulating Wald Tests for Nonlinear Restrictions," *Econometrica*, 53, 1465–1468.
- Hajivassiliou, V. A. (1993), "Simulation Estimation Methods for Limited Dependent Variable Models," in G. S. Maddala, C. R. Rao, and H. D. Vinod, eds., *Econometrics*, volume 11 of *Handbook of Statistics*, New York: Elsevier Science.
- Hajivassiliou, V. A. and McFadden, D. (1998), "The Method of Simulated Scores for the Estimation of LDV Models," *Econometrica*, 66, 863–896.
- Hajivassiliou, V. A., McFadden, D. L., and Ruud, P. A. (1996), "Simulation of Multivariate Normal Rectangle Probabilities and Their Derivatives: Theoretical and Computational Results," *Journal of Econometrics*, 72, 85–134.
- Heckman, J. J. (1978), "Dummy Endogenous Variables in a Simultaneous Equation System," *Econometrica*, 46, 931–959.
- Hinkley, D. V. (1975), "On Power Transformations to Symmetry," *Biometrika*, 62, 101–111.
- Jondrow, J., Lovell, C. A. K., Materov, I. S., and Schmidt, P. (1982), "On the Estimation of Technical Efficiency in the Stochastic Frontier Production Function Model," *Journal of Econometrics*, 19, 233–238.

- Kim, M. and Hill, R. C. (1993), "The Box-Cox Transformation-of-Variables in Regression," *Empirical Economics*, 18, 307–319.
- Kumbhakar, S. C. and Lovell, C. A. K. (2000), *Stochastic Frontier Analysis*, New York: Cambridge University Press.
- Lee, L.-F. (1981), "Simultaneous Equations Models with Discrete and Censored Dependent Variables," in C. F. Manski and D. McFadden, eds., *Structural Analysis of Discrete Data with Econometric Applications*, Cambridge, MA: MIT Press.
- Long, J. S. (1997), *Regression Models for Categorical and Limited Dependent Variables*, Thousand Oaks, CA: Sage Publications.
- McFadden, D. (1974), "Conditional Logit Analysis of Qualitative Choice Behavior," in P. Zarembka, ed., *Frontiers in Econometrics*, New York: Academic Press.
- McFadden, D. (1981), "Econometric Models of Probabilistic Choice," in C. F. Manski and D. McFadden, eds., *Structural Analysis of Discrete Data with Econometric Applications*, Cambridge, MA: MIT Press.
- McKelvey, R. D. and Zavoina, W. (1975), "A Statistical Model for the Analysis of Ordinal Level Dependent Variables," *Journal of Mathematical Sociology*, 4, 103–120.
- Meeusen, W. and van den Broeck, J. (1977), "Efficiency Estimation from Cobb-Douglas Production Functions with Composed Error," *International Economic Review*, 18, 435–444.
- Mroz, T. A. (1987), "The Sensitivity of an Empirical Model of Married Women's Work to Economic and Statistical Assumptions," *Econometrica*, 55, 765–799.
- Mroz, T. A. (1999), "Discrete Factor Approximations in Simultaneous Equation Models: Estimating the Impact of a Dummy Endogenous Variable on a Continuous Outcome," *Journal of Econometrics*, 92, 233–274.
- Nawata, K. (1994), "Estimation of Sample Selection Bias Models by the Maximum Likelihood Estimator and Heckman's Two-Step Estimator," *Economics Letters*, 45, 33–40.
- Parks, R. W. (1967), "Efficient Estimation of a System of Regression Equations When Disturbances Are Both Serially and Contemporaneously Correlated," *Journal of the American Statistical Association*, 62, 500–509.
- Phillips, C. B. and Park, J. Y. (1988), "On Formulating Wald Tests of Nonlinear Restrictions," *Econometrica*, 56, 1065–1083.
- Powers, D. A. and Xie, Y. (2000), *Statistical Methods for Categorical Data Analysis*, San Diego: Academic Press.
- Rivers, D. and Vuong, Q. H. (1988), "Limited Information Estimators and Exogeneity Tests for Simultaneous Probit Models," *Journal of Econometrics*, 39, 347–366.
- Roberts, G. O., Gelman, A., and Gilks, W. R. (1997), "Weak Convergence and Optimal Scaling of Random Walk Metropolis Algorithms," *Annals of Applied Probability*, 7, 110–120.
- Roberts, G. O. and Rosenthal, J. S. (2001), "Optimal Scaling for Various Metropolis-Hastings Algorithms," *Statistical Science*, 16, 351–367.

Schervish, M. J. (1995), *Theory of Statistics*, New York: Springer-Verlag.

Wooldridge, J. M. (2002), *Econometric Analysis of Cross Section and Panel Data*, Cambridge, MA: MIT Press.

Subject Index

- bounds on parameter estimates, [1541](#)
- BOUNDS statement, [1541](#)
- censored regression models
 - QLIM procedure, [1557](#)
- covariates
 - heteroscedasticity models, [1546](#)
- gamma distribution
 - definition of (QLIM), [1589](#)
 - QLIM procedure, [1589](#)
- Gaussian distribution
 - definition of (QLIM), [1589](#)
 - QLIM procedure, [1589](#)
- heteroscedasticity models
 - covariates, [1546](#)
- initial values
 - QLIM procedure, [1579](#)
- inverse-gamma distribution
 - definition of (QLIM), [1589](#)
 - QLIM procedure, [1589](#)
- Lagrange multiplier test
 - nonlinear hypotheses, [1570](#)
- limited dependent variable models
 - QLIM procedure, [1557](#)
- logit
 - QLIM Procedure, [1520](#)
- maximum a posteriori
 - QLIM procedure, [1579](#)
- nonlinear hypotheses
 - Lagrange multiplier test, [1570](#)
 - Wald test, [1570](#)
- normal distribution
 - definition of (QLIM), [1589](#)
 - QLIM procedure, [1589](#)
- ordinal discrete choice modeling
 - QLIM procedure, [1554](#)
- output ODS Graphics table names
 - QLIM procedure, [1598](#)
- output table names
 - QLIM procedure, [1597](#)
- prior distribution
 - distribution specification (QLIM), [1551](#)
- probit
 - QLIM Procedure, [1520](#)
- QLIM Procedure, [1520](#)
 - logit, [1520](#)
 - probit, [1520](#)
 - selection, [1520](#)
 - Tobit model, [1520](#)
- QLIM procedure
 - bivariate limited dependent variable modeling, [1564](#)
 - Box-Cox modeling, [1564](#)
 - BY groups, [1542](#)
 - censored regression models, [1557](#)
 - frontier, [1561](#)
 - gamma distribution, [1589](#)
 - Gaussian distribution, [1589](#)
 - heteroscedasticity, [1563](#)
 - initial values, [1579](#)
 - inverse-gamma distribution , [1589](#)
 - limited dependent variable models, [1557](#)
 - maximum a posteriori, [1579](#)
 - multivariate limited dependent models, [1569](#)
 - normal distribution, [1589](#)
 - ordinal discrete choice modeling, [1554](#)
 - output, [1590](#)
 - output ODS Graphics table names, [1598](#)
 - output table names, [1597](#)
 - selection models, [1565](#)
 - standard distributions, [1588](#)
 - syntax, [1527](#)
 - t distribution, [1590](#)
 - tests on parameters, [1569](#)
 - truncated regression models, [1560](#)
 - types of Tobit model, [1558](#)
 - uniform distribution, [1590](#)
- selection
 - QLIM Procedure, [1520](#)
- standard distributions
 - QLIM procedure, [1588](#)
- t distribution
 - definition of (QLIM), [1590](#)
 - QLIM procedure, [1590](#)
- Tobit model
 - QLIM Procedure, [1520](#)
- truncated regression models
 - QLIM procedure, [1560](#)

types of Tobit model

QLIM procedure, [1558](#)

uniform distribution

definition of (QLIM), [1590](#)

QLIM procedure, [1590](#)

Wald test

nonlinear hypotheses, [1570](#)

Syntax Index

- ACCURACY= option
 - BAYES statement (QLIM), 1536
- AGGREGATION= option
 - BAYES statement (QLIM), 1536
- ALL option
 - TEST statement (QLIM), 1552
- ATTEMPTS= number
 - BAYES statement (QLIM), 1536, 1537
- AUTOMCMC option
 - BAYES statement (QLIM), 1536
- BAYES statement
 - QLIM procedure, 1536
- BCX option
 - MODEL statement (QLIM), 1548
- BETA
 - PRIOR statement (QLIM), 1551
- BOUNDS statement
 - QLIM procedure, 1541
- BY statement
 - QLIM procedure, 1542
- CENSORED option
 - ENDOGENOUS statement (QLIM), 1543
- CONDITIONAL
 - OUTPUT statement (QLIM), 1550
- CORRB option
 - QLIM procedure, 1531
- CORROUT option
 - PROC QLIM statement, 1531
- COST option
 - ENDOGENOUS statement (QLIM), 1544
- COVB option
 - QLIM procedure, 1531
- COVEST= option
 - QLIM procedure, 1532
- COVOUT option
 - PROC QLIM statement, 1531
- DATA= option
 - PROC QLIM statement, 1531
- DIAGNOSTICS= option
 - BAYES statement (QLIM), 1537
- DISCRETE option
 - ENDOGENOUS statement (QLIM), 1543
- DISTRIBUTION= option
 - ENDOGENOUS statement (QLIM), 1543
- ENDOTEST option
 - ENDOGENOUS statement (QLIM), 1545
- ERRSTD
 - OUTPUT statement (QLIM), 1550
- EXPECTED
 - OUTPUT statement (QLIM), 1550
- FRONTIER option
 - ENDOGENOUS statement (QLIM), 1544
- GAMMA
 - PRIOR statement (QLIM), 1551
- HECKIT option
 - QLIM procedure, 1532
- IGAMMA
 - PRIOR statement (QLIM), 1551
- INIT statement
 - QLIM procedure, 1547
- ITPRINT option
 - QLIM procedure, 1531
- LIMIT1= option
 - MODEL statement (QLIM), 1548
- LM option
 - TEST statement (QLIM), 1553
- LOWERBOUND= option
 - ENDOGENOUS statement (QLIM), 1544
- LR option
 - TEST statement (QLIM), 1553
- MARGINAL
 - OUTPUT statement (QLIM), 1550
- MAXNMCMC=number
 - BAYES statement (QLIM), 1537
- MAXTUNE= option
 - BAYES statement (QLIM), 1539
- METHOD= option
 - QLIM procedure, 1533
- MILLS
 - OUTPUT statement (QLIM), 1550
- MINTUNE= option
 - BAYES statement (QLIM), 1539
- MODEL statement
 - QLIM procedure, 1548
- MULTIPLIER= value
 - BAYES statement (QLIM), 1537
- NBI= option

- BAYES statement (QLIM), 1539
- NDRAW= option
 - QLIM procedure, 1533
- NLOPTIONS statement
 - QLIM procedure, 1549
- NMC= option
 - BAYES statement (QLIM), 1539
- NMCPRIOR= option
 - BAYES statement (QLIM), 1539
- NOINT option
 - MODEL statement (QLIM), 1548
- NONORMALIZE option
 - WEIGHT statement (QLIM), 1554
- NOPRINT option
 - PROC QLIM statement, 1531
- NORMAL
 - PRIOR statement (QLIM), 1551
- NTRDS= option
 - BAYES statement (QLIM), 1539
- NTU= option
 - BAYES statement (QLIM), 1539
- ORDER= option
 - ENDOGENOUS statement (QLIM), 1543
- OUT= option
 - OUTPUT statement (QLIM), 1550
- OUTEST= option
 - PROC QLIM statement, 1531
- OUTPOST= option
 - BAYES statement (QLIM), 1539
- OUTPRIOR= option
 - BAYES statement (QLIM), 1539
- OUTPUT statement
 - QLIM procedure, 1550
- OVERID option
 - ENDOGENOUS statement (QLIM), 1546
- PLOTS option
 - QLIM statement (QLIM), 1533
- PREDICTED
 - OUTPUT statement (QLIM), 1550
- PRINTALL option
 - PROC QLIM statement, 1531
- PRIOR statement
 - QLIM procedure, 1551
- PROB
 - OUTPUT statement (QLIM), 1550
- PROBALL
 - OUTPUT statement (QLIM), 1550
- PRODUCTION option
 - ENDOGENOUS statement (QLIM), 1544
- PROPCOV= option
 - BAYES statement (QLIM), 1540
- PROPORTION= value

- BAYES statement (QLIM), 1537
- QLIM procedure, 1527
 - PRIOR statement, 1551
 - syntax, 1527
- QLIM procedure, CLASS statement, 1542
- QLIM procedure, FREQ statement, 1546
- QLIM procedure, TEST statement, 1552
- QLIM procedure, WEIGHT statement, 1554
- RANDINIT option
 - BAYES statement (QLIM), 1537
- RESIDUAL
 - OUTPUT statement (QLIM), 1550
- RESTRICT statement
 - QLIM procedure, 1551
- RLLIMITS= number
 - BAYES statement (QLIM), 1536
- SAMPLING= option
 - BAYES statement (QLIM), 1540
- SEED= option
 - BAYES statement (QLIM), 1540
 - QLIM procedure, 1533
- SELECT option
 - ENDOGENOUS statement (QLIM), 1545
- SELECTVAR option
 - MODEL statement (QLIM), 1549
- SIMTIME option
 - BAYES statement (QLIM), 1540
- STATIONARITY= option
 - BAYES statement (QLIM), 1537
- STATISTICS option
 - BAYES statement (QLIM), 1540
- T
 - PRIOR statement (QLIM), 1551
- TARGETESS= number
 - BAYES statement (QLIM), 1536
- TARGETSTATS<= option>
 - BAYES statement (QLIM), 1536
- TE1
 - OUTPUT statement (QLIM), 1550
- TE2
 - OUTPUT statement (QLIM), 1551
- THIN= option
 - BAYES statement (QLIM), 1541
- TOL= value
 - BAYES statement (QLIM), 1537
- TRUNCATED option
 - ENDOGENOUS statement (QLIM), 1544
- UNIFORM
 - PRIOR statement (QLIM), 1551
- UPPERBOUND= option

ENDOGENOUS statement (QLIM), [1544](#)

WALD option

TEST statement (QLIM), [1552](#)

XBETA

OUTPUT statement (QLIM), [1550](#)