



THE  
POWER  
TO KNOW.

# **SAS/STAT® 9.22 User's Guide**

## **Introduction to Discriminant**

### **Procedures**

#### **(Book Excerpt)**



This document is an individual chapter from *SAS/STAT® 9.22 User's Guide*.

The correct bibliographic citation for the complete manual is as follows: SAS Institute Inc. 2010. *SAS/STAT® 9.22 User's Guide*. Cary, NC: SAS Institute Inc.

Copyright © 2010, SAS Institute Inc., Cary, NC, USA

All rights reserved. Produced in the United States of America.

**For a Web download or e-book:** Your use of this publication shall be governed by the terms established by the vendor at the time you acquire this publication.

**U.S. Government Restricted Rights Notice:** Use, duplication, or disclosure of this software and related documentation by the U.S. government is subject to the Agreement with SAS Institute and the restrictions set forth in FAR 52.227-19, Commercial Computer Software-Restricted Rights (June 1987).

SAS Institute Inc., SAS Campus Drive, Cary, North Carolina 27513.

1st electronic book, May 2010

SAS® Publishing provides a complete selection of books and electronic products to help customers use SAS software to its fullest potential. For more information about our e-books, e-learning products, CDs, and hard-copy books, visit the SAS Publishing Web site at [support.sas.com/publishing](http://support.sas.com/publishing) or call 1-800-727-3228.

SAS® and all other SAS Institute Inc. product or service names are registered trademarks or trademarks of SAS Institute Inc. in the USA and other countries. ® indicates USA registration.

Other brand and product names are registered trademarks or trademarks of their respective companies.

# Chapter 10

## Introduction to Discriminant Procedures

### Contents

Overview: Discriminant Procedures . . . . .	199
Background: Discriminant Procedures . . . . .	200
Example: Contrasting Univariate and Multivariate Analyses . . . . .	201
References . . . . .	206

---

### Overview: Discriminant Procedures

The SAS procedures for discriminant analysis fit data with one classification variable and several quantitative variables. The purpose of discriminant analysis can be to find one or more of the following:

- a mathematical rule, or *discriminant function*, for guessing to which class an observation belongs, based on knowledge of the quantitative variables only
- a set of linear combinations of the quantitative variables that best reveals the differences among the classes
- a subset of the quantitative variables that best reveals the differences among the classes

The SAS discriminant procedures are as follows:

DISCRIM	computes various discriminant functions for classifying observations. Linear or quadratic discriminant functions can be used for data with approximately multivariate normal within-class distributions. Nonparametric methods can be used without making any assumptions about these distributions.
CANDISC	performs a canonical analysis to find linear combinations of the quantitative variables that best summarize the differences among the classes.
STEPDISC	uses forward selection, backward elimination, or stepwise selection to try to find a subset of quantitative variables that best reveals differences among the classes.

---

## Background: Discriminant Procedures

The term *discriminant analysis* (Fisher 1936; Cooley and Lohnes 1971; Tatsuoaka 1971; Kshirsagar 1972; Lachenbruch 1975, 1979; Gnanadesikan 1977; Klecka 1980; Hand 1981, 1982; Silverman 1986) refers to several different types of analyses. Classificatory discriminant analysis is used to classify observations into two or more known groups on the basis of one or more quantitative variables. Classification can be done by either a parametric method or a nonparametric method in the DISCRIM procedure. A parametric method is appropriate only for approximately normal within-class distributions. The method generates either a linear discriminant function (the within-class covariance matrices are assumed to be equal) or a quadratic discriminant function (the within-class covariance matrices are assumed to be unequal).

When the distribution within each group is not assumed to have any specific distribution or is assumed to have a distribution different from the multivariate normal distribution, nonparametric methods can be used to derive classification criteria. These methods include the kernel method and nearest-neighbor methods. The kernel method uses uniform, normal, Epanechnikov, biweight, or triweight kernels in estimating the group-specific density at each observation. The within-group covariance matrices or the pooled covariance matrix can be used to scale the data.

The performance of a discriminant function can be evaluated by estimating error rates (probabilities of misclassification). Error count estimates and posterior probability error rate estimates can be evaluated with PROC DISCRIM. When the input data set is an ordinary SAS data set, the error rates can also be estimated by cross validation.

In multivariate statistical applications, the data collected are largely from distributions different from the normal distribution. Various forms of nonnormality can arise, such as qualitative variables or variables with underlying continuous but nonnormal distributions. If the multivariate normality assumption is violated, the use of parametric discriminant analysis might not be appropriate. When a parametric classification criterion (linear or quadratic discriminant function) is derived from a nonnormal population, the resulting error rate estimates might be biased.

If your quantitative variables are not normally distributed, or if you want to classify observations on the basis of categorical variables, you should consider using the CATMOD or LOGISTIC procedure to fit a categorical linear model with the classification variable as the dependent variable. Press and Wilson (1978) compare logistic regression and parametric discriminant analysis and conclude that logistic regression is preferable to parametric discriminant analysis in cases for which the variables do not have multivariate normal distributions within classes. However, if you do have normal within-class distributions, logistic regression is less efficient than parametric discriminant analysis. Efron (1975) shows that with two normal populations having a common covariance matrix, logistic regression is between one-half and two-thirds as effective as the linear discriminant function in achieving asymptotically the same error rate.

Do not confuse discriminant analysis with cluster analysis. All varieties of discriminant analysis require prior knowledge of the classes, usually in the form of a sample from each class. In cluster analysis, the data do not include information about class membership; the purpose is to construct a classification. See Chapter 11, [“Introduction to Clustering Procedures.”](#)



Canonical discriminant analysis is a dimension-reduction technique related to principal components and canonical correlation, and it can be performed by both the CANDISC and DISCRIM procedures. A discriminant criterion is always derived in PROC DISCRIM. If you want canonical discriminant analysis without the use of a discriminant criterion, you should use PROC CANDISC. Stepwise discriminant analysis is a variable-selection technique implemented by the STEPDISC procedure. After selecting a subset of variables with PROC STEPDISC, use any of the other discriminant procedures to obtain more detailed analyses. PROC CANDISC and PROC STEPDISC perform hypothesis tests that require the within-class distributions to be approximately normal, but these procedures can be used descriptively with nonnormal data.

Another alternative to discriminant analysis is to perform a series of univariate one-way ANOVAs. All three discriminant procedures provide summaries of the univariate ANOVAs. The advantage of the multivariate approach is that two or more classes that overlap considerably when each variable is viewed separately might be more distinct when examined from a multivariate point of view.

---

## Example: Contrasting Univariate and Multivariate Analyses

Consider an artificial data set with two classes of observations indicated by 'H' and 'O'. The following statements generate and plot the data:

```
data random;
  drop n;

  Group = 'H';
  do n = 1 to 20;
    x = 4.5 + 2 * normal(57391);
    y = x + .5 + normal(57391);
    output;
  end;


  Group = 'O';
  do n = 1 to 20;
    x = 6.25 + 2 * normal(57391);
    y = x - 1 + normal(57391);
    output;
  end;

run;

proc sgplot noautolegend;
  scatter y=y x=x / markerchar=group group=group;
run;
```

The plot is shown in [Figure 10.1](#).

**Figure 10.1** Groups for Contrasting Univariate and Multivariate Analyses



The following statements perform a canonical discriminant analysis and display the results in Figure 10.2:

```
proc candisc anova;  
  class Group;  
  var x y;  
run;
```

**Figure 10.2** Contrasting Univariate and Multivariate Analyses

The CANDISC Procedure			
Total Sample Size	40	DF Total	39
Variables	2	DF Within Classes	38
Classes	2	DF Between Classes	1
Number of Observations Read		40	
Number of Observations Used		40	

Figure 10.2 continued

Class Level Information							
Group	Variable Name	Frequency	Weight	Proportion			
H	H	20	20.0000	0.500000			
O	O	20	20.0000	0.500000			
The CANDISC Procedure							
Univariate Test Statistics							
F Statistics, Num DF=1, Den DF=38							
Variable	Total Standard Deviation	Pooled Standard Deviation	Between Standard Deviation	R-Square	R-Square / (1-RSq)	F Value	Pr > F
x	2.1776	2.1498	0.6820	0.0503	0.0530	2.01	0.1641
y	2.4215	2.4486	0.2047	0.0037	0.0037	0.14	0.7105
Average R-Square							
Unweighted				0.0269868			
Weighted by Variance				0.0245201			
Multivariate Statistics and Exact F Statistics							
S=1 M=0 N=17.5							
Statistic	Value	F Value	Num DF	Den DF	Pr > F		
Wilks' Lambda	0.64203704	10.31	2	37	0.0003		
Pillai's Trace	0.35796296	10.31	2	37	0.0003		
Hotelling-Lawley Trace	0.55754252	10.31	2	37	0.0003		
Roy's Greatest Root	0.55754252	10.31	2	37	0.0003		

Figure 10.2 continued

The CANDISC Procedure					
	Canonical Correlation	Adjusted Canonical Correlation	Approximate Standard Error	Squared Canonical Correlation	
1	0.598300	0.589467	0.102808	0.357963	
Eigenvalues of Inv(E)*H = CanRsqr/(1-CanRsqr)					
	Eigenvalue	Difference	Proportion	Cumulative	
1	0.5575		1.0000	1.0000	
Test of H0: The canonical correlations in the current row and all that follow are zero					
	Likelihood Ratio	Approximate F Value	Num DF	Den DF	Pr > F
1	0.64203704	10.31	2	37	0.0003
NOTE: The F statistic is exact.					
The CANDISC Procedure					
Total Canonical Structure					
Variable		Can1			
x		-0.374883			
y		0.101206			
Between Canonical Structure					
Variable		Can1			
x		-1.000000			
y		1.000000			
Pooled Within Canonical Structure					
Variable		Can1			
x		-0.308237			
y		0.081243			

**Figure 10.2** *continued*

The CANDISC Procedure	
Total-Sample Standardized Canonical Coefficients	
Variable	Can1
x	-2.625596855
y	2.446680169
Pooled Within-Class Standardized Canonical Coefficients	
Variable	Can1
x	-2.592150014
y	2.474116072
Raw Canonical Coefficients	
Variable	Can1
x	-1.205756217
y	1.010412967
Class Means on Canonical Variables	
Group	Can1
H	0.7277811475
O	-.7277811475

The univariate R squares are very small, 0.0503 for x and 0.0037 for y, and neither variable shows a significant difference between the classes at the 0.10 level.

The multivariate test for differences between the classes is significant at the 0.0003 level. Thus, the multivariate analysis has found a highly significant difference, whereas the univariate analyses failed to achieve even the 0.10 level. The raw canonical coefficients for the first canonical variable, Can1, show that the classes differ most widely on the linear combination  $-1.205756217x + 1.010412967y$  or approximately  $y - 1.2x$ . The R square between Can1 and the class variable is 0.357963 as given by the squared canonical correlation, which is much higher than either univariate R square.

In this example, the variables are highly correlated within classes. If the within-class correlation were smaller, there would be greater agreement between the univariate and multivariate analyses.

---

## References

- Cooley, W. W. and Lohnes, P. R. (1971), *Multivariate Data Analysis*, New York: John Wiley & Sons.
- Efron, B. (1975), "The Efficiency of Logistic Regression Compared to Normal Discriminant Analysis," *Journal of the American Statistical Association*, 70, 892–898.
- Fisher, R. A. (1936), "The Use of Multiple Measurements in Taxonomic Problems," *Annals of Eugenics*, 7, 179–188.
- Gnanadesikan, R. (1977), *Methods for Statistical Data Analysis of Multivariate Observations*, New York: John Wiley & Sons.
- Hand, D. J. (1981), *Discrimination and Classification*, New York: John Wiley & Sons.
- Hand, D. J. (1982), *Kernel Discriminant Analysis*, New York: Research Studies Press.
- Klecka, W. R. (1980), *Discriminant Analysis*, Sage University Paper Series on Quantitative Applications in the Social Sciences, 07-019, Beverly Hills and London: Sage Publications.
- Kshirsagar, A. M. (1972), *Multivariate Analysis*, New York: Marcel Dekker.
- Lachenbruch, P. A. (1975), *Discriminant Analysis*, New York: Hafner Publishing.
- Lachenbruch, P. A. (1979), "Discriminant Analysis," *Biometrics*.
- Press, S. J. and Wilson, S. (1978), "Choosing between Logistic Regression and Discriminant Analysis," *Journal of the American Statistical Association*, 73, 699–705.
- Silverman, B. W. (1986), *Density Estimation for Statistics and Data Analysis*, New York: Chapman & Hall.
- Tatsuoka, M. M. (1971), *Multivariate Analysis*, New York: John Wiley & Sons.

## Your Turn

---

We welcome your feedback.

- If you have comments about this book, please send them to **`yourturn@sas.com`**. Include the full title and page numbers (if applicable).
- If you have comments about the software, please send them to **`suggest@sas.com`**.





# SAS® Publishing Delivers!

Whether you are new to the work force or an experienced professional, you need to distinguish yourself in this rapidly changing and competitive job market. SAS® Publishing provides you with a wide range of resources to help you set yourself apart. Visit us online at [support.sas.com/bookstore](http://support.sas.com/bookstore).

## SAS® Press

Need to learn the basics? Struggling with a programming problem? You'll find the expert answers that you need in example-rich books from SAS Press. Written by experienced SAS professionals from around the world, SAS Press books deliver real-world insights on a broad range of topics for all skill levels.

**[support.sas.com/saspress](http://support.sas.com/saspress)**

## SAS® Documentation

To successfully implement applications using SAS software, companies in every industry and on every continent all turn to the one source for accurate, timely, and reliable information: SAS documentation. We currently produce the following types of reference documentation to improve your work experience:

- Online help that is built into the software.
- Tutorials that are integrated into the product.
- Reference documentation delivered in HTML and PDF – **free** on the Web.
- Hard-copy books.

**[support.sas.com/publishing](http://support.sas.com/publishing)**

## SAS® Publishing News

Subscribe to SAS Publishing News to receive up-to-date information about all new SAS titles, author podcasts, and new Web site features via e-mail. Complete instructions on how to subscribe, as well as access to past issues, are available at our Web site.

**[support.sas.com/spn](http://support.sas.com/spn)**



**THE  
POWER  
TO KNOW®**

