



THE
POWER
TO KNOW.

SAS/STAT[®] 9.22 User's Guide

The SURVEYMEANS Procedure

(Book Excerpt)



This document is an individual chapter from *SAS/STAT® 9.22 User's Guide*.

The correct bibliographic citation for the complete manual is as follows: SAS Institute Inc. 2010. *SAS/STAT® 9.22 User's Guide*. Cary, NC: SAS Institute Inc.

Copyright © 2010, SAS Institute Inc., Cary, NC, USA

All rights reserved. Produced in the United States of America.

For a Web download or e-book: Your use of this publication shall be governed by the terms established by the vendor at the time you acquire this publication.

U.S. Government Restricted Rights Notice: Use, duplication, or disclosure of this software and related documentation by the U.S. government is subject to the Agreement with SAS Institute and the restrictions set forth in FAR 52.227-19, Commercial Computer Software-Restricted Rights (June 1987).

SAS Institute Inc., SAS Campus Drive, Cary, North Carolina 27513.

1st electronic book, May 2010

SAS® Publishing provides a complete selection of books and electronic products to help customers use SAS software to its fullest potential. For more information about our e-books, e-learning products, CDs, and hard-copy books, visit the SAS Publishing Web site at support.sas.com/publishing or call 1-800-727-3228.

SAS® and all other SAS Institute Inc. product or service names are registered trademarks or trademarks of SAS Institute Inc. in the USA and other countries. ® indicates USA registration.

Other brand and product names are registered trademarks or trademarks of their respective companies.

Chapter 86

The SURVEYMEANS Procedure

Contents

Overview: SURVEYMEANS Procedure	7240
Getting Started: SURVEYMEANS Procedure	7241
Simple Random Sampling	7241
Stratified Sampling	7243
Output Data Sets	7246
Syntax: SURVEYMEANS Procedure	7247
PROC SURVEYMEANS Statement	7248
BY Statement	7257
CLASS Statement	7258
CLUSTER Statement	7258
DOMAIN Statement	7259
RATIO Statement	7260
REPWEIGHTS Statement	7262
STRATA Statement	7263
VAR Statement	7264
WEIGHT Statement	7264
Details: SURVEYMEANS Procedure	7265
Missing Values	7265
Survey Data Analysis	7266
Specification of Population Totals and Sampling Rates	7266
Primary Sampling Units (PSUs)	7267
Domain Analysis	7267
Statistical Computations	7267
Definitions and Notation	7268
Mean	7269
Variance and Standard Error of the Mean	7270
<i>t</i> Test for the Mean	7271
Degrees of Freedom	7271
Confidence Limits for the Mean	7272
Coefficient of Variation	7273
Proportions	7273
Total	7273
Variance and Standard Deviation of the Total	7273
Confidence Limits for the Total	7274

Ratio	7275
Domain Statistics	7276
Quantiles	7278
Replication Methods for Variance Estimation	7280
Balanced Repeated Replication (BRR) Method	7280
Fay's BRR Method	7282
Jackknife Method	7282
Hadamard Matrix	7284
Output Data Sets	7284
Replicate Weights Output Data Set	7284
Jackknife Coefficients Output Data Set	7285
Rectangular and Stacking Structures in an Output Data Set	7285
Displayed Output	7287
Data and Sample Design Summary	7287
Class Level Information	7287
Stratum Information	7288
Variance Estimation	7288
Statistics	7289
Quantiles	7290
Domain Analysis	7291
Ratio Analysis	7291
Domain Ratio Analysis	7292
Hadamard Matrix	7292
ODS Table Names	7293
Examples: SURVEYMEANS Procedure	7294
Example 86.1: Stratified Cluster Sample Design	7294
Example 86.2: Domain Analysis	7298
Example 86.3: Ratio Analysis	7302
Example 86.4: Analyzing Survey Data with Missing Values	7302
Example 86.5: Variance Estimation Using Replication Methods	7304
References	7307

Overview: SURVEYMEANS Procedure

The SURVEYMEANS procedure estimates characteristics of a survey population by using statistics computed from a survey sample. You can estimate statistics such as means, totals, proportions, quantiles, and ratios. PROC SURVEYMEANS also provides domain analysis, which computes estimates for subpopulations or domains. The procedure also estimates variances and confidence limits and performs t tests for these statistics. PROC SURVEYMEANS uses either the Taylor series (linearization) method or replication (subsampling) methods to estimate sampling errors of estimators based on complex sample designs. The sample design can be a complex survey sample design with stratification, clustering, and unequal weighting. See Lohr (2009), Särndal, Swensson, and Wretman (1992), and Wolter (1985) for more details.

Getting Started: SURVEYMEANS Procedure

This section demonstrates how you can use the SURVEYMEANS procedure to produce descriptive statistics from sample survey data. For a complete description of PROC SURVEYMEANS, see the section “[Syntax: SURVEYMEANS Procedure](#)” on page 7247. The section “[Examples: SURVEYMEANS Procedure](#)” on page 7294 provides more complicated examples to illustrate the applications of PROC SURVEYMEANS.

Simple Random Sampling

This example illustrates how you can use PROC SURVEYMEANS to estimate population means and proportions from sample survey data. The study population is a junior high school with a total of 4,000 students in grades 7, 8, and 9. Researchers want to know how much these students spend weekly for ice cream, on average, and what percentage of students spend at least \$10 weekly for ice cream.

To answer these questions, 40 students were selected from the entire student population by using simple random sampling (SRS). Selection by simple random sampling means that all students have an equal chance of being selected and no student can be selected more than once. Each student selected for the sample was asked how much he or she spends for ice cream per week, on average. The SAS data set IceCream saves the responses of the 40 students:

```
data IceCream;
  input Grade Spending @@;
  if (Spending < 10) then Group='less';
  else Group='more';
  datalines;
7 7 7 7 8 12 9 10 7 1 7 10 7 3 8 20 8 19 7 2
7 2 9 15 8 16 7 6 7 6 7 6 9 15 8 17 8 14 9 8
9 8 9 7 7 3 7 12 7 4 9 14 8 18 9 9 7 2 7 1
7 4 7 11 9 8 8 10 8 13 7 2 9 6 9 11 7 2 7 9
;
```

The variable Grade contains a student's grade. The variable Spending contains a student's response regarding how much he spends per week for ice cream, in dollars. The variable Group is created to indicate whether a student spends at least \$10 weekly for ice cream: Group='more' if a student spends at least \$10, or Group='less' if a student spends less than \$10.

You can use PROC SURVEYMEANS to produce estimates for the entire student population, based on this random sample of 40 students:

```
title1 'Analysis of Ice Cream Spending';
title2 'Simple Random Sample Design';
proc surveymeans data=IceCream total=4000;
  var Spending Group;
run;
```

The PROC SURVEYMEANS statement invokes the procedure. The TOTAL=4000 option specifies the total number of students in the study population, or school. The procedure uses this total to adjust variance estimates for the effects of sampling from a finite population. The VAR statement names the variables to analyze, Spending and Group.

Figure 86.1 displays the results from this analysis. There are a total of 40 observations used in the analysis. The “Class Level Information” table lists the two levels of the variable Group. This variable is a character variable, and so PROC SURVEYMEANS provides a categorical analysis for it, estimating the relative frequency or proportion for each level. If you want a categorical analysis for a numeric variable, you can name that variable in the CLASS statement.

Figure 86.1 Analysis of Ice Cream Spending

Analysis of Ice Cream Spending						
Simple Random Sample Design						
The SURVEYMEANS Procedure						
Data Summary						
Number of Observations			40			
Class Level Information						
Class Variable		Levels	Values			
Group		2	less more			
Statistics						
Variable	Level	N	Mean	Std Error of Mean	95% CL for Mean	
Spending		40	8.750000	0.845139	7.04054539	10.4594546
Group	less	23	0.575000	0.078761	0.41568994	0.7343101
	more	17	0.425000	0.078761	0.26568994	0.5843101

The “Statistics” table displays the estimates for each analysis variable. By default, PROC SURVEYMEANS displays the number of observations, the estimate of the mean, its standard error, and the 95% confidence limits for the mean. You can obtain other statistics by specifying the corresponding *statistic-keywords* in the PROC SURVEYMEANS statement.

The estimate of the average weekly ice cream expense is \$8.75 for students at this school. The standard error of this estimate is \$0.85, and the 95% confidence interval for weekly ice cream expense is from \$7.04 to \$10.46. The analysis variable Group is a character variable, and so PROC SURVEYMEANS analyzes it as categorical, estimating the relative frequency or proportion for each level or category. These estimates are displayed in the Mean column of the “Statistics” table. It is estimated that 57.5% of all students spend less than \$10 weekly on ice cream, while 42.5% of the students spend at least \$10 weekly. The standard error of each estimate is 7.9%.

Stratified Sampling

Suppose that the sample of students described in the previous section was actually selected by using stratified random sampling. In stratified sampling, the study population is divided into nonoverlapping strata, and samples are selected from each stratum independently.

The list of students in this junior high school was stratified by grade, yielding three strata: grades 7, 8, and 9. A simple random sample of students was selected from each grade. [Table 86.1](#) shows the total number of students in each grade.

Table 86.1 Number of Students by Grade

Grade	Number of Students
7	1,824
8	1,025
9	1,151
Total	4,000

To analyze this stratified sample, you need to provide the population totals for each stratum to PROC SURVEYMEANS. The SAS data set StudentTotals contains the information from [Table 86.1](#):

```
data StudentTotals;
    input Grade _total_;
    datalines;
7 1824
8 1025
9 1151
;
```

The variable Grade is the stratum identification variable, and the variable _TOTAL_ contains the total number of students for each stratum. PROC SURVEYMEANS requires you to use the variable name _TOTAL_ for the stratum population totals.

The procedure uses the stratum population totals to adjust variance estimates for the effects of sampling from a finite population. If you do not provide population totals or sampling rates, then the procedure assumes that the proportion of the population in the sample is very small, and the computation does not involve a finite population correction.

In a stratified sample design, when the sampling rates in the strata are unequal, you need to use sampling weights to reflect this information in order to produce an unbiased mean estimator. In this example, the appropriate sampling weights are reciprocals of the probabilities of selection. You can use the following DATA step to create the sampling weights:

```

data IceCream;
  set IceCream;
  if Grade=7 then Prob=20/1824;
  if Grade=8 then Prob=9/1025;
  if Grade=9 then Prob=11/1151;
  Weight=1/Prob;
run;

```

When you use PROC SURVEYSELECT to select your sample, the procedure creates these sampling weights for you.

The following SAS statements perform the stratified analysis of the survey data:

```

title1 'Analysis of Ice Cream Spending';
title2 'Stratified Sample Design';
proc surveymeans data=IceCream total=StudentTotals;
  stratum Grade / list;
  var Spending Group;
  weight Weight;
run;

```

The PROC SURVEYMEANS statement invokes the procedure. The DATA= option names the SAS data set IceCream as the input data set to be analyzed. The TOTAL= option names the data set StudentTotals as the input data set that contains the stratum population totals. Comparing this to the analysis in the section “[Simple Random Sampling](#)” on page 7241, notice that the TOTAL=StudentTotals option is used here instead of the TOTAL=4000 option. In this stratified sample design, the population totals are different for different strata, and so you need to provide them to PROC SURVEYMEANS in a SAS data set.

The STRATA statement identifies the stratification variable Grade. The LIST option in the STRATA statement requests that the procedure display stratum information. The WEIGHT statement tells the procedure that the variable Weight contains the sampling weights.

[Figure 86.2](#) displays information about the input data set. There are three strata in the design and 40 observations in the sample. The categorical variable Group has two levels, ‘less’ and ‘more.’

[Figure 86.3](#) displays information for each stratum. The table displays a stratum index and the values of the STRATA variable. The stratum index identifies each stratum by a sequentially assigned number. For each stratum, the table gives the population total (total number of students), the sampling rate, and the sample size. The stratum sampling rate is the ratio of the number of students in the sample to the number of students in the population for that stratum. The table also lists each analysis variable and the number of stratum observations for that variable. For categorical variables, the table lists each level and the number of sample observations in that level.

Figure 86.4 shows the following:

- The estimate of average weekly ice cream expense is \$9.14 for students in this school, with a standard error of \$0.53, and a 95% confidence interval from \$8.06 to \$10.22.
- An estimate of 54.5% of all students spend less than \$10 weekly on ice cream, and 45.5% spend more, with a standard error of 5.8%.

Figure 86.4 Analysis of Ice Cream Spending

		Statistics				
Variable	Level	N	Mean	Std Error of Mean	95% CL for Mean	
Spending		40	9.141298	0.531799	8.06377052	10.2188254
Group	less	23	0.544555	0.058424	0.42617678	0.6629323
	more	17	0.455445	0.058424	0.33706769	0.5738232

Output Data Sets

PROC SURVEYMEANS uses the Output Delivery System (ODS) to create output data sets. This is a departure from older SAS procedures that provide OUTPUT statements for similar functionality. For more information about ODS, see Chapter 20, “[Using the Output Delivery System](#).”

For example, to save the “Statistics” table shown in [Figure 86.4](#) in the previous section in an output data set, you use the ODS OUTPUT statement as follows:

```

title1 'Analysis of Ice Cream Spending';
title2 'Stratified Sample Design';
proc surveymeans data=IceCream total=StudentTotals;
    stratum Grade / list;
    var Spending Group;
    weight Weight;
    ods output Statistics=MyStat;
run;

```

The statement

```
ods output Statistics=MyStat;
```

requests that the “Statistics” table that appears in [Figure 86.4](#) be placed in a SAS data set MyStat.

The PRINT procedure displays observations of the data set MyStat:

```

proc print data=MyStat;
run;

```

[Figure 86.5](#) displays the data set MyStat. The section “[ODS Table Names](#)” on page 7293 gives the complete list of tables produced by PROC SURVEYMEANS.

Figure 86.5 Output Data Set MyStat

Analysis of Ice Cream Spending Stratified Sample Design							
Obs	VarName	Var Level	N	Mean	StdErr	Lower CLMean	Upper CLMean
1	Spending		40	9.141298	0.531799	8.06377052	10.2188254
2	Group	less	23	0.544555	0.058424	0.42617678	0.6629323
3	Group	more	17	0.455445	0.058424	0.33706769	0.5738232

Syntax: SURVEYMEANS Procedure

The following statements are available in PROC SURVEYMEANS:

```

PROC SURVEYMEANS < options > < statistic-keywords > ;
  BY variables ;
  CLASS variables ;
  CLUSTER variables ;
  DOMAIN variables < variable*variable variable*variable*variable ... > < / option > ;
  RATIO < 'label' > variables / variables ;
  REPWEIGHTS variables < / options > ;
  STRATA variables < / option > ;
  VAR variables ;
  WEIGHT variable ;

```

The PROC SURVEYMEANS statement invokes the procedure. It optionally names the input data sets, specifies statistics for the procedure to compute, and specifies the variance estimation method. The PROC SURVEYMEANS statement is required.

The VAR statement identifies the variables to be analyzed. The CLASS statement identifies those numeric variables that are to be analyzed as categorical variables. The STRATA statement lists the variables that form the strata in a stratified sample design. The CLUSTER statement specifies cluster identification variables in a clustered sample design. The DOMAIN statement lists the variables that define domains for subpopulation analysis. The RATIO statement requests ratio analysis for means or proportions of analysis variables. The WEIGHT statement names the sampling weight variable. The REPWEIGHTS statement names replicate weight variables for BRR or jackknife variance estimation. You can use a BY statement with PROC SURVEYMEANS to obtain separate analyses for groups defined by the BY variables.

All statements can appear multiple times except the PROC SURVEYMEANS statement and the WEIGHT statement, each of which can appear only once.

The rest of this section gives detailed syntax information for the BY, CLASS, CLUSTER, DOMAIN, RATIO, REPWEIGHTS, STRATA, VAR, and WEIGHT statements in alphabetical order after the description of the PROC SURVEYMEANS statement.

PROC SURVEYMEANS Statement

PROC SURVEYMEANS < options > *statistic-keywords* ;

The PROC SURVEYMEANS statement invokes the procedure. In this statement, you identify the data set to be analyzed, specify the variance estimation method, and provide sample design information. The DATA= option names the input data set to be analyzed. The VARMETHOD= option specifies the variance estimation method, which is the Taylor series method by default. For Taylor series variance estimation, you can include a finite population correction factor in the analysis by providing either the sampling rate or population total with the RATE= or TOTAL= option. If your design is stratified, with different sampling rates or totals for different strata, then you can input these stratum rates or totals in a SAS data set that contains the stratification variables.

In the PROC SURVEYMEANS statement, you also can use *statistic-keywords* to specify statistics, such as population mean and population total, for the procedure to compute. You can also request data set summary information and sample design information.

You can specify the following options in the PROC SURVEYMEANS statement:

ALPHA= α

sets the confidence level for confidence limits. The value of the ALPHA= option must be between 0 and 1, and the default value is 0.05. A confidence level of α produces $100(1 - \alpha)\%$ confidence limits. The default of ALPHA=0.05 produces 95% confidence limits.

DATA=SAS-data-set

specifies the SAS data set to be analyzed by PROC SURVEYMEANS. If you omit the DATA= option, the procedure uses the most recently created SAS data set.

MISSING

treats missing values as a valid (nonmissing) category for all categorical variables, which include CLASS, STRATA, CLUSTER, and DOMAIN variables.

By default, if you do not specify the MISSING option, an observation is excluded from the analysis if it has a missing value. For more information, see the section “[Missing Values](#)” on page 7265.

NOMCAR

requests that the procedure treat missing values in the variance computation as *not missing completely at random* (NOMCAR) for Taylor series variance estimation. When you specify the NOMCAR option, PROC SURVEYMEANS computes variance estimates by analyzing the nonmissing values as a domain (subpopulation), where the entire population includes both nonmissing and missing domains. See the section “[Missing Values](#)” on page 7265 for more details.

By default, PROC SURVEYMEANS completely excludes an observation from analysis if that observation has a missing value, unless you specify the MISSING option for categorical variables. Note that the NOMCAR option has no effect on a categorical variable when you specify the MISSING option, which treats missing values as a valid nonmissing level.

The NOMCAR option applies only to Taylor series variance estimation. The replication methods, which you request with the **VARMETHOD=BRR** and **VARMETHOD=JACKKNIFE** options, do not use the NOMCAR option.

NONSYMCL

requests nonsymmetric confidence limits for quantiles when you request quantiles with **PERCENTILE=** or **QUANTILE=** option. See the section “[Confidence Limits](#)” on page 7280 for more details.

NOSPARSE

suppresses the display of analysis variables with zero frequency. By default, the procedure displays all continuous variables and all levels of categorical variables.

ORDER=DATA | FORMATTED | FREQ | INTERNAL

specifies the order in which the values of the categorical variables are to be reported.

Note that the specified order also determines the ordering for levels of the STRATA, CLUSTER, and DOMAIN variables.

The following shows how PROC SURVEYMEANS interprets values of the ORDER= option:

DATA	orders values according to their order in the input data set.
FORMATTED	orders values by their formatted values. This order is operating environment dependent. By default, the order is ascending.
FREQ	order values by descending frequency count; the most observed value comes first in the order.
INTERNAL	orders values by their unformatted values, which yields the same order that the SORT procedure does. This order is operating environment dependent.

By default, ORDER=FORMATTED.

For more information about sorting order, see the chapter on the SORT procedure in the *Base SAS Procedures Guide* and the discussion of BY-group processing in *SAS Language Reference: Concepts*.

PERCENTILE=(values)

specifies percentiles you want the procedure to compute. You can separate values with blanks or commas. Each value must be between 0 and 100. You can also use the *statistic-keywords* DECILES, MEDIAN, Q1, Q3, and QUANTILES to request common percentiles.

PROC SURVEYMEANS uses Woodruff’s method (Dorfman and Valliant 1993; Särndal, Swensson, and Wretman 1992; and Francisco and Fuller 1991) to estimate the variances of quantiles. BRR and jackknife variance estimation do not apply to the variance estimation for percentiles. See the section “[Quantiles](#)” on page 7278 for more details.

QUANTILE=(values)

specifies quantiles you want the procedure to compute. You can separate values with blanks or commas. Each value must be between 0 and 1. You can also use the *statistic-keywords* DECILES, MEDIAN, Q1, Q3, and QUANTILES to request common quantiles.

PROC SURVEYMEANS uses Woodruff's method (Dorfman and Valliant 1993; Särndal, Swensson, and Wretman 1992; Francisco and Fuller 1991) to estimate the variances of quantiles. BRR and jackknife variance estimation do not apply to the variance estimation for quantiles. See the section “[Quantiles](#)” on page 7278 for more details.

RATE=*value* | *SAS-data-set*

R=*value* | *SAS-data-set*

specifies the sampling rate as a nonnegative *value*, or specifies an input data set that contains the stratum sampling rates. The procedure uses this information to compute a finite population correction for Taylor series variance estimation. The procedure does not use the RATE= option for BRR or jackknife variance estimation, which you request with the [VARMETHOD=BRR](#) or [VARMETHOD=JACKKNIFE](#) option.

If your sample design has multiple stages, you should specify the *first-stage sampling rate*, which is the ratio of the number of PSUs selected to the total number of PSUs in the population.

For a nonstratified sample design, or for a stratified sample design with the same sampling rate in all strata, you should specify a nonnegative *value* for the RATE= option. If your design is stratified with different sampling rates in the strata, then you should name a SAS data set that contains the stratification variables and the sampling rates. See the section “[Specification of Population Totals and Sampling Rates](#)” on page 7266 for more details.

The *value* in the RATE= option or the values of `_RATE_` in the secondary data set must be nonnegative numbers. You can specify *value* as a number between 0 and 1. Or you can specify *value* in percentage form as a number between 1 and 100, and PROC SURVEYMEANS converts that number to a proportion. The procedure treats the value 1 as 100% instead of 1%.

If you do not specify the [TOTAL=](#) or RATE= option, then the Taylor series variance estimation does not include a finite population correction. You cannot specify both the TOTAL= and RATE= options.

STACKING

requests that the procedure produce the output data sets by using a stacking table structure, which was the default before SAS 9. The new default is to produce a rectangular table structure in the output data sets.

A rectangular structure creates one observation for each analysis variable in the data set. A stacking structure creates only one observation in the output data set for all analysis variables.

The STACKING option affects the following tables:

- Domain
- Ratio
- Statistics
- StrataInfo

See the section “[Rectangular and Stacking Structures in an Output Data Set](#)” on page 7285 for more details.

TOTAL=*value* | *SAS-data-set*

N=*value* | *SAS-data-set*

specifies the total number of primary sampling units in the study population as a positive *value*, or specifies an input data set that contains the stratum population totals. The procedure uses this information to compute a finite population correction for Taylor series variance estimation. The procedure does not use the TOTAL= option for BRR or jackknife variance estimation, which you request with the **VARMETHOD=BRR** or **VARMETHOD=JACKKNIFE** option.

For a nonstratified sample design, or for a stratified sample design with the same population total in all strata, you should specify a positive *value* for the TOTAL= option. If your sample design is stratified with different population totals in the strata, then you should name a SAS data set that contains the stratification variables and the population totals. See the section “[Specification of Population Totals and Sampling Rates](#)” on page 7266 for more details.

If you do not specify the TOTAL= or **RATE=** option, then the Taylor series variance estimation does not include a finite population correction. You cannot specify both the TOTAL= and RATE= options.

statistic-keywords

specifies the statistics for the procedure to compute. If you do not specify any *statistic-keywords*, PROC SURVEYMEANS computes the NOBS, MEAN, STDERR, and CLM statistics by default.

The statistics produced depend on the type of the analysis variable. If you name a numeric variable in the CLASS statement, then the procedure analyzes that variable as a categorical variable. The procedure always analyzes character variables as categorical. See the section “[CLASS Statement](#)” on page 7258 for more information.

PROC SURVEYMEANS computes MIN, MAX, and RANGE for numeric variables but not for categorical variables. For numeric variables, the keyword MEAN produces the mean, but for categorical variables it produces the proportion in each category or level. Also, for categorical variables, the keyword NOBS produces the number of observations for each variable level, and the keyword NMISS produces the number of missing observations for each level. If you request the keyword NCLUSTER for a categorical variable, PROC SURVEYMEANS displays for each level the number of clusters with observations in that level. PROC SURVEYMEANS computes SUMWGT in the same way for both categorical and numeric variables, as the sum of the weights over all nonmissing observations.

PROC SURVEYMEANS performs univariate analysis, analyzing each variable separately. Thus the number of nonmissing and missing observations might not be the same for all analysis variables. See the section “[Missing Values](#)” on page 7265 for more information.

The following statistics are available for ratios (which you request with a **RATIO** statement): N, NCLU, SUMWGT, RATIO, STDERR, DF, T, PROBT, and CLM, as shown in the following list. If no statistics are requested, the procedure computes the ratio and its standard error by default.

The valid *statistic-keywords* are as follows:

ALL	all statistics listed
CLM	$100(1 - \alpha)\%$ two-sided confidence limits for the MEAN, where α is determined by the ALPHA= option ; the default is $\alpha = 0.05$
CLSUM	$100(1 - \alpha)\%$ two-sided confidence limits for the SUM, where α is determined by the ALPHA= option ; the default is $\alpha = 0.05$
CV	coefficient of variation for MEAN
CVSUM	coefficient of variation for SUM
DECILES	the 10th, the 20th, . . . , and the 90th percentiles including their standard errors and confidence limits
DF	degrees of freedom for the t test
LCLM	$100(1 - \alpha)\%$ one-sided lower confidence limit of the MEAN, where α is determined by the ALPHA= option ; the default is $\alpha = 0.05$
LCLSUM	$100(1 - \alpha)\%$ one-sided lower confidence limit of the SUM, where α is determined by the ALPHA= option ; the default is $\alpha = 0.05$
MAX	maximum value
MEAN	mean for a numeric variable, or the proportion in each category for a categorical variable
MEDIAN	median for a numeric variable
MIN	minimum value
NCLUSTER	number of clusters
NMISS	number of missing observations
NOBS	number of nonmissing observations
Q1	lower quartile
Q3	upper quartile
QUARTILES	lower quartile (25%th percentile), median (50%th percentile), and upper quartile (75%th percentile), including their standard errors and confidence limits
RANGE	range, MAX–MIN
STD	standard deviation of the SUM. When you request SUM, the procedure computes STD by default.
STDERR	standard error of the MEAN or RATIO. When you request MEAN or RATIO, the procedure computes STDERR by default.
SUM	weighted sum, $\sum w_i y_i$, or estimated population total when the appropriate sampling weights are used
SUMWGT	sum of the weights, $\sum w_i$
T	t -value and its corresponding p -value with DF degrees of freedom for $H_0 : \theta = 0$, where θ is a requested statistic

UCLM	$100(1 - \alpha)\%$ one-sided upper confidence limit of the MEAN, where α is determined by the ALPHA= option; the default is $\alpha = 0.05$
UCLSUM	$100(1 - \alpha)\%$ one-sided upper confidence limit of the SUM, where α is determined by the ALPHA= option; the default is $\alpha = 0.05$
VAR	variance of the MEAN or RATIO
VARSUM	variance of the SUM

See the section “[Statistical Computations](#)” on page 7267 for details about how PROC SURVEYMEANS computes these statistics.

VARMETHOD=BRR <(method-options)>

VARMETHOD=JACKKNIFE | **JK** <(method-options)>

VARMETHOD=TAYLOR

specifies the variance estimation method. **VARMETHOD=TAYLOR** requests the Taylor series method, which is the default if you do not specify the **VARMETHOD=** option or the **REPWEIGHTS** statement. **VARMETHOD=BRR** requests variance estimation by balanced repeated replication (BRR), and **VARMETHOD=JACKKNIFE** requests variance estimation by the delete-1 jackknife method.

For **VARMETHOD=BRR** and **VARMETHOD=JACKKNIFE** you can specify *method-options* in parentheses. [Table 86.2](#) summarizes the available *method-options*.

Table 86.2 Variance Estimation Options

VARMETHOD=	Variance Estimation Method	Method-Options
BRR	Balanced repeated replication	DFADJ FAY <=value> HADAMARD=SAS-data-set OUTWEIGHTS=SAS-data-set PRINTH REPS=number
JACKKNIFE	Jackknife	DFADJ OUTJKCOEFS=SAS-data-set OUTWEIGHTS=SAS-data-set
TAYLOR	Taylor series linearization	None

Method-options must be enclosed in parentheses following the method keyword. For example:

```
varmethod=BRR (reps=60 outweights=myReplicateWeights)
```

The following values are available for the **VARMETHOD=** option:

BRR <(method-options)> requests [balanced repeated replication](#) (BRR) variance estimation. The BRR method requires a stratified sample design with two primary sampling units (PSUs) per stratum. See the section “[Balanced Repeated Replication \(BRR\) Method](#)” on page 7280 for more information.

You can specify the following *method-options* in parentheses following VARMETHOD=BRR:

DFADJ

computes the degrees of freedom as the number of nonmissing strata for an analysis variable. The degrees of freedom for VARMETHOD=BRR equal the number of strata, which by default is based on all valid observations in the data set. But if you specify the DFADJ *method-option*, PROC SURVEYMEANS does not count any empty strata that are due to all observations containing missing values for an analysis variable.

See the section “[Degrees of Freedom](#)” on page 7271 for more information. See the section “[Data and Sample Design Summary](#)” on page 7287 for details about valid observations.

The DFADJ *method-option* has no effect on categorical variables when you specify the [MISSING](#) option, which treats missing values as a valid nonmissing level.

The DFADJ *method-option* cannot be used when you provide replicate weights with a [REPWEIGHTS](#) statement. When you use a REPWEIGHTS statement, the degrees of freedom equal the number of REPWEIGHTS variables (or replicates), unless you specify an alternative value in the DF= option in the REPWEIGHTS statement.

FAY <=value>

requests [Fay’s method](#), a modification of the [BRR](#) method, for variance estimation. See the section “[Fay’s BRR Method](#)” on page 7282 for more information.

You can specify the *value* of the Fay coefficient, which is used in converting the original sampling weights to replicate weights. The Fay coefficient must be a nonnegative number less than 1. By default, the value of the Fay coefficient equals 0.5.

HADAMARD=SAS-data-set

H=SAS-data-set

names a SAS data set that contains the [Hadamard matrix](#) for BRR replicate construction. If you do not provide a Hadamard matrix with the HADAMARD= *method-option*, PROC SURVEYMEANS generates an appropriate Hadamard matrix for replicate construction. See the sections “[Balanced Repeated Replication \(BRR\) Method](#)” on page 7280 and “[Hadamard Matrix](#)” on page 7284 for details.

If a Hadamard matrix of a given dimension exists, it is not necessarily unique. Therefore, if you want to use a specific Hadamard matrix, you must provide the matrix as a SAS data set in the HADAMARD= *method-option*.

In the HADAMARD= input data set, each variable corresponds to a column of the Hadamard matrix, and each observation corresponds to a row of the matrix. You can use any variable names in the HADAMARD= data set. All values in the data set must equal either 1 or -1 . You must ensure that the matrix you provide is indeed a Hadamard matrix—that is, $\mathbf{A}'\mathbf{A} = R\mathbf{I}$, where \mathbf{A} is the Hadamard matrix of dimension R and \mathbf{I} is an identity matrix. PROC SURVEYMEANS does not check the validity of the Hadamard matrix that you provide.

The HADAMARD= input data set must contain at least H variables, where H denotes the number of first-stage strata in your design. If the data set contains more than H variables, the procedure uses only the first H variables. Similarly, the HADAMARD= input data set must contain at least H observations.

If you do not specify the REPS= *method-option*, then the number of replicates is taken to be the number of observations in the HADAMARD= input data set. If you specify the number of replicates—for example, REPS=*nreps*—then the first *nreps* observations in the HADAMARD= data set are used to construct the replicates.

You can specify the PRINTH option to display the Hadamard matrix that the procedure uses to construct replicates for BRR.

OUTWEIGHTS=SAS-data-set

names a SAS data set that contains replicate weights. See the section “[Balanced Repeated Replication \(BRR\) Method](#)” on page 7280 for information about replicate weights. See the section “[Replicate Weights Output Data Set](#)” on page 7284 for more details about the contents of the OUTWEIGHTS= data set.

The OUTWEIGHTS= *method-option* is not available when you provide replicate weights with the [REPWEIGHTS](#) statement.

PRINTH

displays the Hadamard matrix.

When you provide your own Hadamard matrix with the HADAMARD= *method-option*, only the rows and columns of the Hadamard matrix that are used by the procedure are displayed. See the sections “[Balanced Repeated Replication \(BRR\) Method](#)” on page 7280 and “[Hadamard Matrix](#)” on page 7284 for details.

The PRINTH *method-option* is not available when you provide replicate weights with the [REPWEIGHTS](#) statement because the procedure does not use a Hadamard matrix in this case.

REPS=number

specifies the number of replicates for BRR variance estimation. The value of *number* must be an integer greater than 1.

If you do not provide a Hadamard matrix with the **HADAMARD=method-option**, the number of replicates should be greater than the number of strata and should be a multiple of 4. See the section “**Balanced Repeated Replication (BRR) Method**” on page 7280 for more information. If a Hadamard matrix cannot be constructed for the REPS= value that you specify, the value is increased until a Hadamard matrix of that dimension can be constructed. Therefore, it is possible for the actual number of replicates used to be larger than the REPS= value that you specify.

If you provide a Hadamard matrix with the **HADAMARD=method-option**, the value of REPS= must not be less than the number of rows in the Hadamard matrix. If you provide a Hadamard matrix and do not specify the REPS= *method-option*, the number of replicates equals the number of rows in the Hadamard matrix.

If you do not specify the REPS= or **HADAMARD=method-option** and do not include a **REPWEIGHTS** statement, the number of replicates equals the smallest multiple of 4 that is greater than the number of strata.

If you provide replicate weights with the **REPWEIGHTS** statement, the procedure does not use the REPS= *method-option*. With a **REPWEIGHTS** statement, the number of replicates equals the number of **REPWEIGHTS** variables.

JACKKNIFE | JK *<(method-options)>* requests variance estimation by the delete-1 jackknife method. See the section “**Jackknife Method**” on page 7282 for details. If you provide replicate weights with a **REPWEIGHTS** statement, **VARMETHOD=JACKKNIFE** is the default variance estimation method.

You can specify the following *method-options* in parentheses following **VARMETHOD=JACKKNIFE**:

DFADJ

computes the degrees of freedom as the number of nonmissing strata for an analysis variable. The degrees of freedom for **VARMETHOD=JACKKNIFE** equal the number of clusters (or number of observations if there is no clusters) minus the number of strata (or one if there is no strata). By default, the number of strata is based on all valid observations in the data set. But if you specify the **DFADJ** *method-option*, PROC SURVEYMEANS does not count any empty strata that are due to all observations containing missing values for an analysis variable.

See the section “**Degrees of Freedom**” on page 7271 for more information. See the section “**Data and Sample Design Summary**” on page 7287 for details about valid observations.

The DFADJ *method-option* has no effect on categorical variables when you specify the [MISSING](#) option, which treats missing values as a valid nonmissing level.

The DFADJ *method-option* cannot be used when you provide replicate weights with a [REPWEIGHTS](#) statement. When you use a [REPWEIGHTS](#) statement, the degrees of freedom equal the number of [REPWEIGHTS](#) variables (or replicates), unless you specify an alternative value in the DF= option in the [REPWEIGHTS](#) statement.

OUTJKCOEFS=SAS-data-set

names a SAS data set that contains jackknife coefficients. See the section “[Jackknife Method](#)” on page 7282 for information about [jackknife coefficients](#). See the section “[Jackknife Coefficients Output Data Set](#)” on page 7285 for more details about the contents of the OUTJKCOEFS= data set.

OUTWEIGHTS=SAS-data-set

names a SAS data set that contains replicate weights. See the section “[Jackknife Method](#)” on page 7282 for information about replicate weights. See the section “[Replicate Weights Output Data Set](#)” on page 7284 for more details about the contents of the OUTWEIGHTS= data set.

The OUTWEIGHTS= *method-option* is not available when you provide replicate weights with the [REPWEIGHTS](#) statement.

TAYLOR

requests Taylor series variance estimation. This is the default method if you do not specify the VARMETHOD= option or a [REPWEIGHTS](#) statement. See the section “[Taylor Series Method](#)” on page 7270 for more information.

BY Statement

BY variables ;

You can specify a BY statement with PROC SURVEYMEANS to obtain separate analyses on observations in groups that are defined by the BY variables. When a BY statement appears, the procedure expects the input data set to be sorted in order of the BY variables. If you specify more than one BY statement, only the last one specified is used.

If your input data set is not sorted in ascending order, use one of the following alternatives:

- Sort the data by using the SORT procedure with a similar BY statement.
- Specify the NOTSORTED or DESCENDING option in the BY statement for the SURVEYMEANS procedure. The NOTSORTED option does not mean that the data are unsorted but rather that the data are arranged in groups (according to values of the BY variables) and that these groups are not necessarily in alphabetical or increasing numeric order.

- Create an index on the BY variables by using the DATASETS procedure (in Base SAS software).

Note that using a BY statement provides completely separate analyses of the BY groups. It does not provide a statistically valid domain (subpopulation) analysis, where the total number of units in the subpopulation is not known with certainty. You should use the DOMAIN statement to obtain domain analysis. For more information about subpopulation analysis for sample survey data, see Cochran (1977).

For more information about BY-group processing, see the discussion in *SAS Language Reference: Concepts*. For more information about the DATASETS procedure, see the discussion in the *Base SAS Procedures Guide*.

CLASS Statement

CLASS *variables* ;

The CLASS statement names variables to be analyzed as categorical variables. For categorical variables, PROC SURVEYMEANS estimates the proportion in each category or level, instead of the overall mean. PROC SURVEYMEANS always analyzes character variables as categorical. If you want categorical analysis for a numeric variable, you must include that variable in the CLASS statement.

The CLASS *variables* are one or more variables in the DATA= input data set. These variables can be either character or numeric. The formatted values of the CLASS variables determine the categorical variable levels. Thus, you can use formats to group values into levels. See the FORMAT procedure in the *Base SAS Procedures Guide* and the FORMAT statement and SAS formats in *SAS Language Reference: Dictionary* for more information.

When determining levels of a CLASS variable, an observation with missing values for this CLASS variable is excluded, unless you specify the MISSING option. For more information, see the section “Missing Values” on page 7265.

You can use multiple CLASS statements to specify categorical variables.

When you specify classification variables, you can use the SAS system option SUMSIZE= to limit (or to specify) the amount of memory that is available for data analysis. See the chapter on SAS system options in *SAS Language Reference: Dictionary* for a description of the SUMSIZE= option.

CLUSTER Statement

CLUSTER *variables* ;

The CLUSTER statement names variables that identify the clusters in a clustered sample design. The combinations of categories of CLUSTER variables define the clusters in the sample. If there is a STRATA statement, clusters are nested within strata.

If you provide replicate weights for BRR or jackknife variance estimation with the [REPWEIGHTS](#) statement, you do not need to specify a CLUSTER statement.

If your sample design has clustering at multiple stages, you should identify only the first-stage clusters (primary sampling units (PSUs)), in the CLUSTER statement. See the section “[Primary Sampling Units \(PSUs\)](#)” on page 7267 for more information.

The CLUSTER *variables* are one or more variables in the DATA= input data set. These variables can be either character or numeric. The formatted values of the CLUSTER variables determine the CLUSTER variable levels. Thus, you can use formats to group values into levels. See the FORMAT procedure in the *Base SAS Procedures Guide* and the FORMAT statement and SAS formats in *SAS Language Reference: Dictionary* for more information.

When determining levels of a CLUSTER variable, an observation with missing values for this CLUSTER variable is excluded, unless you specify the [MISSING](#) option. For more information, see the section “[Missing Values](#)” on page 7265.

You can use multiple CLUSTER statements to specify cluster variables. The procedure uses variables from all CLUSTER statements to create clusters.

DOMAIN Statement

DOMAIN *variables* < *variable*variable variable*variable*variable ...* > < / *option* > ;

The DOMAIN statement requests analysis for domains (subpopulations) in addition to analysis for the entire study population. The DOMAIN statement names the variables that identify domains, which are called domain variables.

It is common practice to compute statistics for domains. The formation of these domains might be unrelated to the sample design. Therefore, the sample sizes for the domains are random variables. Use a DOMAIN statement to incorporate this variability into the variance estimation.

Note that a DOMAIN statement is different from a [BY](#) statement. In a BY statement, you treat the sample sizes as fixed in each subpopulation, and you perform analysis within each BY group independently. See the section “[Domain Analysis](#)” on page 7267 for more details.

Use the DOMAIN statement on the entire data set to perform a domain analysis. Creating a new data set from a single domain and analyzing that with PROC SURVEYMEANS yields inappropriate estimates of variance.

A domain variable can be either character or numeric. The procedure treats domain variables as categorical variables. If a variable appears by itself in a DOMAIN statement, each level of this variable determines a domain in the study population. If two or more variables are joined by asterisks (*), then every possible combination of levels of these variables determines a domain. The procedure performs a descriptive analysis within each domain that is defined by the domain variables.

When determining levels of a DOMAIN variable, an observation with missing values for this DOMAIN variable is excluded, unless you specify the [MISSING](#) option. For more information, see the section “[Missing Values](#)” on page 7265.

The formatted values of the domain variables determine the categorical variable levels. Thus, you can use formats to group values into levels. See the **FORMAT** procedure in the *Base SAS Procedures Guide* and the **FORMAT** statement and SAS formats in *SAS Language Reference: Dictionary* for more information.

You can specify the following option in the **DOMAIN** statement after a slash (/):

DFADJ

computes the degrees of freedom by using the number of non-empty strata for an analysis variable in a domain.

In a domain analysis, it is possible that some strata contain no sampling units for a specific domain. Or some strata in the domain might be empty due to missing values. By default, the procedure counts these empty strata when computing the degrees of freedom.

However, if you specify the **DFADJ** option, the procedure excludes any empty strata when computing the degrees of freedom. Prior to SAS 9.2, the procedure excluded empty strata by default.

See the section “[Degrees of Freedom](#)” on page 7271 for more information. See the section “[Data and Sample Design Summary](#)” on page 7287 for details about valid observations.

The **DFADJ** option has no effect on categorical variables when you specify the **MISSING** option, which treats missing values as a valid nonmissing level.

RATIO Statement

RATIO < 'label' > *variables* / *variables* ;

The **RATIO** statement requests ratio analysis for means or proportions of analysis variables. A ratio statement names the variables whose means are used as numerators or denominators in a ratio. Variables that appear before the slash (/) are called *numerator variables* and are used as numerators. Variables that appear after the slash (/) are called *denominator variables* and are used as denominators. These *variables* can be any number of analysis variables, either continuous or categorical, except those named in the **BY**, **CLUSTER**, **REPWEIGHTS**, **STRATA**, and **WEIGHT** statements.

You can optionally specify a label for each **RATIO** statement to identify the ratios in the output. Labels must be enclosed in single quotes.

The computation of ratios depends on whether the numerator and denominator variables are continuous or categorical.

For continuous variables, ratios are calculated from the variable means. For example, for continuous variables *X*, *Y*, *Z*, and *T*, the following **RATIO** statement requests that the procedure analyze the ratios \bar{x}/\bar{z} , \bar{x}/\bar{t} , \bar{y}/\bar{z} , and \bar{y}/\bar{t} :

```
ratio x y / z t;
```

If a continuous variable appears as both a numerator and a denominator variable, the ratio of this variable to itself is ignored.

For categorical variables, ratios are calculated with the proportions for the categories. For example, if the categorical variable Gender has the values ‘Male’ and ‘Female,’ with the proportions $p_m = \text{Pr}(\text{Gender}=\text{‘Male’})$ and $p_f = \text{Pr}(\text{Gender}=\text{‘Female’})$, and Y is a continuous variable, then the following RATIO statement requests that the procedure analyze the ratios p_m/p_f , p_f/p_m , \bar{y}/p_m , and \bar{y}/p_f :

```
ratio Gender y / Gender;
```

If a categorical variable appears as both a numerator and denominator variable, then the ratios of the proportions for all categories are computed, except the ratio of each category to itself.

You can have more than one RATIO statement. Each RATIO statement produces ratios independently by using its own numerator and denominator variables. Each RATIO statement also produces its own ratio analysis table.

Available statistics for a ratio are as follows:

- N, number of observations used to compute the ratio
- NCLU, number of clusters
- SUMWGT, sum of weights
- RATIO, ratio
- STDERR, standard error of ratio
- VAR, variance of ratio
- T, *t*-value of ratio
- PROBT, *p*-value of *t*
- DF, degrees of freedom of *t*
- CLM, two-sided confidence limits of ratio
- UCLM, one-sided upper confidence limit of ratio
- LCLM, one-sided lower confidence limit of ratio

The procedure calculates these statistics based on the *statistic-keywords* that you specify in the PROC SURVEYMEANS statement. If a *statistic-keyword* is not appropriate for a RATIO statement, that *statistic-keyword* is ignored for the ratios. If no valid statistics are requested for a RATIO statement, the procedure computes the ratio and its standard error by default.

When the means or proportions for the numerator and denominator variables in a ratio are calculated, an observation is excluded if it has a missing value for a continuous numerator or denominator variable. The procedure also excludes an observation with a missing value for a categorical numerator or denominator variable unless you specify the **MISSING** option.

When the denominator for a ratio is zero, then the value of the ratio is displayed as ‘–Infy’, ‘Infy’, or a missing value, depending on whether the numerator is negative, positive, or zero, respectively, and the corresponding internal value is the special missing value ‘.M’, the special missing value ‘.I’, or the usual missing value, respectively.

REPWEIGHTS Statement

REPWEIGHTS *variables* < / *options* > ;

The REPWEIGHTS statement names variables that provide replicate weights for BRR or jackknife variance estimation, which you request with the **VARMETHOD=BRR** or **VARMETHOD=JACKKNIFE** option in the PROC SURVEYMEANS statement. If you do not provide replicate weights for these methods by using a REPWEIGHTS statement, then the procedure constructs replicate weights for the analysis. See the sections “[Balanced Repeated Replication \(BRR\) Method](#)” on page 7280 and “[Jackknife Method](#)” on page 7282 for information about replicate weights.

Each REPWEIGHTS variable should contain the weights for a single replicate, and the number of replicates equals the number of REPWEIGHTS variables. The REPWEIGHTS variables must be numeric, and the variable values must be nonnegative numbers.

If you provide replicate weights with a REPWEIGHTS statement, you do not need to specify a **CLUSTER** or **STRATA** statement. If you use a REPWEIGHTS statement and do not specify the **VARMETHOD=** option in the PROC SURVEYMEANS statement, the procedure uses **VARMETHOD=JACKKNIFE** by default.

If you specify a REPWEIGHTS statement but do not include a **WEIGHT** statement, the procedure uses the average of replicate weights of each observation as the observation’s weight.

You can specify the following options in the REPWEIGHTS statement after a slash (/):

DF=*df*

specifies the degrees of freedom for the analysis. The value of *df* must be a positive number. By default, the degrees of freedom equals the number of REPWEIGHTS variables.

JKCOEFS=*value*

specifies a [jackknife coefficient](#) for **VARMETHOD=JACKKNIFE**. The coefficient *value* must be a nonnegative number. See the section “[Jackknife Method](#)” on page 7282 for details about jackknife coefficients.

You can use this option to specify a single value of the jackknife coefficient, which the procedure uses for all replicates. To specify different coefficients for different replicates, use the **JKCOEFS=values** or **JKCOEFS=SAS-data-set** option.

JKCOEFS=*values*

specifies jackknife coefficients for **VARMETHOD=JACKKNIFE**, where each coefficient corresponds to an individual replicate that is identified by a REPWEIGHTS variable. You can separate *values* with blanks or commas. The coefficient *values* must be nonnegative numbers. The number of *values* must equal the number of replicate weight variables named in the REPWEIGHTS statement. List these values in the same order in which you list the corresponding replicate weight variables in the REPWEIGHTS statement.

See the section “[Jackknife Method](#)” on page 7282 for details about jackknife coefficients.

To specify different coefficients for different replicates, you can also use the **JKCOEFS=SAS-data-set** option. To specify a single jackknife coefficient for all replicates, use the **JKCOEFS=value** option.

JKCOEFS=SAS-data-set

names a SAS data set that contains the jackknife coefficients for **VARMETHOD=JACKKNIFE**. You provide the jackknife coefficients in the JKCOEFS= data set variable JKCoefficient. Each coefficient value must be a nonnegative number. The observations in the JKCOEFS= data set should correspond to the replicates that are identified by the REPWEIGHTS variables. Arrange the coefficients or observations in the JKCOEFS= data set in the same order in which you list the corresponding replicate weight variables in the REPWEIGHTS statement. The number of observations in the JKCOEFS= data set must not be less than the number of REPWEIGHTS variables.

See the section “[Jackknife Method](#)” on page 7282 for details about jackknife coefficients.

To specify different coefficients for different replicates, you can also use the **JKCOEFS=values** option. To specify a single jackknife coefficient for all replicates, use the **JKCOEFS=value** option.

STRATA Statement

STRATA *variables* < / *option* > ;

The STRATA statement specifies variables that form the strata in a stratified sample design. The combinations of categories of STRATA variables define the strata in the sample.

If your sample design has stratification at multiple stages, you should identify only the first-stage strata in the STRATA statement. See the section “[Specification of Population Totals and Sampling Rates](#)” on page 7266 for more information.

If you provide replicate weights for BRR or jackknife variance estimation with the **REPWEIGHTS** statement, you do not need to specify a STRATA statement.

The STRATA *variables* are one or more variables in the DATA= input data set. These variables can be either character or numeric. The formatted values of the STRATA variables determine the levels. Thus, you can use formats to group values into levels. See the FORMAT procedure in the *Base SAS Procedures Guide* and the FORMAT statement and SAS formats in *SAS Language Reference: Dictionary* for more information.

When determining levels of a STRATA variable, an observation with missing values for this STRATA variable is excluded, unless you specify the **MISSING** option. For more information, see the section “[Missing Values](#)” on page 7265.

You can use multiple STRATA statements to specify stratum variables.

You can specify the following option in the STRATA statement after a slash (/):

LIST

displays a “Stratum Information” table, which includes values of the STRATA variables and the number of observations, number of clusters, population total, and sampling rate for each stratum. See the section “[Stratum Information](#)” on page 7288 for more details.

VAR Statement

VAR *variables* ;

The VAR statement names the variables to be analyzed.

If you want a categorical analysis for a numeric variable, you must also name that variable in the **CLASS** statement. For categorical variables, PROC SURVEYMEANS estimates the proportion in each category or level, instead of the overall mean. Character variables are always analyzed as categorical variables. See the section “**CLASS Statement**” on page 7258 for more information.

When you specify a variable in a **RATIO** statement, but not in a VAR statement, the procedure includes this variable as an analysis variable.

If you do not specify a VAR statement, then PROC SURVEYMEANS analyzes all variables in the DATA= input data set, except those named in the **BY**, **CLUSTER**, **DOMAIN**, **REPWEIGHTS**, **STRATA**, and **WEIGHT** statements.

WEIGHT Statement

WEIGHT *variable* ;

The WEIGHT statement names the variable that contains the sampling weights. This variable must be numeric, and the sampling weights must be positive numbers. If an observation has a weight that is nonpositive or missing, then the procedure omits that observation from the analysis. See the section “**Missing Values**” on page 7265 for more information. If you specify more than one WEIGHT statement, the procedure uses only the first WEIGHT statement and ignores the rest.

If you do not specify a WEIGHT statement but provide replicate weights with a **REPWEIGHTS** statement, PROC SURVEYMEANS uses the average of replicate weights of each observation as the observation’s weight.

If you do not specify a WEIGHT statement or a REPWEIGHTS statement, PROC SURVEYMEANS assigns all observations a weight of one.

Details: SURVEYMEANS Procedure

Missing Values

If you have missing values in your survey data for any reason, such as nonresponse, this can compromise the quality of your survey results. If the respondents are different from the nonrespondents with regard to a survey effect or outcome, then survey estimates might be biased and cannot accurately represent the survey population. There are a variety of techniques in sample design and survey operations that can reduce nonresponse. After data collection is complete, you can use imputation to replace missing values with acceptable values, and/or you can use sampling weight adjustments to compensate for nonresponse. You should complete this data preparation and adjustment before you analyze your data with PROC SURVEYMEANS. See Cochran (1977); Kalton and Kaspyzyk (1986); and Brick and Kalton (1996) for more information.

If an observation has a missing value or a nonpositive value for the **WEIGHT** variable, then that observation is excluded from the analysis.

An observation is also excluded from the analysis if it has a missing value for any design (**STRATA**, **CLUSTER**, or **DOMAIN**) variable, unless you specify the **MISSING** option in the PROC SURVEYMEANS statement. If you specify the **MISSING** option, the procedure treats missing values as a valid (nonmissing) category for all categorical variables.

By default, when computing statistics for an analysis variable, PROC SURVEYMEANS omits observations with missing values for that analysis variable. The procedure computes statistics for each variable based only on observations that have nonmissing values for that variable. This treatment is based on the assumption that the missing values are missing completely at random (MCAR). However, this assumption is sometimes not true. For example, evidence from other surveys might suggest that observations with missing values are systematically different from observations without missing values. If you believe that missing values are not missing completely at random, then you can specify the **NOMCAR** option to let variance estimation include these observations with missing values in the analysis variables.

Whether or not you specify the **NOMCAR** option, the procedure always excludes observations with missing or invalid values for the **WEIGHT**, **STRATA**, **CLUSTER**, and **DOMAIN** variables, unless you specify the **MISSING** option.

When you specify the **NOMCAR** option, the procedure treats observations with and without missing values for analysis variables as two different domains, and it performs a domain analysis in the domain of nonmissing observations.

The procedure performs univariate analysis and analyzes each VAR variable separately. Thus, the number of missing observations might be different for different variables. You can specify the keyword **NMISS** in the PROC SURVEYMEANS statement to display the number of missing values for each analysis variable in the “Statistics” table.

When you specify a **RATIO** statement, the procedure excludes any observation that has a missing value for a continuous numerator or denominator variable. The procedure also excludes an observation with a missing value for a categorical numerator or denominator variable unless you specify the **MISSING** option.

If you use a **REPWEIGHTS** statement, all **REPWEIGHTS** variables must contain nonmissing values.

Survey Data Analysis

Specification of Population Totals and Sampling Rates

To include a finite population correction (*fpc*) in Taylor series variance estimation, you can input either the sampling rate or the population total by using the **RATE=** or **TOTAL=** option in the PROC SURVEYMEANS statement. (You cannot specify both of these options in the same PROC SURVEYMEANS statement.) The **RATE=** and **TOTAL=** options apply only to Taylor series variance estimation. The procedure does not use a finite population correction for BRR or jackknife variance estimation.

If you do not specify the **RATE=** or **TOTAL=** option, the Taylor series variance estimation does not include a finite population correction. For fairly small sampling fractions, it is appropriate to ignore this correction. See Cochran (1977) and Kish (1965) for more information.

If your design has multiple stages of selection and you are specifying the **RATE=** option, you should input the first-stage sampling rate, which is the ratio of the number of PSUs in the sample to the total number of PSUs in the study population. If you are specifying the **TOTAL=** option for a multistage design, you should input the total number of PSUs in the study population. See the section “[Primary Sampling Units \(PSUs\)](#)” on page 7267 for more details.

For a nonstratified sample design, or for a stratified sample design with the same sampling rate or the same population total in all strata, you can use the **RATE=value** or **TOTAL=value** option. If your sample design is stratified with different sampling rates or population totals in different strata, use the **RATE=SAS-data-set** or **TOTAL=SAS-data-set** option to name a SAS data set that contains the stratum sampling rates or totals. This data set is called a *secondary data set*, as opposed to the *primary data set* that you specify with the **DATA=** option.

The secondary data set must contain all the stratification variables listed in the **STRATA** statement and all the variables in the **BY** statement. If there are formats associated with the **STRATA** variables and the **BY** variables, then the formats must be consistent in the primary and the secondary data sets. If you specify the **TOTAL=SAS-data-set** option, the secondary data set must have a variable named **_TOTAL_** that contains the stratum population totals. Or if you specify the **RATE=SAS-data-set** option, the secondary data set must have a variable named **_RATE_** that contains the stratum sampling rates. If the secondary data set contains more than one observation for any one stratum, then the procedure uses the first value of **_TOTAL_** or **_RATE_** for that stratum and ignores the rest.

The *value* in the **RATE=** option or the values of **_RATE_** in the secondary data set must be nonnegative numbers. You can specify *value* as a number between 0 and 1. Or you can specify *value* in percentage form as a number between 1 and 100, and PROC SURVEYMEANS converts that number to a proportion. The procedure treats the value 1 as 100% instead of 1%.

If you specify the `TOTAL=value` option, *value* must not be less than the sample size. If you provide stratum population totals in a secondary data set, these values must not be less than the corresponding stratum sample sizes.

Primary Sampling Units (PSUs)

When you have clusters, or primary sampling units (PSUs), in your sample design, the procedure estimates variance from the variation among PSUs when the Taylor series variance method is used. See the section “[Variance and Standard Error of the Mean](#)” on page 7270 and the section “[Variance and Standard Deviation of the Total](#)” on page 7273 for more information.

BRR or jackknife variance estimation methods draw multiple replicates (or subsamples) from the full sample by following a specific resampling scheme. These subsamples are constructed by deleting PSUs from the full sample.

If you use a `REPWEIGHTS` statement to provide replicate weights for BRR or jackknife variance estimation, you do not need to specify a `CLUSTER` statement. Otherwise, you should specify a `CLUSTER` statement whenever your design includes clustering at the first stage of sampling. If you do not specify a `CLUSTER` statement, then PROC SURVEYMEANS treats each observation as a PSU.

Domain Analysis

It is common practice to compute statistics for domains (subpopulations), in addition to computing statistics for the entire study population. Analysis for domains that uses the entire sample is called *domain analysis* (also called subgroup analysis, subpopulation analysis, or subdomain analysis). The formation of these subpopulations of interest might be unrelated to the sample design. Therefore, the sample sizes for the subpopulations might actually be random variables.

Use a `DOMAIN` statement to incorporate this variability into the variance estimation. Note that using a `BY` statement provides completely separate analyses of the BY groups. It does not provide a statistically valid subpopulation or domain analysis, where the total number of units in the subpopulation is not known with certainty.

For more detailed information about domain analysis, see Kish (1965).

Statistical Computations

The SURVEYMEANS procedure uses the Taylor series (linearization) method or replication (re-sampling) methods to estimate sampling errors of estimators based on complex sample designs. For details see Wolter (1985); Lohr (2009); Kalton (1983); Hidiroglou, Fuller, and Hickman (1980); Fuller et al. (1989); Lee, Forthoffer, and Lorimor (1989); Cochran (1977); Kish (1965); Hansen, Hurwitz, and Madow (1953); Rust (1985); Dippo, Fay, and Morganstein (1984); Rao and Shao (1999); Rao, Wu, and Yue (1992); and Rao and Shao (1996). You can use the `VARMETHOD=` option to specify a variance estimation method to use. By default, the Taylor series method is used.

The Taylor series method obtains a linear approximation for the estimator and then uses the variance estimate for this approximation to estimate the variance of the estimate itself (Woodruff 1971, Fuller 1975). When there are clusters, or PSUs, in the sample design, the procedure estimates variance from the variation among PSUs. When the design is stratified, the procedure pools stratum variance estimates to compute the overall variance estimate. For t tests of the estimates, the degrees of freedom equal the number of clusters minus the number of strata in the sample design.

For a multistage sample design, the Taylor series estimation depends only on the first stage of the sample design. Therefore, the required input includes only first-stage cluster (PSU) and first-stage stratum identification. You do not need to input design information about any additional stages of sampling. This variance estimation method assumes that the first-stage sampling fraction is small, or that the first-stage sample is drawn with replacement, as it often is in practice.

Quite often in complex surveys, respondents have unequal weights, which reflect unequal selection probabilities and adjustments for nonresponse. In such surveys, the appropriate sampling weights must be used to obtain valid estimates for the study population.

However, replication methods have recently gained popularity for estimating variances in complex survey data analysis. One reason for this popularity is the relative simplicity of replication-based estimates, especially for nonlinear estimators; another is that modern computational capacity has made replication methods feasible for practical survey analysis.

Replication methods draw multiple replicates (also called subsamples) from a full sample according to a specific resampling scheme. The most commonly used resampling schemes are the *balanced repeated replication* (BRR) method and the *jackknife* method. For each replicate, the original weights are modified for the PSUs in the replicates to create replicate weights. The population parameters of interest are estimated by using the replicate weights for each replicate. Then the variances of parameters of interest are estimated by the variability among the estimates derived from these replicates. You can use a **REPWEIGHTS** statement to provide your own replicate weights for variance estimation. For more information about using replication methods to analyze sample survey data, see the section “[Replication Methods for Variance Estimation](#)” on page 7280.

Definitions and Notation

For a stratified clustered sample design, together with the sampling weights, the sample can be represented by an $n \times (P + 1)$ matrix

$$\begin{aligned} (\mathbf{w}, \mathbf{Y}) &= (w_{hij}, \mathbf{y}_{hij}) \\ &= (w_{hij}, y_{hij}^{(1)}, y_{hij}^{(2)}, \dots, y_{hij}^{(P)}) \end{aligned}$$

where

- $h = 1, 2, \dots, H$ is the stratum index
- $i = 1, 2, \dots, n_h$ is the cluster index within stratum h
- $j = 1, 2, \dots, m_{hi}$ is the unit index within cluster i of stratum h

- $p = 1, 2, \dots, P$ is the analysis variable number, with a total of P variables
- $n = \sum_{h=1}^H \sum_{i=1}^{n_h} m_{hi}$ is the total number of observations in the sample
- w_{hij} denotes the sampling weight for unit j in cluster i of stratum h
- $y_{hij} = (y_{hij}^{(1)}, y_{hij}^{(2)}, \dots, y_{hij}^{(P)})$ are the observed values of the analysis variables for unit j in cluster i of stratum h , including both the values of numerical variables and the values of indicator variables for levels of categorical variables.

For a categorical variable C , let l denote the number of levels of C , and denote the level values as c_1, c_2, \dots, c_l . Let $y^{(q)}$ ($q \in \{1, 2, \dots, P\}$) be an indicator variable for the category $C = c_k$ ($k = 1, 2, \dots, l$) with the observed value in unit j in cluster i of stratum h :

$$y_{hij}^{(q)} = I_{\{C=c_k\}}(h, i, j) = \begin{cases} 1 & \text{if } C_{hij} = c_k \\ 0 & \text{otherwise} \end{cases}$$

Note that the indicator variable $y_{hij}^{(q)}$ is set to missing when C_{hij} is missing. Therefore, the total number of analysis variables, P , is the total number of numerical variables plus the total number of levels of all categorical variables.

The sampling rate f_h for stratum h , which is used in Taylor series variance estimation, is the fraction of first-stage units (PSUs) selected for the sample. You can use the TOTAL= or RATE= option to input population totals or sampling rates. See the section “[Specification of Population Totals and Sampling Rates](#)” on page 7266 for details. If you input stratum totals, PROC SURVEYMEANS computes f_h as the ratio of the stratum sample size to the stratum total. If you input stratum sampling rates, PROC SURVEYMEANS uses these values directly for f_h . If you do not specify the TOTAL= or RATE= option, then the procedure assumes that the stratum sampling rates f_h are negligible, and a finite population correction is not used when computing variances. Replication methods specified by the [VARMETHOD=BRR](#) or the [VARMETHOD=JACKKNIFE](#) option do not use this finite population correction f_h .

Mean

When you specify the keyword MEAN, the procedure computes the estimate of the mean (mean per element) from the survey data. Also, the procedure computes the mean by default if you do not specify any *statistic-keywords* in the PROC SURVEYMEANS statement.

PROC SURVEYMEANS computes the estimate of the mean as

$$\hat{Y} = \left(\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} w_{hij} y_{hij} \right) / w_{...}$$

where

$$w_{...} = \sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} w_{hij}$$

is the sum of the weights over all observations in the sample.

Variance and Standard Error of the Mean

When you specify the keyword `STDERR`, the procedure computes the standard error of the mean. Also, the procedure computes the standard error by default if you specify the keyword `MEAN`, or if you do not specify any *statistic-keywords* in the `PROC SURVEYMEANS` statement. The keyword `VAR` requests the variance of the mean.

Taylor Series Method

When you use `VARMETHOD=TAYLOR`, or by default if you do not specify the `VARMETHOD=` option, `PROC SURVEYMEANS` uses the Taylor series method to estimate the variance of the mean $\hat{\bar{Y}}$. The procedure computes the estimated variance as

$$\widehat{V}(\hat{\bar{Y}}) = \sum_{h=1}^H \widehat{V}_h(\hat{\bar{Y}})$$

where if $n_h > 1$,

$$\begin{aligned} \widehat{V}_h(\hat{\bar{Y}}) &= \frac{n_h(1-f_h)}{n_h-1} \sum_{i=1}^{n_h} (e_{hi\cdot} - \bar{e}_{h\cdot\cdot})^2 \\ e_{hi\cdot} &= \left(\sum_{j=1}^{m_{hi}} w_{hij} (y_{hij} - \hat{\bar{Y}}) \right) / w_{hi\cdot} \\ \bar{e}_{h\cdot\cdot} &= \left(\sum_{i=1}^{n_h} e_{hi\cdot} \right) / n_h \end{aligned}$$

and if $n_h = 1$,

$$\widehat{V}_h(\hat{\bar{Y}}) = \begin{cases} \text{missing} & \text{if } n_{h'} = 1 \text{ for } h' = 1, 2, \dots, H \\ 0 & \text{if } n_{h'} > 1 \text{ for some } 1 \leq h' \leq H \end{cases}$$

Replication Methods

When you specify `VARMETHOD=BRR` or `VARMETHOD=JACKKNIFE`, the procedure computes the variance $\widehat{V}(\hat{\bar{Y}})$ with replication methods by using the variability among replicate estimates to estimate the overall variance. See the section “[Replication Methods for Variance Estimation](#)” on page 7280 for more details.

Standard Error

The standard error of the mean is the square root of the estimated variance.

$$\text{StdErr}(\hat{\bar{Y}}) = \sqrt{\widehat{V}(\hat{\bar{Y}})}$$

***t* Test for the Mean**

If you specify the keyword **T**, PROC SURVEYMEANS computes the *t*-value for testing that the population mean equals zero, $H_0 : \bar{Y} = 0$. The test statistic equals

$$t(\hat{Y}) = \hat{Y} / \text{StdErr}(\hat{Y})$$

The two-sided *p*-value for this test is

$$\text{Prob}(|T| > |t(\hat{Y})|)$$

where *T* is a random variable with the *t* distribution with *df* degrees of freedom.

Degrees of Freedom

PROC SURVEYMEANS computes degrees of freedom *df* to obtain the 100(1 – α)% confidence limits for means, proportions, totals, ratios, and other statistics. The degrees of freedom computation depends on the variance estimation method that you request. Missing values can affect the degrees of freedom computation. See the section “[Missing Values](#)” on page 7265 for details.

Taylor Series Variance Estimation

For the Taylor series method, PROC SURVEYMEANS calculates the degrees of freedom for the *t* test as the number of clusters minus the number of strata. If there are no clusters, then the degrees of freedom equal the number of observations minus the number of strata. If the design is not stratified, then the degrees of freedom equal the number of PSUs minus one.

If all observations in a stratum are excluded from the analysis due to missing values, then that stratum is called an *empty stratum*. Empty strata are not counted in the total number of strata for the table. Similarly, empty clusters and missing observations are not included in the total counts of cluster and observations that are used to compute the degrees of freedom for the analysis.

If you specify the **MISSING** option, missing values are treated as valid nonmissing levels for a categorical variable and are included in computing degrees of freedom. If you specify the **NOMCAR** option for Taylor series variance estimation, observations with missing values for an analysis variable are included in computing degrees of freedom.

Replicate-Based Variance Estimation

When there is a **REPWEIGHTS** statement, the degrees of freedom equal the number of **REPWEIGHTS** variables, unless you specify an alternative in the **DF=** option in a **REPWEIGHTS** statement.

For BRR or jackknife variance estimation without a **REPWEIGHT** statement, by default PROC SURVEYMEANS computes the degrees of freedom by using all valid observations in the input data set. A valid observation is an observation that has a positive value of the **WEIGHT** variable and nonmissing values of the **STRATA** and **CLUSTER** variables unless you specify the **MISSING**

option. See the section “Data and Sample Design Summary” on page 7287 for details about valid observations.

For BRR variance estimation (including [Fay’s method](#)) without a REPWEIGHTS statement, PROC SURVEYMEANS calculates the degrees of freedom as the number of strata. PROC SURVEYMEANS bases the number of strata on all valid observations in the data set, unless you specify the [DFADJ method-option](#) for **VARMETHOD=BRR**. When you specify the DFADJ option, the procedure computes the degrees of freedom as the number of nonmissing strata for an analysis variable. This excludes any empty strata that occur when observations with missing values of that analysis variable are removed.

For jackknife variance estimation without a REPWEIGHTS statement, PROC SURVEYMEANS calculates the degrees of freedom as the number of clusters (or number of observations if there are no clusters) minus the number of strata (or one if there are no strata). For jackknife variance estimation, PROC SURVEYMEANS bases the number of strata and clusters on all valid observations in the data set, unless you specify the [DFADJ method-option](#) for **VARMETHOD=JACKKNIFE**. When you specify the DFADJ option, the procedure computes the degrees of freedom from the number of nonmissing strata and clusters for an analysis variable. This excludes any empty strata or clusters that occur when observations with missing values of an analysis variable are removed.

The procedure displays the degrees of freedom for the t test if you specify the keyword DF in the PROC SURVEYMEANS statement.

Confidence Limits for the Mean

If you specify the keyword CLM, the procedure computes two-sided confidence limits for the mean. Also, the procedure includes the confidence limits by default if you do not specify any [statistic-keywords](#) in the PROC SURVEYMEANS statement.

The confidence coefficient is determined by the value of the ALPHA= option, which by default equals 0.05 and produces 95% confidence limits. The confidence limits are computed as

$$\hat{Y} \pm \text{StdErr}(\hat{Y}) \cdot t_{df, \alpha/2}$$

where \hat{Y} is the estimate of the mean, $\text{StdErr}(\hat{Y})$ is the standard error of the mean, and $t_{df, \alpha/2}$ is the $100(1 - \alpha/2)$ th percentile of the t distribution with df calculated as described in the section “[t Test for the Mean](#)” on page 7271.

If you specify the keyword UCLM, the procedure computes the one-sided upper $100(1 - \alpha)\%$ confidence limit for the mean:

$$\hat{Y} + \text{StdErr}(\hat{Y}) \cdot t_{df, \alpha}$$

If you specify the keyword LCLM, the procedure computes the one-sided lower $100(1 - \alpha)\%$ confidence limit for the mean:

$$\hat{Y} - \text{StdErr}(\hat{Y}) \cdot t_{df, \alpha}$$

Coefficient of Variation

If you specify the keyword CV, PROC SURVEYMEANS computes the coefficient of variation, which is the ratio of the standard error of the mean to the estimated mean:

$$cv(\bar{Y}) = \text{StdErr}(\hat{\bar{Y}}) / \hat{\bar{Y}}$$

If you specify the keyword CVSUM, PROC SURVEYMEANS computes the coefficient of variation for the estimated total, which is the ratio of the standard deviation of the sum to the estimated total:

$$cv(Y) = \text{Std}(\hat{Y}) / \hat{Y}$$

Proportions

If you specify the keyword MEAN for a categorical variable, PROC SURVEYMEANS estimates the proportion, or relative frequency, for each level of the categorical variable. If you do not specify any *statistic-keywords* in the PROC SURVEYMEANS statement, the procedure estimates the proportions for levels of the categorical variables, together with their standard errors and confidence limits.

The procedure estimates the proportion in level c_k for variable C as

$$\hat{p} = \frac{\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} w_{hij} y_{hij}^{(q)}}{\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} w_{hij}}$$

where $y_{hij}^{(q)}$ is the value of the indicator function for level $C = c_k$, defined in the section “[Definitions and Notation](#)” on page 7268, and $y_{hij}^{(q)}$ equals 1 if the observed value of variable C equals c_k , and $y_{hij}^{(q)}$ equals 0 otherwise. Since the proportion estimator is actually an estimator of the mean for an indicator variable, the procedure computes its variance and standard error according to the method outlined in the section “[Variance and Standard Error of the Mean](#)” on page 7270. Similarly, the procedure computes confidence limits for proportions as described in the section “[Confidence Limits for the Mean](#)” on page 7272.

Total

If you specify the keyword SUM, the procedure computes the estimate of the population total from the survey data. The estimate of the total is the weighted sum over the sample:

$$\hat{Y} = \sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} w_{hij} y_{hij}$$

For a categorical variable level, \hat{Y} estimates its total frequency in the population.

Variance and Standard Deviation of the Total

When you specify the keyword STD or the keyword SUM, the procedure estimates the standard deviation of the total. The keyword VARSUM requests the variance of the total.

Taylor Series Method

When you use **VARMETHOD=TAYLOR**, or by default, PROC SURVEYMEANS uses the Taylor series method to estimate the variance of the total as

$$\widehat{V}(\widehat{Y}) = \sum_{h=1}^H \widehat{V}_h(\widehat{Y})$$

where if $n_h > 1$,

$$\widehat{V}_h(\widehat{Y}) = \frac{n_h(1-f_h)}{n_h-1} \sum_{i=1}^{n_h} (y_{hi\cdot} - \bar{y}_{h\cdot\cdot})^2$$

$$y_{hi\cdot} = \sum_{j=1}^{m_{hi}} w_{hij} y_{hij}$$

$$\bar{y}_{h\cdot\cdot} = \left(\sum_{i=1}^{n_h} y_{hi\cdot} \right) / n_h$$

and if $n_h = 1$,

$$\widehat{V}_h(\widehat{Y}) = \begin{cases} \text{missing} & \text{if } n_{h'} = 1 \text{ for } h' = 1, 2, \dots, H \\ 0 & \text{if } n_{h'} > 1 \text{ for some } 1 \leq h' \leq H \end{cases}$$

Replication Methods

When you specify **VARMETHOD=BRR** or **VARMETHOD=JACKKNIFE** option, the procedure computes the variance $\widehat{V}(\widehat{Y})$ with replication methods by measuring the variability among the estimates derived from these replicates. See the section “[Replication Methods for Variance Estimation](#)” on page 7280 for more details.

Standard Deviation

The standard deviation of the total equals

$$\text{Std}(\widehat{Y}) = \sqrt{\widehat{V}(\widehat{Y})}$$

Confidence Limits for the Total

If you specify the keyword **CLSUM**, the procedure computes confidence limits for the total. The confidence coefficient is determined by the value of the **ALPHA=** option, which by default equals 0.05 and produces 95% confidence limits. The confidence limits are computed as

$$\widehat{Y} \pm \text{Std}(\widehat{Y}) \cdot t_{df, \alpha/2}$$

where \hat{Y} is the estimate of the total, $\text{Std}(\hat{Y})$ is the estimated standard deviation, and $t_{df, \alpha/2}$ is the $100(1 - \alpha/2)$ th percentile of the t distribution with df calculated as described in the section “[t Test for the Mean](#)” on page 7271.

If you specify the keyword UCLSUM, the procedure computes the one-sided upper $100(1 - \alpha)\%$ confidence limit for the sum:

$$\hat{Y} + \text{Std}(\hat{Y}) \cdot t_{df, \alpha}$$

If you specify the keyword LCLSUM, the procedure computes the one-sided lower $100(1 - \alpha)\%$ confidence limit for the sum:

$$\hat{Y} - \text{Std}(\hat{Y}) \cdot t_{df, \alpha}$$

Ratio

When you use a **RATIO** statement, the procedure produces statistics requested by the *statistic-keywords* in the PROC SURVEYMEANS statement.

Suppose that you want to calculate the ratio of variable Y to variable X . Let x_{hij} be the value of variable X for the j th member in cluster i in the h th stratum.

The ratio of Y to X is

$$\hat{R} = \frac{\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} w_{hij} y_{hij}}{\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} w_{hij} x_{hij}}$$

PROC SURVEYMEANS uses the Taylor series method to estimate the variance of the ratio \hat{R} as

$$\hat{V}(\hat{R}) = \sum_{h=1}^H \hat{V}_h(\hat{R})$$

where if $n_h > 1$,

$$\begin{aligned} \hat{V}_h(\hat{R}) &= \frac{n_h(1 - f_h)}{n_h - 1} \sum_{i=1}^{n_h} (g_{hi\cdot} - \bar{g}_{h\cdot\cdot})^2 \\ g_{hi\cdot} &= \frac{\sum_{j=1}^{m_{hi}} w_{hij} (y_{hij} - x_{hij} \hat{R})}{\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} w_{hij} x_{hij}} \\ \bar{g}_{h\cdot\cdot} &= \left(\sum_{i=1}^{n_h} g_{hi\cdot} \right) / n_h \end{aligned}$$

and if $n_h = 1$,

$$\hat{V}_h(\hat{R}) = \begin{cases} \text{missing} & \text{if } n_{h'} = 1 \text{ for } h' = 1, 2, \dots, H \\ 0 & \text{if } n_{h'} > 1 \text{ for some } 1 \leq h' \leq H \end{cases}$$

The standard error of the ratio is the square root of the estimated variance:

$$\text{StdErr}(\widehat{R}) = \sqrt{\widehat{V}(\widehat{R})}$$

When the denominator for a ratio is zero, then the value of the ratio is displayed as ‘-Infy’, ‘Infy’, or a missing value, depending on whether the numerator is negative, positive, or zero, respectively; and the corresponding internal value is the special missing value ‘.M’, the special missing value ‘.I’, or the usual missing value, respectively.

Domain Statistics

When you use a **DOMAIN** statement to request a domain analysis, the procedure computes the requested statistics for each domain.

For a domain D , let I_D be the corresponding indicator variable:

$$I_D(h, i, j) = \begin{cases} 1 & \text{if observation } (h, i, j) \text{ belongs to } D \\ 0 & \text{otherwise} \end{cases}$$

Let

$$v_{hij} = w_{hij} I_D(h, i, j) = \begin{cases} w_{hij} & \text{if observation } (h, i, j) \text{ belongs to } D \\ 0 & \text{otherwise} \end{cases}$$

The requested statistics for variable y in domain D are computed by using the new weights v .

Domain Mean

The estimated mean of y in the domain D is

$$\widehat{\bar{Y}}_D = \left(\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} v_{hij} y_{hij} \right) / v_{\dots}$$

where

$$v_{\dots} = \sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} v_{hij}$$

The variance of $\widehat{\bar{Y}}_D$ is estimated by

$$\widehat{V}(\widehat{\bar{Y}}_D) = \sum_{h=1}^H \widehat{V}_h(\widehat{\bar{Y}}_D)$$

where if $n_h > 1$,

$$\begin{aligned}\widehat{V}_h(\widehat{Y}_D) &= \frac{n_h(1-f_h)}{n_h-1} \sum_{i=1}^{n_h} (r_{hi\cdot} - \bar{r}_{h\cdot\cdot})^2 \\ r_{hi\cdot} &= \left(\sum_{j=1}^{m_{hi}} v_{hij} (y_{hij} - \widehat{Y}_D) \right) / v_{hi\cdot} \\ \bar{r}_{h\cdot\cdot} &= \left(\sum_{i=1}^{n_h} r_{hi\cdot} \right) / n_h\end{aligned}$$

and if $n_h = 1$,

$$\widehat{V}_h(\widehat{Y}_D) = \begin{cases} \text{missing} & \text{if } n_{h'} = 1 \text{ for } h' = 1, 2, \dots, H \\ 0 & \text{if } n_{h'} > 1 \text{ for some } 1 \leq h' \leq H \end{cases}$$

Domain Total

The estimated total in domain D is

$$\widehat{Y}_D = \sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} v_{hij} y_{hij}$$

and its estimated variance is

$$\widehat{V}(\widehat{Y}_D) = \sum_{h=1}^H \widehat{V}_h(\widehat{Y}_D)$$

where if $n_h > 1$,

$$\begin{aligned}\widehat{V}_h(\widehat{Y}_D) &= \frac{n_h(1-f_h)}{n_h-1} \sum_{i=1}^{n_h} (z_{hi\cdot} - \bar{z}_{h\cdot\cdot})^2 \\ z_{hi\cdot} &= \sum_{j=1}^{m_{hi}} v_{hij} z_{hij} \\ \bar{z}_{h\cdot\cdot} &= \left(\sum_{i=1}^{n_h} z_{hi\cdot} \right) / n_h\end{aligned}$$

and if $n_h = 1$,

$$\widehat{V}_h(\widehat{Y}_D) = \begin{cases} \text{missing} & \text{if } n_{h'} = 1 \text{ for } h' = 1, 2, \dots, H \\ 0 & \text{if } n_{h'} > 1 \text{ for some } 1 \leq h' \leq H \end{cases}$$

Domain Ratio

The estimated ratio of Y to X in domain D is

$$\hat{R}_D = \frac{\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} v_{hij} y_{hij}}{\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} v_{hij} x_{hij}}$$

and its estimated variance is

$$\hat{V}(\hat{R}_D) = \sum_{h=1}^H \hat{V}_h(\hat{R}_D)$$

where if $n_h > 1$,

$$\begin{aligned} \hat{V}_h(\hat{R}_D) &= \frac{n_h(1-f_h)}{n_h-1} \sum_{i=1}^{n_h} (g_{hi\cdot} - \bar{g}_{h\cdot\cdot})^2 \\ g_{hi\cdot} &= \frac{\sum_{j=1}^{m_{hi}} v_{hij} (y_{hij} - x_{hij} \hat{R}_D)}{\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} v_{hij} x_{hij}} \\ \bar{g}_{h\cdot\cdot} &= \left(\sum_{i=1}^{n_h} g_{hi\cdot} \right) / n_h \end{aligned}$$

and if $n_h = 1$,

$$\hat{V}_h(\hat{R}_D) = \begin{cases} \text{missing} & \text{if } n_{h'} = 1 \text{ for } h' = 1, 2, \dots, H \\ 0 & \text{if } n_{h'} > 1 \text{ for some } 1 \leq h' \leq H \end{cases}$$

Quantiles

Let Y be the variable of interest in a complex survey. Denote $F(t) = Pr(Y \leq t)$ as the cumulative distribution for Y . For $0 < p < 1$, the p th quantile of the population cumulative distribution function is

$$Y_p = \inf\{y : F(y) \geq p\}$$

Estimate of Quantile

Let $\{y_{hij}, w_{hij}\}$ be the observed values for variable Y associated with sampling weights, where (h, i, j) are the stratum index, cluster index, and member index, respectively, as shown in the section “Definitions and Notation” on page 7268. Let $y_{(1)} < y_{(2)} < \dots < y_{(d)}$ denote the sample order statistics for variable Y .

An estimate of quantile Y_p is

$$\hat{Y}_p = \begin{cases} y_{(1)} & \text{if } p < \hat{F}(y_{(1)}) \\ y_{(k)} + \frac{p - \hat{F}(y_{(k)})}{\hat{F}(y_{(k+1)}) - \hat{F}(y_{(k)})} (y_{(k+1)} - y_{(k)}) & \text{if } \hat{F}(y_{(k)}) \leq p < \hat{F}(y_{(k+1)}) \\ y_{(d)} & \text{if } p = 1 \end{cases}$$

where $\hat{F}(t)$ is the estimated cumulative distribution for Y :

$$\hat{F}(t) = \frac{\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} w_{hij} I(y_{hij} \leq t)}{\sum_{h=1}^H \sum_{i=1}^{n_h} \sum_{j=1}^{m_{hi}} w_{hij}}$$

and $I(\cdot)$ is the indicator function.

Standard Error

PROC SURVEYMEANS uses Woodruff's method (Dorfman and Valliant 1993; Särndal, Swensson, and Wretman 1992; and Francisco and Fuller 1991) to estimate the variances of quantiles. This method first constructs a confidence interval on a quantile. Then it uses the width of the confidence interval to estimate the standard error of a quantile.

In order to estimate the variance for \hat{Y}_p , first the procedure estimates the variance of the estimated distribution function $\hat{F}(\hat{Y}_p)$ by

$$\hat{V}(\hat{F}(\hat{Y}_p)) = \sum_{h=1}^H \frac{n_h(1-f_h)}{n_h-1} \sum_{i=1}^{n_h} (d_{hi\cdot} - \bar{d}_{h\cdot\cdot})^2$$

where

$$d_{hi\cdot} = \left(\sum_{j=1}^{m_{hi}} w_{hij} (I(y_{hij} \leq \hat{Y}_p) - \hat{F}(\hat{Y}_p)) \right) / w_{h\cdot\cdot}$$

$$\bar{d}_{h\cdot\cdot} = \left(\sum_{i=1}^{n_h} d_{hi\cdot} \right) / n_h$$

Then $100(1-\alpha)\%$ confidence limits of $\hat{F}(\hat{Y}_p)$ can be constructed by

$$(\hat{p}_L, \hat{p}_U) = \left(\hat{F}(\hat{Y}_p) - t_{df, \alpha/2} \sqrt{\hat{V}(\hat{F}(\hat{Y}_p))}, \hat{F}(\hat{Y}_p) + t_{df, \alpha/2} \sqrt{\hat{V}(\hat{F}(\hat{Y}_p))} \right)$$

where $t_{df, \alpha/2}$ is the $100(1-\alpha/2)$ th percentile of the t distribution with df degrees of freedom, described in the section “[Degrees of Freedom](#)” on page 7271.

When (\hat{p}_L, \hat{p}_U) is out of the range of $[0,1]$, the procedure does not compute the standard error.

The \hat{p}_L th quantile is defined as

$$\hat{Y}_{\hat{p}_L} = \begin{cases} y_{(1)} & \text{if } \hat{p}_L < \hat{F}(y_{(1)}) \\ y_{(k_L)} + \frac{\hat{p}_L - \hat{F}(y_{(k_L)})}{\hat{F}(y_{(k_L+1)}) - \hat{F}(y_{(k_L)})} (y_{(k_L+1)} - y_{(k_L)}) & \text{if } \hat{F}(y_{(k_L)}) \leq \hat{p}_L < \hat{F}(y_{(k_L+1)}) \\ y_{(d)} & \text{if } \hat{p}_L = 1 \end{cases}$$

and the \hat{p}_U th quantile is defined as

$$\hat{Y}_{\hat{p}_U} = \begin{cases} y_{(1)} & \text{if } \hat{p}_U < \hat{F}(y_{(1)}) \\ y_{(k_U)} + \frac{\hat{p}_U - \hat{F}(y_{(k_U)})}{\hat{F}(y_{(k_U+1)}) - \hat{F}(y_{(k_U)})} (y_{(k_U+1)} - y_{(k_U)}) & \text{if } \hat{F}(y_{(k_U)}) \leq \hat{p}_U < \hat{F}(y_{(k_U+1)}) \\ y_{(d)} & \text{if } \hat{p}_U = 1 \end{cases}$$

The standard error of \hat{Y}_p then is estimated by

$$sd(\hat{Y}_p) = \frac{\hat{Y}_{\hat{p}_U} - \hat{Y}_{\hat{p}_L}}{2t_{df, \alpha/2}}$$

where $t_{df, \alpha/2}$ is the $100(1 - \alpha/2)$ th percentile of the t distribution with df degrees of freedom.

Confidence Limits

Symmetric $100(1 - \alpha)\%$ confidence limits are computed as

$$\left(\hat{Y}_p - sd(\hat{Y}_p) \cdot t_{df, \alpha/2}, \quad \hat{Y}_p + sd(\hat{Y}_p) \cdot t_{df, \alpha/2} \right)$$

If you specify the **NONSYMCL** option in the SURVEYMEANS statement, the procedure computes $100(1 - \alpha)\%$ nonsymmetric confidence limits:

$$\left(\hat{Y}_{\hat{p}_L}, \quad \hat{Y}_{\hat{p}_U} \right)$$

Replication Methods for Variance Estimation

Recently replication methods have gained popularity for estimating variances in complex survey data analysis. One reason for this popularity is the relative simplicity of replication-based estimates, especially for nonlinear estimators; another is that modern computational capacity has made replication methods feasible for practical survey analysis. For details see Lohr (2009); Wolter (1985); Rust (1985); Dippo, Fay, and Morganstein (1984); Rao and Shao (1999); Rao, Wu, and Yue (1992); and Rao and Shao (1996).

Replication methods draw multiple replicates (also called subsamples) from a full sample according to a specific resampling scheme. The most commonly used resampling schemes are the *balanced repeated replication* (BRR) method and the *jackknife* method. For each replicate, the original weights are modified for the PSUs in the replicates to create replicate weights. The statistics of interest are estimated by using the replicate weights for each replicate. Then the variances of parameters of interest are estimated by the variability among the estimates derived from these replicates. You can use the **REPWEIGHTS** statement to provide your own replicate weights for variance estimation.

Balanced Repeated Replication (BRR) Method

The balanced repeated replication (BRR) method requires that the full sample be drawn by using a stratified sample design with two primary sampling units (PSUs) per stratum. Let H be the total number of strata. The total number of replicates R is the smallest multiple of 4 that is greater than H . However, if you prefer a larger number of replicates, you can specify the **REPS=number** option. If a $number \times number$ Hadamard matrix cannot be constructed, the number of replicates is increased until a Hadamard matrix becomes available.

Each replicate is obtained by deleting one PSU per stratum according to the corresponding [Hadamard matrix](#) and adjusting the original weights for the remaining PSUs. The new weights are called replicate weights.

Replicates are constructed by using the first H columns of the $R \times R$ [Hadamard matrix](#). The r th ($r = 1, 2, \dots, R$) replicate is drawn from the full sample according to the r th row of the Hadamard matrix as follows:

- If the (r, h) th element of the Hadamard matrix is 1, then the first PSU of stratum h is included in the r th replicate and the second PSU of stratum h is excluded.
- If the (r, h) th element of the Hadamard matrix is -1 , then the second PSU of stratum h is included in the r th replicate and the first PSU of stratum h is excluded.

The replicate weights of the remaining PSUs in each half-sample are then doubled to their original weights. For more details about the BRR method, see Wolter (1985) and Lohr (2009).

By default, an appropriate Hadamard matrix is generated automatically to create the replicates. You can request that the Hadamard matrix be displayed by specifying the [VARMETHOD=BRR\(PRINTH\)](#) *method-option*. If you provide a Hadamard matrix by specifying the [VARMETHOD=BRR\(HADAMARD=\)](#) *method-option*, then the replicates are generated according to the provided Hadamard matrix.

You can use the [VARMETHOD=BRR\(OUTWEIGHTS=\)](#) *method-option* to save the replicate weights into a SAS data set.

Suppose that θ is a population parameter of interest. Let $\hat{\theta}$ be the estimate from the full sample for θ . Let $\hat{\theta}_r$ be the estimate from the r th replicate subsample by using replicate weights. PROC SURVEYMEANS estimates the variance of $\hat{\theta}$ by

$$\widehat{V}(\hat{\theta}) = \frac{1}{R} \sum_{r=1}^R (\hat{\theta}_r - \hat{\theta})^2$$

with H degrees of freedom, where H is the number of strata.

If a parameter cannot be computed from one or more replicates, then the variance estimate is computed by using those replicates from which the parameter can be estimated. For example, suppose the parameter is a ratio. If a replicate r contains observations such that the denominator of the ratio is zero, then the ratio cannot be computed from replicate r . In this case, the BRR variance estimate is computed as

$$\widehat{V}(\hat{\theta}) = \frac{1}{R'} \sum_{r=1}^{R'} (\hat{\theta}_r - \hat{\theta})^2$$

where the summation is over the replicates where the parameter θ can be computed, and R' is the number of those replicates.

Fay's BRR Method

Fay's method is a modification of the BRR method, and it requires a stratified sample design with two primary sampling units (PSUs) per stratum. The total number of replicates R is the smallest multiple of 4 that is greater than the total number of strata H . However, if you prefer a larger number of replicates, you can specify the `REPS=method-option`.

For each replicate, Fay's method uses a Fay coefficient $0 \leq \epsilon < 1$ to impose a perturbation of the original weights in the full sample that is gentler than using only half-samples, as in the traditional BRR method. The Fay coefficient $0 \leq \epsilon < 1$ can be set by specifying the `FAY = ϵ method-option`. By default, $\epsilon = 0.5$ if the `FAY method-option` is specified without providing a value for ϵ (Judkins 1990; Rao and Shao 1999). When $\epsilon = 0$, Fay's method becomes the traditional BRR method. For more details, see Dippo, Fay, and Morganstein (1984), Fay (1984), Fay (1989), and Judkins (1990).

Let H be the number of strata. Replicates are constructed by using the first H columns of the $R \times R$ Hadamard matrix, where R is the number of replicates, $R > H$. The r th ($r = 1, 2, \dots, R$) replicate is created from the full sample according to the r th row of the Hadamard matrix as follows:

- If the (r, h) th element of the Hadamard matrix is 1, then the full sample weight of the first PSU in stratum h is multiplied by ϵ and the full sample weight of the second PSU is multiplied by $2 - \epsilon$ to obtain the r th replicate weights.
- If the (r, h) th element of the Hadamard matrix is -1 , then the full sample weight of the first PSU in stratum h is multiplied by $2 - \epsilon$ and the full sample weight of the second PSU is multiplied by ϵ to obtain the r th replicate weights.

You can use the `VARMETHOD=BRR(OUTWEIGHTS=) method-option` to save the replicate weights into a SAS data set.

By default, an appropriate Hadamard matrix is generated automatically to create the replicates. You can request that the Hadamard matrix be displayed by specifying the `VARMETHOD=BRR(PRINTH) method-option`. If you provide a Hadamard matrix by specifying the `VARMETHOD=BRR(HADAMARD=) method-option`, then the replicates are generated according to the provided Hadamard matrix.

Suppose that θ is a population parameter of interest. Let $\hat{\theta}$ be the estimate from the full sample for θ . Let $\hat{\theta}_r$ be the estimate from the r th replicate subsample by using replicate weights. PROC SURVEYMEANS estimates the variance of θ by

$$\widehat{V}(\hat{\theta}) = \frac{1}{R(1 - \epsilon)^2} \sum_{r=1}^R (\hat{\theta}_r - \hat{\theta})^2$$

with H degrees of freedom, where H is the number of strata.

Jackknife Method

The jackknife method of variance estimation deletes one PSU at a time from the full sample to create replicates. The total number of replicates R is the same as the total number of PSUs. In each

replicate, the sample weights of the remaining PSUs are modified by the jackknife coefficient α_r . The modified weights are called replicate weights.

The jackknife coefficient and replicate weights are described as follows.

Without Stratification If there is no stratification in the sample design (no **STRATA** statement), the jackknife coefficients α_r are the same for all replicates:

$$\alpha_r = \frac{R-1}{R} \quad \text{where } r = 1, 2, \dots, R$$

Denote the original weight in the full sample for the j th member of the i th PSU as w_{ij} . If the i th PSU is included in the r th replicate ($r = 1, 2, \dots, R$), then the corresponding replicate weight for the j th member of the i th PSU is defined as

$$w_{ij}^{(r)} = w_{ij} / \alpha_r$$

With Stratification If the sample design involves stratification, each stratum must have at least two PSUs to use the jackknife method.

Let stratum \tilde{h}_r be the stratum from which a PSU is deleted for the r th replicate. Stratum \tilde{h}_r is called the *donor stratum*. Let $n_{\tilde{h}_r}$ be the total number of PSUs in the donor stratum \tilde{h}_r . The jackknife coefficients are defined as

$$\alpha_r = \frac{n_{\tilde{h}_r} - 1}{n_{\tilde{h}_r}} \quad \text{where } r = 1, 2, \dots, R$$

Denote the original weight in the full sample for the j th member of the i th PSU as w_{ij} . If the i th PSU is included in the r th replicate ($r = 1, 2, \dots, R$), then the corresponding replicate weight for the j th member of the i th PSU is defined as

$$w_{ij}^{(r)} = \begin{cases} w_{ij} & \text{if } i\text{th PSU is not in the donor stratum } \tilde{h}_r \\ w_{ij} / \alpha_r & \text{if } i\text{th PSU is in the donor stratum } \tilde{h}_r \end{cases}$$

You can use the **VARMETHOD=JACKKNIFE(OUTJKCOEFS=)** *method-option* to save the jackknife coefficients into a SAS data set and use the **VARMETHOD=JACKKNIFE(OUTWEIGHTS=)** *method-option* to save the replicate weights into a SAS data set.

If you provide your own replicate weights with a **REPWEIGHTS** statement, then you can also provide corresponding jackknife coefficients with the **JKCOEFS=** option.

Suppose that θ is a population parameter of interest. Let $\hat{\theta}$ be the estimate from the full sample for θ . Let $\hat{\theta}_r$ be the estimate from the r th replicate subsample by using replicate weights. PROC SURVEYMEANS estimates the variance of $\hat{\theta}$ by

$$\widehat{V}(\hat{\theta}) = \sum_{r=1}^R \alpha_r (\hat{\theta}_r - \hat{\theta})^2$$

with $R - H$ degrees of freedom, where R is the number of replicates and H is the number of strata, or $R - 1$ when there is no stratification.

Hadamard Matrix

A Hadamard matrix \mathbf{H} is a square matrix whose elements are either 1 or -1 such that

$$\mathbf{H}\mathbf{H}' = k\mathbf{I}$$

where k is the dimension of \mathbf{H} and \mathbf{I} is the identity matrix of order k . The order k is necessarily 1, 2, or a positive integer that is a multiple of 4.

For example, the following matrix is a Hadamard matrix of dimension $k = 8$:

1	1	1	1	1	1	1	1
1	-1	1	-1	1	-1	1	-1
1	1	-1	-1	1	1	-1	-1
1	-1	-1	1	1	-1	-1	1
1	1	1	1	-1	-1	-1	-1
1	-1	1	-1	-1	1	-1	1
1	1	-1	-1	-1	-1	1	1
1	-1	-1	1	-1	1	1	-1

Output Data Sets

You can use the Output Delivery System to create a SAS data set from any piece of PROC SURVEYMEANS output. See the section “[ODS Table Names](#)” on page 7293 for more information.

PROC SURVEYMEANS also provides an output data set that stores the replicate weights for BRR or jackknife variance estimation and an output data set that stores the jackknife coefficients for jackknife variance estimation.

Replicate Weights Output Data Set

If you specify the `OUTWEIGHTS=` *method-option* for `VARMETHOD=BRR` or `JACKKNIFE`, PROC SURVEYMEANS stores the replicate weights in an output data set. The `OUTWEIGHTS=` output data set contains all observations from the `DATA=` input data set that are valid (used in the analysis). (A valid observation is an observation that has a positive value of the `WEIGHT` variable. Valid observations must also have nonmissing values of the `STRATA` and `CLUSTER` variables, unless you specify the `MISSING` option. See the section “[Data and Sample Design Summary](#)” on page 7287 for details about valid observations.)

The `OUTWEIGHTS=` data set contains the following variables:

- all variables in the `DATA=` input data set
- `RepWt_1`, `RepWt_2`, . . . , `RepWt_n`, which are the replicate weight variables

where n is the total number of replicates in the analysis. Each replicate weight variable contains the replicate weights for the corresponding replicate. Replicate weights equal zero for those observations not included in the replicate.

After the procedure creates replicate weights for a particular input data set and survey design, you can use the `OUTWEIGHTS= method-option` to store these replicate weights and then use them again in subsequent analyses, either in PROC SURVEYMEANS or in the other survey procedures. You can use the `REPWEIGHTS` statement to provide replicate weights for the procedure.

Jackknife Coefficients Output Data Set

If you specify the `OUTJKCOEFS= method-option` for `VARMETHOD=JACKKNIFE`, PROC SURVEYMEANS stores the jackknife coefficients in an output data set. The `OUTJKCOEFS=` output data set contains one observation for each replicate. The `OUTJKCOEFS=` data set contains the following variables:

- Replicate, which is the replicate number for the jackknife coefficient
- JKCoefficient, which is the jackknife coefficient
- DonorStratum, which is the stratum of the PSU that was deleted to construct the replicate, if you specify a `STRATA` statement

After the procedure creates jackknife coefficients for a particular input data set and survey design, you can use the `OUTJKCOEFS= method-option` to store these coefficients and then use them again in subsequent analyses, either in PROC SURVEYMEANS or in the other survey procedures. You can use the `JKCOEFS=` option in the `REPWEIGHTS` statement to provide jackknife coefficients for the procedure.

Rectangular and Stacking Structures in an Output Data Set

When you use an ODS output statement to create SAS data sets for certain tables in PROC SURVEYMEANS, there are two possible types of table structure for the output data sets: *rectangular* and *stacking*. A rectangular structure creates one observation for each analysis variable in the data set. A stacking structure creates only one observation in the output data set for all analysis variables.

Before SAS 9, the stacking table structure, similar to the table structure in PROC MEANS, was the default in PROC SURVEYMEANS. Since SAS 9, the new default is to produce a rectangular table in the output data sets. You can use the `STACKING` option to request that the procedure produce the output data sets by using a stacking table structure.

The `STACKING` option affects the following tables:

- Domain
- Ratio
- Statistics
- StrataInfo

Figure 86.6 and Figure 86.7 shows these two structures for analyzing the following data set:

```
data new;
  input sex$ x;
  datalines;
M 12
F 5
M 13
F 23
F 11
;
```

The following statements request the default rectangular structure of the output data set for the statistics table:

```
proc surveymeans data=new mean;
  ods output statistics=rectangle;
run;

proc print data=rectangle;
run;
```

Figure 86.6 shows the rectangular structure.

Figure 86.6 Rectangular Structure in the Output Data Set

Rectangular Structure in the Output Data Set				
Obs	Var Name	Var Level	Mean	StdErr
1	x		12.800000	2.905168
2	sex	F	0.600000	0.244949
3	sex	M	0.400000	0.244949

The following statements specify the **STACKING** option to request that the output data set have a stacking structure:

```
proc surveymeans data=new mean stacking;
  ods output statistics=stacking;
run;

proc print data=stacking;
run;
```

Figure 86.7 shows the stacking structure of the output data set for the statistics table requested by the **STACKING** option.

Figure 86.7 Stacking Structure in the Output Data Set Requested by the STACKING option

Stacking Structure in the Output Data Set					
Obs	x	x_Mean	x_StdErr	sex_F	sex_F_Mean
1	x	12.800000	2.905168	sex=F	0.600000
Obs	sex_F_StdErr	sex_M	sex_M_Mean	sex_M_StdErr	
1	0.244949	sex=M	0.400000	0.244949	

Displayed Output

The SURVEYMEANS procedure produces output that is described in the following sections.

Data and Sample Design Summary

The “Data Summary” table provides information about the input data set and the sample design. This table displays the total number of valid observations, where an observation is considered *valid* if it has nonmissing values for all procedure variables other than the analysis variables—that is, for all specified [STRATA](#), [CLUSTER](#), and [WEIGHT](#) variables. This number might differ from the number of nonmissing observations for an individual analysis variable, which the procedure displays in the “Statistics” table. See the section “[Missing Values](#)” on page 7265 for more information.

PROC SURVEYMEANS displays the following information in the “Data Summary” table:

- Number of Strata, if you specify a STRATA statement
- Number of Clusters, if you specify a CLUSTER statement
- Number of Observations, which is the total number of valid observations
- Sum of Weights, which is the sum over all valid observations, if you specify a WEIGHT statement

Class Level Information

If you use a [CLASS](#) statement to name classification variables for categorical analysis, or if you list any character variables in the VAR statement, then PROC SURVEYMEANS displays a “Class Level Information” table. This table contains the following information for each classification variable:

- Class Variable, which lists each CLASS variable name
- Levels, which is the number of values or levels of the classification variable

- Values, which lists the values of the classification variable. The values are separated by a white space character; therefore, to avoid confusion, you should not include a white space character within a classification variable value.

Stratum Information

If you specify the **LIST** option in the STRATA statement, PROC SURVEYMEANS displays a “Stratum Information” table. This table displays the number of valid observations in each stratum, as well as the number of nonmissing stratum observations for each analysis variable. The “Stratum Information” table provides the following for each stratum:

- Stratum Index, which is a sequential stratum identification number
- STRATA variable(s), which lists the levels of **STRATA** variables for the stratum
- Population Total, if you specify the **TOTAL=** option
- Sampling Rate, if you specify the **TOTAL=** or **RATE=** option. If you specify the **TOTAL=** option, the sampling rate is based on the number of valid observations in the stratum.
- N Obs, which is the number of valid observations
- Variable, which lists each analysis variable name
- Levels, which identifies each level for categorical variables
- N, which is the number of nonmissing observations for the analysis variable
- Clusters, which is the number of clusters, if you specify a **CLUSTER** statement

Variance Estimation

If the variance method is not Taylor series or if the **NOMCAR** option is used, by default, PROC SURVEYMEANS displays the following variance estimation specifications in the “Variance Estimation” table:

- Method, which is the variance estimation method
- Replicate Weights Data Set, which is the name of the SAS data set that contains the replicate weights
- Number of Replicates, which is the number of replicates if you specify the **VARMETHOD=BRR** or **VARMETHOD=JACKKNIFE** option
- Hadamard Data Set, which is the name of the SAS data set for the HADAMARD matrix if you specify the **VARMETHOD=BRR(HADAMARD=)** *method-option*

- Fay Coefficient, which is the value of the FAY coefficient if you specify the **VARMETHOD=BRR(FAY)** *method-option*
- Missing Levels Included (MISSING), if you specify the **MISSING** option
- Missing Levels Included (NOMCAR), if you specify the **NOMCAR** option

Statistics

The “Statistics” table displays all of the statistics that you request with *statistic-keywords* in the PROC SURVEYMEANS statement, except DECILES, MEDIAN, Q1, Q3, and QUARTILES, which are displayed in the “Quantiles” table. If you do not specify any *statistic-keywords*, then by default this table displays the following information for each analysis variable: the sample size, the mean, the standard error of the mean, and the confidence limits for the mean. The “Statistics” table can contain the following information for each analysis variable, depending on which *statistic-keywords* you request:

- Variable name
- Variable Label
- Level, which identifies each level for categorical variables
- N, which is the number of nonmissing observations
- N Miss, which is the number of missing observations
- Minimum
- Maximum
- Range
- Number of Clusters
- Sum of Weights
- DF, which is the degrees of freedom for the t test
- Mean
- Std Error of Mean, which is the standard error of the mean
- Var of Mean, which is the variance of the mean
- t Value, for testing H_0 : population MEAN = 0
- $\text{Pr} > |t|$, which is the two-sided p -value for the t test
- $100(1 - \alpha)\%$ CL for Mean, which are two-sided confidence limits for the mean

- $100(1 - \alpha)\%$ Upper CL for Mean, which is a one-sided upper confidence limit for the mean
- $100(1 - \alpha)\%$ Lower CL for Mean, which is a one-sided lower confidence limit for the mean
- Coeff of Variation, which is the coefficient of variation for the mean
- Sum
- Std Dev, which is the standard deviation of the sum
- Var of Sum, which is the variance of the sum
- $100(1 - \alpha)\%$ CL for Sum, which are two-sided confidence limits for the sum
- $100(1 - \alpha)\%$ Upper CL for Sum, which is a one-sided upper confidence limit for the sum
- $100(1 - \alpha)\%$ Lower CL for Sum, which is a one-sided lower confidence limit for the Sum
- Coeff of Variation for sum, which is the coefficient of variation for the sum

Quantiles

The “Quantiles” table displays all the quantiles that you request with either *statistic-keywords* such as DECILES, MEDIAN, Q1, Q3, and QUARTILES, or the **PERCENTILE=** option, or the **QUANTILE=** option in the PROC SURVEYMEANS statement.

The “Quantiles” table contains the following information for each quantile:

- Variable name
- Variable Label
- Percentile, which is the requested quantile in the format of %
- Percentile Label, which is the corresponding common name for a percentile if it exists—for example, *Median* for 50th percentile
- Estimate, which is the estimate for a requested quantile with respect to the population distribution
- Std Error, which is the standard error of the quantile
- $100(1 - \alpha)\%$ Confidence Limits, which are two-sided confidence limits for the quantile

Domain Analysis

If you specify a **DOMAIN** statement, the procedure displays domain statistics in a “Domain Analysis” table. A “Domain Analysis” table displays all the requested statistics for each level of the domain request. The procedure produces a separate “Domain Analysis” for each separate domain request. For example, the **DOMAIN** statement

```
domain A B*C*D A*C C;
```

specifies four domain requests:

- A: all the levels of A
- C: all the levels of C
- A*C: all the interactive levels of A and C
- B*C*D: all the interactive levels of B, C, and D

The procedure displays four “Domain Analysis” tables, one for each domain definition. If you use an **ODS OUTPUT** statement to create an output data set for domain analysis, the output data set contains a variable **Domain** whose values are these domain definitions.

A “Domain Analysis” table contains all the columns in the “Statistics” table, plus columns of domain variable values.

Ratio Analysis

The “Ratio Analysis” table displays statistics for all the ratios that you request in the **RATIO** statement. If you do not specify any *statistic-keywords* in the **PROC SURVEYMEANS** statement, then by default this table displays the ratios and standard errors. The “Ratio Analysis” table can contain the following information for each ratio, depending on which *statistic-keywords* you request:

- Numerator, which identifies the numerator variable of the ratio
- Denominator, which identifies the denominator variable of the ratio
- N, which is the number of observations used in the ratio analysis
- number of Clusters
- Sum of Weights
- DF, which is the degrees of freedom for the *t* test
- Ratio
- Std Err of Ratio, which is the standard error of the ratio
- Var, which is the variance of the ratio

- t Value, for testing H_0 : population $\text{RATIO} = 0$
- $\text{Pr} > |t|$, which is the two-sided p -value for the t test
- $100(1 - \alpha)\%$ CL for Ratio, which are two-sided confidence limits for the Ratio
- Upper $100(1 - \alpha)\%$ CL for Ratio, which are one-sided upper confidence limits for the Ratio
- Lower $100(1 - \alpha)\%$ CL for Ratio, which are one-sided lower confidence limits for the Ratio

When you use the ODS OUTPUT statement to create an output data set, if you use labels for your RATIO statement, these labels are saved in the variable Ratio Statement in the output data set.

Domain Ratio Analysis

If you specify a **DOMAIN** statement with a **RATIO** statement, the procedure displays domain ratios in a “Domain Ratio Analysis” table. A “Domain Ratio Analysis” table displays all the ratio statistics for each level of the domain request.

A “Domain Ratio Analysis” table contains all the columns in the “Ratio Analysis” table, plus columns of domain variable values.

Hadamard Matrix

If you specify the **VARMETHOD=BRR(PRINTH)** *method-option* in the PROC SURVEYMEANS statement, PROC SURVEYMEANS displays the Hadamard matrix used to construct replicates for BRR variance estimation.

If you provide a Hadamard matrix with the **VARMETHOD=BRR(HADAMARD=)** *method-option* but the procedure does not use the entire matrix, the procedure displays only the rows and columns that are actually used to construct replicates.

ODS Table Names

PROC SURVEYMEANS assigns a name to each table it creates; these names are listed in [Table 86.3](#). You can use these names to refer to tables when you use the Output Delivery System (ODS) to select tables and create output data sets. For more information about ODS, see Chapter 20, “[Using the Output Delivery System](#).”

Table 86.3 ODS Tables Produced by PROC SURVEYMEANS

ODS Table Name	Description	Statement	Option
ClassVarInfo	Class level information	CLASS	Default
Domain	Statistics in domains	DOMAIN	Default
DomainRatio	Statistics for ratios in domains	DOMAIN and RATIO	Default
HadamardMatrix	Hadamard matrix	PROC	PRINTH
Ratio	Statistics for ratios	RATIO	Default
Quantiles	Quantiles	PROC	Default
Statistics	Statistics	PROC	Default
StrataInfo	Stratum information	STRATA	LIST
Summary	Data summary	PROC	Default
VarianceEstimation	Variance estimation	PROC	VARMETHOD=JK BRR or NOMCAR

For example, the following statements create an output data set MyStrata, which contains the “StrataInfo” table, and an output data set MyStat, which contains the “Statistics” table for the ice cream study discussed in the section “[Stratified Sampling](#)” on page 7243:

```

title1 'Analysis of Ice Cream Spending';
proc surveymeans data=IceCream total=StudentTotals;
  strata Grade / list;
  var Spending Group;
  weight Weight;
  ods output
    StrataInfo = MyStrata
    Statistics = MyStat;
run;

```

Examples: SURVEYMEANS Procedure

The section “Getting Started: SURVEYMEANS Procedure” on page 7241 contains examples of analyzing data from simple random sampling and stratified simple random sample designs. This section provides more examples that illustrate how to use PROC SURVEYMEANS.

Example 86.1: Stratified Cluster Sample Design

Consider the example in the section “Stratified Sampling” on page 7243. The study population is a junior high school with a total of 4,000 students in grades 7, 8, and 9. Researchers want to know how much these students spend weekly for ice cream, on the average, and what percentage of students spend at least \$10 weekly for ice cream.

The example in the section “Stratified Sampling” on page 7243 assumes that the sample of students was selected using a stratified simple random sample design. This example shows analysis based on a more complex sample design.

Suppose that every student belongs to a study group and that study groups are formed within each grade level. Each study group contains between two and four students. Table 86.4 shows the total number of study groups for each grade.

Table 86.4 Study Groups and Students by Grade

Grade	Number of Study Groups	Number of Students
7	608	1,824
8	252	1,025
9	403	1,151
Total	617	4,000

It is quicker and more convenient to collect data from students in the same study group than to collect data from students individually. Therefore, this study uses a stratified clustered sample design. The primary sampling units, or clusters, are study groups. The list of all study groups in the school is stratified by grade level. From each grade level, a sample of study groups is randomly selected, and all students in each selected study group are interviewed. The sample consists of eight study groups from the 7th grade, three groups from the 8th grade, and five groups from the 9th grade.

The SAS data set IceCreamStudy saves the responses of the selected students:

```
data IceCreamStudy;
  input Grade StudyGroup Spending @@;
  if (Spending < 10) then Group='less';
  else Group='more';
  datalines;
7 34 7 7 34 7 7 412 4 9 27 14
7 34 2 9 230 15 9 27 15 7 501 2
9 230 8 9 230 7 7 501 3 8 59 20
7 403 4 7 403 11 8 59 13 8 59 17
8 143 12 8 143 16 8 59 18 9 235 9
8 143 10 9 312 8 9 235 6 9 235 11
9 312 10 7 321 6 8 156 19 8 156 14
7 321 3 7 321 12 7 489 2 7 489 9
7 78 1 7 78 10 7 489 2 7 156 1
7 78 6 7 412 6 7 156 2 9 301 8
;
```

In the data set IceCreamStudy, the variable Grade contains a student's grade. The variable StudyGroup identifies a student's study group. It is possible for students from different grades to have the same study group number because study groups are sequentially numbered within each grade. The variable Spending contains a student's response regarding how much he spends per week for ice cream, in dollars. The variable GROUP indicates whether a student spends at least \$10 weekly for ice cream. It is not necessary to store the data in order of grade and study group.

The SAS data set StudyGroups is created to provide PROC SURVEYMEANS with the sample design information shown in [Table 86.4](#):

```
data StudyGroups;
  input Grade _total_;
  datalines;
7 608
8 252
9 403
;
```

The variable Grade identifies the strata, and the variable _TOTAL_ contains the total number of study groups in each stratum. As discussed in the section “[Specification of Population Totals and Sampling Rates](#)” on page 7266, the population totals stored in the variable _TOTAL_ should be expressed in terms of the primary sampling units (PSUs), which are study groups in this example. Therefore, the variable _TOTAL_ contains the total number of study groups for each grade, rather than the total number of students.

In order to obtain unbiased estimates, you create sampling weights by using the following SAS statements:

```
data IceCreamStudy;
    set IceCreamStudy;
    if Grade=7 then Prob=8/608;
    if Grade=8 then Prob=3/252;
    if Grade=9 then Prob=5/403;
    Weight=1/Prob;
run;
```

The sampling weights are the reciprocals of the probabilities of selections. The variable `Weight` contains the sampling weights. Because the sampling design is clustered and all students from each selected cluster are interviewed, the sampling weights equal the inverse of the cluster (or study group) selection probabilities.

The following SAS statements perform the analysis for this sample design:

```
title1 'Analysis of Ice Cream Spending';
proc surveymeans data=IceCreamStudy total=StudyGroups;
    strata Grade / list;
    cluster StudyGroup;
    var Spending Group;
    weight Weight;
run;
```

Output 86.1.1 provides information about the sample design and the input data set. There are three strata in the sample design, and the sample contains 16 clusters and 40 observations. The variable `Group` has two levels, 'less' and 'more.'

Output 86.1.1 Data Summary and Class Information

Analysis of Ice Cream Spending			
The SURVEYMEANS Procedure			
Data Summary			
Number of Strata			3
Number of Clusters			16
Number of Observations			40
Sum of Weights			3162.6
Class Level Information			
Class			
Variable	Levels	Values	
Group	2	less	more

Output 86.1.2 displays information for each stratum. Since the primary sampling units in this design are study groups, the population totals shown in Output 86.1.2 are the total numbers of study groups for each stratum or grade. This differs from Output 86.3, which provides the population totals in terms of students since students were the primary sampling units for that design. Output 86.1.2 also displays the number of clusters for each stratum and analysis variable.

Output 86.1.2 Stratum Information

Stratum Information						
Stratum Index	Grade	Population Total	Sampling Rate	N Obs	Variable	Level
1	7	608	1.32%	20	Spending Group	less more
2	8	252	1.19%	9	Spending Group	less more
3	9	403	1.24%	11	Spending Group	less more

Stratum Information						
Stratum Index	Grade	Population Total	Sampling Rate	N Obs	Variable	N
1	7	608	1.32%	20	Spending Group	20 17 3
2	8	252	1.19%	9	Spending Group	9 0 9
3	9	403	1.24%	11	Spending Group	11 6 5

Stratum Information						
Stratum Index	Grade	Population Total	Sampling Rate	N Obs	Variable	Clusters
1	7	608	1.32%	20	Spending Group	8 8 3
2	8	252	1.19%	9	Spending Group	3 0 3
3	9	403	1.24%	11	Spending Group	5 4 4

Output 86.1.3 displays the estimates of the average weekly ice cream expenditure and the percentage of students spending at least \$10 weekly for ice cream.

Output 86.1.3 Statistics

Statistics						
Variable	Level	N	Mean	Std Error of Mean	95% CL for Mean	
Spending		40	8.923860	0.650859	7.51776370	10.3299565
Group	less	23	0.561437	0.056368	0.43966057	0.6832130
	more	17	0.438563	0.056368	0.31678698	0.5603394

Example 86.2: Domain Analysis

Suppose that you are studying profiles of 800 top-performing companies to provide information about their impact on the economy. You are also interested in the company profiles within each market type. A sample of 66 companies is selected with unequal probability across market types. However, market type is not included in the sample design. Thus, the number of companies within each market type is a random variable in your sample. To obtain statistics within each market type, you should use domain analysis. The data of the 66 companies are saved in the following data set:

```
data Company;
  length Type $14;
  input Type$ Asset Sale Value Profit Employee Weight;
  datalines;
Other      2764.0  1828.0  1850.3   144.0   18.7    9.6
Energy     13246.2  4633.5  4387.7   462.9   24.3   42.6
Finance    3597.7   377.8   93.0    14.0    1.1   12.2
Transportation 6646.1  6414.2  2377.5   348.2   47.1   21.8
HiTech     1068.4  1689.8  1430.2    72.9    4.6    4.3
Manufacturing 1125.0  1719.4  1057.5    98.1   20.4    4.5
Other      1459.0  1241.4   452.7    24.5   20.1    5.5
Finance    2672.3   262.5   296.2    23.1    2.2    9.3
Finance     311.0   566.2   932.0    52.8    2.7    1.9
Energy     1148.6  1014.6   485.1    60.6    4.0    4.5
Finance    5327.0   572.4   372.9    25.2    4.2   17.7
Energy     1602.7   678.4   653.0    75.6    2.8    6.0
Energy     5808.8  1288.4  2007.0   318.8    5.9   19.2
Medical     268.8   204.4   820.9    45.6    3.7    1.8
Transportation 5222.6  2627.8  1910.0   245.6   22.8   17.4
Other       872.7  1419.4   939.3    69.7   12.2    3.7
Retail     4461.7  8946.8  4662.7   289.0  132.1   15.0
HiTech     6719.2  6942.0  8240.2   381.3   85.8   22.1
Retail      833.4  1538.8  1090.3    64.9   15.4    3.5
Finance     415.9   167.3  1126.8    56.8    0.7    2.2
HiTech      442.4  1139.9  1039.9    57.6   22.7    2.3
```

Other	801.5	1157.0	664.2	56.9	15.5	3.4
Finance	4954.8	468.8	366.4	41.7	3.0	16.5
Finance	2661.9	257.9	181.1	21.2	2.1	9.3
Finance	5345.8	530.1	337.4	36.4	4.3	17.8
Energy	3334.3	1644.7	1407.8	157.6	6.4	11.4
Manufacturing	1826.6	2671.7	483.2	71.3	25.3	6.7
Retail	618.8	2354.7	767.7	58.6	19.0	2.9
Retail	1529.1	6534.0	826.3	58.3	65.8	5.7
Manufacturing	4458.4	4824.5	3132.1	28.9	67.0	15.0
HiTech	5831.7	6611.1	9464.7	459.6	86.7	19.3
Medical	6468.3	4199.2	3170.4	270.1	59.5	21.3
Energy	1720.7	473.1	811.1	86.6	1.6	6.3
Energy	1679.7	1379.9	721.1	91.8	4.5	6.2
Retail	4018.2	16823.4	2038.3	178.1	162.0	13.6
Other	227.1	575.8	1083.8	62.6	1.9	1.6
Finance	3872.8	362.0	209.3	27.6	2.4	13.1
Retail	3359.3	4844.7	2651.4	224.1	75.6	11.5
Energy	1295.6	356.9	180.8	162.3	0.6	5.0
Energy	1658.0	626.6	688.0	126.0	3.5	6.1
Finance	12156.7	1345.5	680.7	106.6	9.4	39.2
HiTech	3982.6	4196.0	3946.8	313.9	64.3	13.5
Finance	8760.7	886.4	1006.9	90.0	7.5	28.5
Manufacturing	2362.2	3153.3	1080.0	137.0	25.2	8.4
Transportation	2499.9	3419.0	992.6	47.2	25.3	8.8
Energy	1430.4	1610.0	664.3	77.7	3.5	5.4
Energy	13666.5	15465.4	2736.7	411.4	26.6	43.9
Manufacturing	4069.3	4174.7	2907.6	289.2	38.2	13.7
Energy	2924.7	711.9	1067.8	146.7	3.4	10.1
Transportation	1262.1	1716.0	364.3	71.2	14.5	4.9
Medical	684.4	672.9	287.4	61.8	6.0	3.1
Energy	3069.3	1719.0	1439.0	196.4	4.9	10.6
Medical	246.5	318.8	924.1	43.8	3.1	1.7
Finance	11562.2	1128.5	580.4	64.2	6.7	37.3
Finance	9316.0	1059.4	816.5	95.9	8.0	30.2
Retail	1094.3	3848.0	563.3	29.4	44.7	4.4
Retail	1102.1	4878.3	932.4	65.2	47.3	4.4
HiTech	466.4	675.8	845.7	64.5	5.2	2.4
Manufacturing	10839.4	5468.7	1895.4	232.8	47.8	35.0
Manufacturing	733.5	2135.3	96.6	10.9	2.7	3.2
Manufacturing	10354.2	14477.4	5607.2	321.9	188.5	33.5
Energy	1902.1	2697.9	329.3	34.2	2.2	6.9
Other	2245.2	2132.2	2230.4	198.9	8.0	8.0
Transportation	949.4	1248.3	298.9	35.4	10.4	3.9
Retail	2834.4	2884.6	458.2	41.2	49.8	9.8
Retail	2621.1	6173.8	1992.7	183.7	115.1	9.2

;

For each company in your sample, the variables are defined as follows:

- Type identifies the type of market for the company.
- Asset contains the company's assets, in millions of dollars.
- Sale contains sales, in millions of dollars.

- Value contains the market value of the company, in millions of dollars.
- Profit contains the profit, in millions of dollars.
- Employee contains the number of employees, in thousands.
- Weight contains the sampling weight.

The following SAS statements use PROC SURVEYMEANS to perform the domain analysis, estimating means, and other statistics for the overall population and also for the subpopulations (or domain) defined by market type. The DOMAIN statement specifies Type as the domain variable:

```
title 'Top Companies Profile Study';
proc surveymeans data=Company total=800 mean sum;
  var Asset Sale Value Profit Employee;
  weight Weight;
  domain Type;
run;
```

Output 86.2.1 shows that there are 66 observations in the sample. The sum of the sampling weights equals 799.8, which is close to the total number of companies in the study population.

Output 86.2.1 Company Profile Study

Top Companies Profile Study				
The SURVEYMEANS Procedure				
Data Summary				
Number of Observations		66		
Sum of Weights		799.8		
Statistics				
Variable	Mean	Std Error of Mean	Sum	Std Dev
Asset	6523.488510	720.557075	5217486	1073829
Sale	4215.995799	839.132506	3371953	847885
Value	2145.935121	342.531720	1716319	359609
Profit	188.788210	25.057876	150993	30144
Employee	36.874869	7.787857	29493	7148.003298

The “Statistics” table in Output 86.2.1 displays the estimates of the mean and total for all analysis variables for the entire set of 800 companies, while Output 86.2.2 shows the mean and total estimates for each company type.

Output 86.2.2 Domain Analysis for Company Profile Study

Domain Analysis: Type					
Type	Variable	Mean	Std Error of Mean	Sum	Std Dev
Energy	Asset	7868.302932	1941.699163	1449341	785962
	Sale	5419.679099	2416.214417	998305	673373
	Value	2249.297177	520.295162	414321	213580
	Profit	289.564658	52.512141	53338	25927
	Employee	14.151194	3.974697	2606.650000	1481.777769
Finance	Asset	7890.190264	1057.185336	1855773	704506
	Sale	829.210502	115.762531	195030	74436
	Value	565.068197	76.964547	132904	48156
	Profit	63.716837	10.099341	14986	5801.108513
	Employee	5.806293	0.811555	1365.640000	519.658410
HiTech	Asset	5031.959781	732.436967	321542	183302
	Sale	5464.292019	731.296997	349168	196013
	Value	6707.828482	1194.160584	428630	249154
	Profit	346.407042	42.299004	22135	12223
	Employee	70.766980	8.683595	4522.010000	2524.778281
Manufacturing	Asset	7403.004250	1454.921083	888361	492577
	Sale	7207.638833	2112.444703	864917	501679
	Value	2986.442750	799.121544	358373	196979
	Profit	211.933583	39.993255	25432	13322
	Employee	83.314333	31.089019	9997.720000	6294.309490
Medical	Asset	5046.570609	1218.444638	140799	131942
	Sale	3313.219713	758.216303	92439	85655
	Value	2561.614695	530.802245	71469	64663
	Profit	218.682796	44.051447	6101.250000	5509.560969
	Employee	46.518996	11.135955	1297.880000	1213.651734
Other	Asset	1850.250000	338.128984	58838	31375
	Sale	1620.784906	168.686773	51541	24593
	Value	1432.820755	297.869828	45564	24204
	Profit	115.089937	27.970560	3659.860000	2018.201371
	Employee	14.306604	2.313733	454.950000	216.327710
Retail	Asset	2939.845750	393.692369	235188	94605
	Sale	7395.453500	1746.187580	591636	263263
	Value	2103.863125	529.756409	168309	78304
	Profit	157.171875	31.734253	12574	5478.281027
	Employee	93.624000	15.726743	7489.920000	3093.832061
Transportation	Asset	4712.047359	888.954411	267644	163516
	Sale	4030.233275	1015.555708	228917	142669
	Value	1703.330282	313.841326	96749	58947
	Profit	224.762324	56.168925	12767	8287.585418
	Employee	30.946303	6.786270	1757.750000	1066.586615

Example 86.3: Ratio Analysis

Suppose you are interested in the profit per employee and the sale per employee among the 800 top-performing companies in the data in the previous example. The following SAS statements illustrate how you can use PROC SURVEYMEANS to estimate these ratios:

```
title 'Ratio Analysis in Top Companies Profile Study';
proc surveymeans data=Company total=800 ratio;
  var Profit Sale Employee;
  weight Weight;
  ratio Profit Sale / Employee;
run;
```

The RATIO statement requests the ratio of the profit and the sales to the number of employees.

Output 86.3.1 shows the estimated ratios and their standard errors. Because the profit and the sales figures are in millions of dollars, and the employee numbers are in thousands, the profit per employee is estimated as \$5,120 with a standard error of \$1,059, and the sales per employee are \$114,332 with a standard error of \$20,503.

Output 86.3.1 Estimate Ratios

Ratio Analysis in Top Companies Profile Study			
The SURVEYMEANS Procedure			
Ratio Analysis			
Numerator	Denominator	Ratio	Std Err
Sale	Employee	114.332497	20.502742
Profit	Employee	5.119698	1.058939

Example 86.4: Analyzing Survey Data with Missing Values

As described in the section “[Missing Values](#)” on page 7265, the SURVEYMEANS procedure excludes an observation from the analysis if it has a missing value for the analysis variable or a nonpositive value for the WEIGHT variable.

However, if there is evidence indicating that the nonrespondents are different from the respondents for your study, you can use the NOMCAR option to compute descriptive statistics among respondents while still counting the number of nonrespondents.

We use the ice cream example in the section “[Stratified Sampling](#)” on page 7243 to illustrate how to perform similar analysis when there are missing values.

Suppose that some of the students failed to provide the amounts spent on ice cream, as shown in the following data set, IceCream:

```

data IceCream;
  input Grade Spending @@;
  if Grade=7 then Prob=20/1824;
  if Grade=8 then Prob=9/1025;
  if Grade=9 then Prob=11/1151;
  Weight=1/Prob;
  datalines;
7 7 7 7 8 . 9 10 7 . 7 10 7 3 8 20 8 19 7 2
7 . 9 15 8 16 7 6 7 6 7 6 9 15 8 17 8 14 9 .
9 8 9 7 7 3 7 12 7 4 9 14 8 18 9 9 7 2 7 1
7 4 7 11 9 8 8 . 8 13 7 . 9 . 9 11 7 2 7 9
;

data StudentTotals;
  input Grade _total_;
  datalines;
7 1824
8 1025
9 1151
;

```

Considering the possibility that those students who did not respond spend differently than those students who did respond, you can use the NOMCAR option to request the analysis to treat the respondents as a domain rather than exclude the nonrespondents.

The following SAS statements produce the desired analysis:

```

title 'Analysis of Ice Cream Spending';
proc surveymeans data=IceCream total=StudentTotals nomcar mean sum;
  strata Grade;
  var Spending;
  weight Weight;
run;

```

Output 86.4.1 summarizes the analysis including the variance estimation method.

Output 86.4.1 Analysis of Incomplete Ice Cream Data Excluding Observations with Missing Values

Analysis of Ice Cream Spending	
The SURVEYMEANS Procedure	
Data Summary	
Number of Strata	3
Number of Observations	40
Sum of Weights	4000
Variance Estimation	
Method	Taylor Series
Missing Values	NOMCAR

Output 86.4.2 shows the mean and total estimates when treating respondents as a domain in the student population. Although the point estimates are the same as the analysis without the NOMCAR option, for this particular example, the variance estimations are slightly higher when you assume that the missingness is not completely at random.

Output 86.4.2 Analysis of Incomplete Ice Cream Data Excluding Observations with Missing Values

Variable	Statistics			
	Mean	Std Error of Mean	Sum	Std Dev
Spending	9.770542	0.652347	32139	3515.126876

Example 86.5: Variance Estimation Using Replication Methods

In order to improve service, the San Francisco Municipal Railway (MUNI) conducts a survey to estimate passenger's average waiting time for MUNI's subway system.

The study uses a stratified cluster sample design. Each MUNI subway line is a stratum. The subway lines included in the study are 'J-Church,' 'K-Ingleside,' 'L-Taraval,' 'M-Ocean View,' 'N-Judah,' and the street car 'F-Market & Wharves.' The MUNI vehicles in service for these lines during a day are primary sampling units. Within each stratum, two vehicles (PSUs) are randomly selected. Then the waiting times of passengers for a selected MUNI vehicle are collected.

Table 86.5 shows the number of passengers that are interviewed in each of the selected MUNI vehicles.

Table 86.5 The Sample of the MUNI Waiting Time Study

MUNI Line	Vehicle	Number of Passengers
F-Market & Wharves	1	65
	2	102
J-Church	1	101
	2	142
K-Ingleside	1	145
	2	180
L-Taraval	1	135
	2	185
M-Ocean View	1	139
	2	203
N-Judah	1	306
	2	234

The collected data are saved in the SAS data set MUNIsurvey. The variable Line indicates which MUNI line a passenger is riding. The variable vehicle identifies the vehicle that a passenger is boarding. The variable Waittime is the time (in minutes) that a passenger waited. The variable weight contains the sampling weights, which are determined by selection probabilities within each stratum.

Output 86.5.1 displays the first 10 observations of the data set MUNIsurvey.

Output 86.5.1 First 10 Observations in the Data Set from the MUNI Subway Survey

MUNI Subway Passenger Waiting Time Survey Data					
Obs	line	vehicle	passenger	waittime	weight
1	F-Market & Wharves	1	1	18	59
2	F-Market & Wharves	1	2	0	59
3	F-Market & Wharves	1	3	16	59
4	F-Market & Wharves	1	4	13	59
5	F-Market & Wharves	1	5	5	59
6	F-Market & Wharves	1	6	13	59
7	F-Market & Wharves	1	7	7	59
8	F-Market & Wharves	1	8	5	59
9	F-Market & Wharves	1	9	16	59
10	F-Market & Wharves	1	10	5	59

Using the VARMETHOD=BRR option, the following SAS statements analyze the MUNI subway survey by using the BRR method to estimate the variance:

```

title 'MUNI Passenger Waiting Time Analysis Using BRR';
proc surveymeans data=MUNIsurvey mean varmethod=brr mean clm;
  strata line;
  cluster vehicle;
  var waittime;
  weight weight;
run;

```

The STRATUM variable is line, which corresponds to MUNI lines. The two clusters within each stratum are identified by the variable vehicle. The sampling weights are stored in the variable weight. The mean and confident limits of passenger waiting time (in minutes) are requested statistics.

Output 86.5.2 summarizes the data and indicates that the variance estimation method is BRR with 8 replicates.

Output 86.5.2 MUNI Passenger Waiting Time Analysis Using the BRR Method

MUNI Passenger Waiting Time Analysis Using BRR	
The SURVEYMEANS Procedure	
Data Summary	
Number of Strata	6
Number of Clusters	12
Number of Observations	1937
Sum of Weights	143040
Variance Estimation	
Method	BRR
Number of Replicates	8

Output 86.5.3 reports that the average passenger waiting time for a MUNI vehicle is 7.33 minutes, with an estimated standard of 0.24 minutes, using the BRR method. The 95% confident limits for the mean are estimated as 6.75 to 7.91 minutes.

Output 86.5.3 MUNI Passenger Waiting Time Analysis Using the BRR Method

Statistics			
Variable	Mean	Std Error of Mean	95% CL for Mean
waittime	7.333012	0.237557	6.75172983 7.91429366

Alternatively, the variance can be estimated using the jackknife method if the VARMETHOD=JACKKNIFE option is used. The following SAS statements analyze the MUNI subway survey by using the jackknife method to estimate the variance:

```

title 'MUNI Passenger Waiting Time Analysis Using Jackknife';
proc surveymeans data=MUNIsurvey mean varmethod=jackknife mean clm;
  strata line;
  cluster vehicle;
  var waittime;
  weight weight;
run;

```

Output 86.5.4 summarizes the data and indicates that the variance estimation method is jackknife with 12 replicates.

Output 86.5.4 MUNI Passenger Waiting Time Analysis Using the Jackknife Method

MUNI Passenger Waiting Time Analysis Using Jackknife	
The SURVEYMEANS Procedure	
Data Summary	
Number of Strata	6
Number of Clusters	12
Number of Observations	1937
Sum of Weights	143040
Variance Estimation	
Method	Jackknife
Number of Replicates	12

Output 86.5.5 reports the statistics computed using the jackknife method. Although the average passenger waiting time remains the same (7.33 minutes), the standard error is slightly smaller 0.23 minutes when the jackknife method is used, as opposed to 0.24 minutes when the BRR method is used. The 95% confidence limits are between 6.76 and 7.90 minutes when the jackknife method is used.

Output 86.5.5 MUNI Passenger Waiting Time Analysis Using the Jackknife Method

Statistics			
Variable	Mean	Std Error of Mean	95% CL for Mean
waittime	7.333012	0.232211	6.76481105 7.90121244

References

- Brick, J. M. and Kalton, G. (1996), "Handling Missing Data in Survey Research," *Statistical Methods in Medical Research*, 5, 215–238.
- Cochran, W. G. (1977), *Sampling Techniques*, Third Edition, New York: John Wiley & Sons.
- Dippo, C. S., Fay, R. E., and Morganstein, D. H. (1984), "Computing Variances from Complex Samples with Replicate Weights," *Proceedings of the Survey Research Methods Section, ASA*, 489–494.

- Dorfman, A. and Valliant, R. (1993), "Quantile Variance Estimators in Complex Surveys," *Proceedings of the Survey Research Methods Section, ASA*, 866–871.
- Fay, R. E. (1984), "Some Properties of Estimators of Variance Based on Replication Methods," *Proceedings of the Survey Research Methods Section, ASA*, 495–500.
- Fay, R. E. (1989), "Theory and Application of Replicate Weighting for Variance Calculations," *Proceedings of the Survey Research Methods Section, ASA*, 212–217.
- Francisco, C. A. and Fuller, W. A. (1991), "Quantile Estimation with a Complex Survey Design," *Annals of Statistics*, 19, 454–469.
- Fuller, W. A. (1975), "Regression Analysis for Sample Survey," *Sankhyā*, 37, Series C, Pt. 3, 117–132.
- Fuller, W. A., Kennedy, W., Schnell, D., Sullivan, G., and Park, H. J. (1989), *PC CARP*, Ames: Statistical Laboratory, Iowa State University.
- Hansen, M. H., Hurwitz, W. N., and Madow, W. G. (1953), *Sample Survey Methods and Theory*, Volumes I and II, New York: John Wiley & Sons.
- Hidiroglou, M. A., Fuller, W. A., and Hickman, R. D. (1980), *SUPER CARP*, Ames, IA: Statistical Laboratory, Iowa State University.
- Judkins, D. (1990), "Fay's Method for Variance Estimation," *Journal of Official Statistics*, 6, 223–239.
- Kalton, G. (1983), *Introduction to Survey Sampling*, Sage University Paper series on Quantitative Applications in the Social Sciences, series no. 07-035, Beverly Hills, CA, and London: Sage Publications.
- Kalton, G., and Kaspyzyk, D. (1986), "The Treatment of Missing Survey Data," *Survey Methodology*, 12, 1–16.
- Kish, L. (1965), *Survey Sampling*, New York: John Wiley & Sons.
- Lee, E. S., Forthoffer, R. N., and Lorimor, R. J. (1989), *Analyzing Complex Survey Data*, Sage University Paper series on Quantitative Applications in the Social Sciences, series no. 07-071, Beverly Hills, CA, and London: Sage Publications.
- Lohr, S. L. (2009), *Sampling: Design and Analysis*, Second Edition, Pacific Grove, CA: Duxbury Press.
- Rao, J. N. K. and Shao, J. (1996), "On Balanced Half Sample Variance Estimation in Stratified Sampling," *Journal of the American Statistical Association*, 91, 343–348.
- Rao, J. N. K. and Shao, J. (1999), "Modified Balanced Repeated Replication for Complex Survey Data," *Biometrika*, 86, 403–415.
- Rao, J. N. K., Wu, C. F. J., and Yue, K. (1992), "Some Recent Work on Resampling Methods for Complex Surveys," *Survey Methodology*, 18, 209–217.

- Rust, K. (1985), "Variance Estimation for Complex Estimators in Sample Surveys," *Journal of Official Statistics*, 1, 381–397.
- Särndal, C. E., Swensson, B., and Wretman, J. (1992), *Model Assisted Survey Sampling*, New York: Springer-Verlag.
- Wolter, K. M. (1985), *Introduction to Variance Estimation*, New York: Springer-Verlag.
- Woodruff, R. S. (1971), "A Simple Method for Approximating the Variance of a Complicated Estimate," *Journal of the American Statistical Association*, 66, 411–414.

Subject Index

- alpha level
 - SURVEYMEANS procedure, [7248](#)
- balanced repeated replication
 - SURVEYMEANS procedure, [7280](#)
 - variance estimation (SURVEYMEANS), [7280](#)
- BRR
 - SURVEYMEANS procedure, [7280](#), [7304](#)
- BRR variance estimation
 - SURVEYMEANS procedure, [7280](#)
- categorical variable
 - SURVEYMEANS procedure, [7269](#)
- classification variable
 - SURVEYMEANS procedure, [7258](#), [7269](#), [7273](#)
- clustering
 - SURVEYMEANS procedure, [7258](#)
- coefficient of variation
 - SURVEYMEANS procedure, [7273](#)
- confidence level
 - SURVEYMEANS procedure, [7248](#)
- confidence limits
 - SURVEYMEANS procedure, [7272](#), [7274](#)
- degrees of freedom
 - SURVEYMEANS procedure, [7271](#)
- domain analysis
 - SURVEYMEANS procedure, [7267](#)
- domain statistics
 - SURVEYMEANS procedure, [7276](#)
- donor stratum
 - SURVEYMEANS procedure, [7282](#)
- Fay coefficient
 - SURVEYMEANS procedure, [7254](#), [7282](#)
- Fay's BRR method
 - variance estimation (SURVEYMEANS), [7282](#)
- finite population correction
 - SURVEYMEANS procedure, [7250](#), [7251](#), [7266](#)
- Hadamard matrix
 - SURVEYMEANS procedure, [7254](#), [7284](#)
- jackknife
 - jackknife coefficients
 - SURVEYMEANS procedure, [7282](#), [7285](#)
 - jackknife variance estimation
 - SURVEYMEANS procedure, [7282](#)
 - mean per element
 - SURVEYMEANS procedure, [7269](#)
 - means
 - SURVEYMEANS procedure, [7269](#)
 - missing values
 - SURVEYMEANS procedure, [7248](#), [7265](#), [7302](#)
 - number of replicates
 - SURVEYMEANS procedure, [7256](#), [7280](#), [7282](#)
 - output data sets
 - SURVEYMEANS procedure, [7284](#)
 - output jackknife coefficient
 - SURVEYMEANS procedure, [7285](#)
 - output replicate weights
 - SURVEYMEANS procedure, [7284](#)
 - output table names
 - SURVEYMEANS procedure, [7293](#)
 - percentile
 - SURVEYMEANS procedure, [7278](#)
 - percentiles
 - SURVEYMEANS procedure, [7278](#)
 - primary sampling units (PSUs)
 - SURVEYMEANS procedure, [7267](#)
 - proportion estimation
 - SURVEYMEANS procedure, [7273](#)
 - quantile
 - SURVEYMEANS procedure, [7278](#)
 - quantiles
 - SURVEYMEANS procedure, [7278](#)
 - ratio analysis
 - SURVEYMEANS procedure, [7260](#), [7275](#)
 - ratios
 - SURVEYMEANS procedure, [7260](#), [7275](#)
 - rectangular table
 - SURVEYMEANS procedure, [7250](#), [7285](#)
 - replication methods
 - SURVEYMEANS procedure, [7253](#), [7280](#), [7304](#)
- jackknife
 - SURVEYMEANS procedure, [7280](#), [7282](#), [7304](#)

- sampling rates
 - SURVEYMEANS procedure, 7250, 7266
- sampling weights
 - SURVEYMEANS procedure, 7262, 7264
- simple random sampling
 - SURVEYMEANS procedure, 7241
- stacking table
 - SURVEYMEANS procedure, 7250, 7285
- standard deviations
 - SURVEYMEANS procedure, 7273
- standard errors
 - SURVEYMEANS procedure, 7270
- statistic-keywords
 - SURVEYMEANS procedure, 7251
- statistical computations
 - SURVEYMEANS procedure, 7267
- stratification
 - SURVEYMEANS procedure, 7263
- stratified cluster sample
 - SURVEYMEANS procedure, 7294
- stratified sampling
 - SURVEYMEANS procedure, 7243
- subdomain analysis, *see also* domain analysis
- subgroup analysis, *see also* domain analysis
- subpopulation analysis, *see also* domain analysis
- survey sampling
 - descriptive statistics, 7240
- SURVEYMEANS procedure, 7240
 - alpha level, 7248
 - balanced repeated replication, 7280
 - BRR, 7280, 7304
 - BRR variance estimation, 7280
 - categorical variable, 7258, 7269, 7273
 - class level information table, 7287
 - classification variable, 7269
 - clustering, 7258
 - coefficient of variation, 7273
 - confidence level, 7248
 - confidence limits, 7272, 7274
 - data and sample design summary table, 7287
 - degrees of freedom, 7271
 - denominator variable, 7260
 - domain analysis, 7267
 - domain analysis table, 7291
 - domain means, 7276
 - domain ratio, 7278
 - domain ratio analysis table, 7292
 - domain statistics, 7276
 - domain totals, 7277
 - domain variable, 7259
 - donor stratum, 7282
 - estimated frequencies, 7273
 - estimated totals, 7273
 - Fay coefficient, 7254, 7282
 - Fay's BRR variance estimation, 7282
 - finite population correction, 7250, 7251, 7266
 - first-stage sampling rate, 7250
 - Hadamard matrix, 7254, 7284, 7292
 - jackknife, 7280, 7282, 7304
 - jackknife coefficients, 7282, 7285
 - jackknife variance estimation, 7282
 - list of strata, 7263
 - mean per element, 7269
 - means, 7269
 - missing values, 7248, 7265, 7302
 - number of replicates, 7256, 7280, 7282
 - numerator variable, 7260
 - ODS table names, 7293
 - output data sets, 7246, 7284
 - output jackknife coefficient, 7285
 - output replicate weights, 7284
 - output table names, 7293
 - percentile, 7278
 - percentiles, 7278
 - population totals, 7251, 7266
 - primary sampling units (PSUs), 7267
 - proportion estimation, 7273
 - quantile, 7278
 - quantiles, 7278
 - quantiles table, 7290
 - ratio analysis, 7260, 7275
 - ratio analysis table, 7291
 - ratios, 7260, 7275
 - rectangular table, 7250, 7285
 - replication methods, 7253, 7280, 7304
 - sampling rates, 7250, 7266
 - sampling weights, 7262, 7264
 - simple random sampling, 7241
 - stacking table, 7250, 7285
 - standard deviations of totals, 7273
 - standard errors, 7270
 - standard errors of means, 7270
 - standard errors of ratios, 7275
 - statistic-keywords, 7251
 - statistical computations, 7267
 - statistics table, 7289
 - stratification, 7263
 - stratified cluster sample, 7294
 - stratified sampling, 7243
 - stratum information table, 7288
 - t* test, 7271
 - Taylor series variance estimation, 7257, 7270, 7271, 7274
 - valid observation, 7287
 - variance estimation, 7267
 - variance estimation table, 7288
 - variances of means, 7270

- variances of totals, [7273](#)
- VARMETHOD=BRR option, [7280](#)
- VARMETHOD=JACKKNIFE option, [7282](#)
- VARMETHOD=JK option, [7282](#)
- weighting, [7262](#), [7264](#)

t test

- SURVEYMEANS procedure, [7271](#)

Taylor series variance estimation

- SURVEYMEANS procedure, [7257](#), [7270](#),
[7271](#), [7274](#)

variance estimation

- BRR (SURVEYMEANS), [7280](#)
- jackknife (SURVEYMEANS), [7282](#)
- SURVEYMEANS procedure, [7267](#)
- Taylor series (SURVEYMEANS), [7257](#)
- Taylor series (SURVEYMEANS), [7270](#),
[7271](#), [7274](#)

variances of totals

- SURVEYMEANS procedure, [7273](#)

VARMETHOD=BRR option

- SURVEYMEANS procedure, [7280](#)

VARMETHOD=JACKKNIFE option

- SURVEYMEANS procedure, [7282](#)

VARMETHOD=JK option

- SURVEYMEANS procedure, [7282](#)

weighting

- SURVEYMEANS procedure, [7262](#), [7264](#)

Syntax Index

ALPHA= option
PROC SURVEYMEANS statement, [7248](#)

BY statement
SURVEYMEANS procedure, [7257](#)

CLASS statement
SURVEYMEANS procedure, [7258](#)

CLUSTER statement
SURVEYMEANS procedure, [7258](#)

DATA= option
PROC SURVEYMEANS statement, [7248](#)

DF= option
REPWEIGHTS statement
(SURVEYMEANS), [7262](#)

DFADJ option
DOMAIN statement (SURVEYMEANS),
[7260](#)
VARMETHOD=BRR (PROC
SURVEYMEANS statement), [7254](#)
VARMETHOD=JACKKNIFE (PROC
SURVEYMEANS statement), [7256](#)
VARMETHOD=JK (PROC
SURVEYMEANS statement), [7256](#)

DOMAIN statement
SURVEYMEANS procedure, [7259](#)

FAY= option
VARMETHOD=BRR (PROC
SURVEYMEANS statement), [7254](#)

H= option
VARMETHOD=BRR (PROC
SURVEYMEANS statement), [7254](#)

HADAMARD= option
VARMETHOD=BRR (PROC
SURVEYMEANS statement), [7254](#)

JKCOEFS= option
REPWEIGHTS statement
(SURVEYMEANS), [7262](#)

LIST option
STRATA statement (SURVEYMEANS),
[7263](#)

MISSING option
PROC SURVEYMEANS statement, [7248](#)

N= option
PROC SURVEYMEANS statement, [7251](#)

NOMCAR option
PROC SURVEYMEANS statement, [7248](#)

NONSYMCL option
PROC SURVEYMEANS statement, [7249](#)

NOSPARSE option
PROC SURVEYMEANS statement, [7249](#)

ORDER= option
PROC SURVEYMEANS statement, [7249](#)

OUTJKCOEFS= option
VARMETHOD=JACKKNIFE (PROC
SURVEYMEANS statement), [7257](#)
VARMETHOD=JK (PROC
SURVEYMEANS statement), [7257](#)

OUTWEIGHTS= option
VARMETHOD=BRR (PROC
SURVEYMEANS statement), [7255](#)
VARMETHOD=JACKKNIFE (PROC
SURVEYMEANS statement), [7257](#)
VARMETHOD=JK (PROC
SURVEYMEANS statement), [7257](#)

PERCENTILE= option
PROC SURVEYMEANS statement, [7249](#)

PRINTH option
VARMETHOD=BRR (PROC
SURVEYMEANS statement), [7255](#)
PROC SURVEYMEANS statement, *see*
SURVEYMEANS procedure

QUANTILE= option
PROC SURVEYMEANS statement, [7249](#)

R= option
PROC SURVEYMEANS statement, [7250](#)

RATE= option
PROC SURVEYMEANS statement, [7250](#)

RATIO statement
SURVEYMEANS procedure, [7260](#)

REPS= option
VARMETHOD=BRR (PROC
SURVEYMEANS statement), [7256](#)

REPWEIGHTS statement
SURVEYMEANS procedure, [7262](#)

STACKING option
PROC SURVEYMEANS statement, [7250](#)

- STRATA statement
 - SURVEYMEANS procedure, [7263](#)
- SUBGROUP statement
 - SURVEYMEANS procedure, [7259](#)
- SURVEYMEANS procedure
 - syntax, [7247](#)
- SURVEYMEANS procedure, BY statement, [7257](#)
- SURVEYMEANS procedure, CLASS statement, [7258](#)
- SURVEYMEANS procedure, CLUSTER statement, [7258](#)
- SURVEYMEANS procedure, DOMAIN statement, [7259](#)
 - DFADJ option, [7260](#)
- SURVEYMEANS procedure, PROC SURVEYMEANS statement, [7248](#)
 - ALPHA= option, [7248](#)
 - DATA= option, [7248](#)
 - DFADJ option (VARMETHOD=BRR), [7254](#)
 - DFADJ option (VARMETHOD=JACKKNIFE), [7256](#)
 - DFADJ option (VARMETHOD=JK), [7256](#)
 - FAY= option (VARMETHOD=BRR), [7254](#)
 - H= option (VARMETHOD=BRR), [7254](#)
 - HADAMARD= option (VARMETHOD=BRR), [7254](#)
 - MISSING option, [7248](#)
 - N= option, [7251](#)
 - NOMCAR option, [7248](#)
 - NONSYMCL option, [7249](#)
 - NOSPARSE option, [7249](#)
 - ORDER= option, [7249](#)
 - OUTJKCOEFS= option (VARMETHOD=JACKKNIFE), [7257](#)
 - OUTJKCOEFS= option (VARMETHOD=JK), [7257](#)
 - OUTWEIGHTS= option (VARMETHOD=BRR), [7255](#)
 - OUTWEIGHTS= option (VARMETHOD=JACKKNIFE), [7257](#)
 - OUTWEIGHTS= option (VARMETHOD=JK), [7257](#)
 - PERCENTILE= option, [7249](#)
 - PRINTH option (VARMETHOD=BRR), [7255](#)
 - QUANTILE= option, [7249](#)
 - R= option, [7250](#)
 - RATE= option, [7250](#)
 - REPS= option (VARMETHOD=BRR), [7256](#)
 - STACKING option, [7250](#)
 - TOTAL= option, [7251](#)
 - VARMETHOD= option, [7253](#)
- SURVEYMEANS procedure, RATIO statement, [7260](#)
- SURVEYMEANS procedure, REPWEIGHTS statement, [7262](#)
 - DF= option, [7262](#)
 - JKCOEFS= option, [7262](#)
- SURVEYMEANS procedure, STRATA statement, [7263](#)
 - LIST option, [7263](#)
- SURVEYMEANS procedure, VAR statement, [7264](#)
- SURVEYMEANS procedure, WEIGHT statement, [7264](#)
- TOTAL= option
 - PROC SURVEYMEANS statement, [7251](#)
- VAR statement
 - SURVEYMEANS procedure, [7264](#)
- VARMETHOD= option
 - PROC SURVEYMEANS statement, [7253](#)
- WEIGHT statement
 - SURVEYMEANS procedure, [7264](#)

Your Turn

We welcome your feedback.

- If you have comments about this book, please send them to **`yourturn@sas.com`**. Include the full title and page numbers (if applicable).
- If you have comments about the software, please send them to **`suggest@sas.com`**.

SAS® Publishing Delivers!

Whether you are new to the work force or an experienced professional, you need to distinguish yourself in this rapidly changing and competitive job market. SAS® Publishing provides you with a wide range of resources to help you set yourself apart. Visit us online at support.sas.com/bookstore.

SAS® Press

Need to learn the basics? Struggling with a programming problem? You'll find the expert answers that you need in example-rich books from SAS Press. Written by experienced SAS professionals from around the world, SAS Press books deliver real-world insights on a broad range of topics for all skill levels.

support.sas.com/saspress

SAS® Documentation

To successfully implement applications using SAS software, companies in every industry and on every continent all turn to the one source for accurate, timely, and reliable information: SAS documentation. We currently produce the following types of reference documentation to improve your work experience:

- Online help that is built into the software.
- Tutorials that are integrated into the product.
- Reference documentation delivered in HTML and PDF – **free** on the Web.
- Hard-copy books.

support.sas.com/publishing

SAS® Publishing News

Subscribe to SAS Publishing News to receive up-to-date information about all new SAS titles, author podcasts, and new Web site features via e-mail. Complete instructions on how to subscribe, as well as access to past issues, are available at our Web site.

support.sas.com/spn



**THE
POWER
TO KNOW®**

